

GRADED RINGS OF RANK 2 SARKISOV LINKS

GAVIN BROWN AND FRANCESCO ZUCCONI

ABSTRACT. We compute a class of Sarkisov links from Fano 3-folds embedded in weighted Grassmannians using explicit methods for describing graded rings associated to a variation of GIT quotient.

1. INTRODUCTION

The main aim of this paper is to describe a class of Sarkisov links from Fano 3-folds embedded in weighted Grassmannians $w\text{Grass}(2, 5)$. Altınok [Alt98] lists 69 families of such Fano 3-folds. These are listed in [BDK⁺], and they are discussed from the point of view of weighted Grassmannians in [CR02]. The general member of each family lies in weighted projective space (wps) as a codimension 3 quasismooth variety defined by the five maximal Pfaffians of a skew 5×5 matrix.

Theorem 1.1 gives a concise statement of the result; the proof provides a great deal of additional intricate information about the links, as described in Section 4.3.

We make a detailed study of links from Type I centres on these Fano 3-folds. (A Type I centre is a cyclic quotient singularity $\frac{1}{r}(n_1, n_2, n_3)$ on X for which the weights n_1, n_2, n_3 of the $\mathbb{Z}/r\mathbb{Z}$ action are those of independent global variables; see Definition 4.1 below. The Kawamata blowup is then simply the weighted blowup with weights n_1, n_2, n_3 .)

Theorem 1.1. *Let X be a general member of one of the 69 families of codimension 3 Pfaffian Fano 3-folds and $p \in X$ be a Type I centre. Suppose that the three quasilinear equations defining the orbifold tangent space at p have weights $a_1 \leq a_2 \leq a_3$. Then there is a Sarkisov link that starts with the Kawamata blowup of p followed by a flop:*

$$X \longleftarrow Y \dashrightarrow Y_1.$$

The link is completed in one of four ways:

$a_1 = a_2 = a_3$: $Y_1 = Y' \rightarrow \mathbb{P}^2$ is a conic bundle;

$a_1 < a_2 = a_3$: $Y_1 = Y' \rightarrow X'$ is a divisorial contraction to a line on a Fano 3-fold X' ;

$a_1 = a_2 < a_3$: $Y_1 \dashrightarrow Y' \rightarrow \mathbb{P}^1$ is a Mori flip followed by a del Pezzo fibration;

$a_1 < a_2 < a_3$: $Y_1 \dashrightarrow Y' \rightarrow X'$ is a Mori flip followed by a divisorial contraction to a point on a Fano 3-fold X' .

2000 *Mathematics Subject Classification.* 14J30 (Primary); 14J45, 14E30, 14E08 (Secondary).

Corresponding author: Gavin Brown, IMSAS, University of Kent, Canterbury, CT2 7NF, UK.

Fax: +44 (0)1227 827932. Email: gdb@kent.ac.uk

The first author was partially supported by EPSRC grants GR/S03980/01 and EP/E000258/1 and ‘MIUR PRIN 2003: Spazi di moduli e teoria di Lie’.

Francesco Zucconi, DIMI, Università di Udine, via delle Scienze 208, 33100 Udine, Italy. Email: zucconi@dimi.uniud.it.

More detailed information about each of the four cases is given in Section 4.3.4, including a list of conic bundles that occur and, in Table 1, a list of del Pezzo fibrations that occur.

Sarkisov links are a tool to study the birational rigidity or otherwise of Fano 3-folds and Mori fibre spaces. In every link arising in Theorem 1.1, the 3-fold at the end of the link (either X' or Y' , depending on the link) is not isomorphic to 3-fold X at the beginning. Thus we have:

Corollary 1.2. *Let X be a general member of one of the 64 families of codimension 3 Pfaffian Fano 3-folds that admit a Type I centre. Then X is not birationally rigid.*

The remaining five families of 5×5 Pfaffian Fano 3-folds do not have a Type I centre (two of them have no projections at all, the rest have more complicated projections that will need a new analysis starting with [Rei02] and generalisations of [Pap06]) and so we cannot find the links of Theorem 1.1 in those cases.

1.1. Explicit Sarkisov links in the literature. Theorem 1.1 is a descendent of other explicit calculations in the literature. We describe some of these; we do not discuss work, such as that of Grinenko [Gri01] or Abramovich et al [AKMW02], which is very relevant but proceeds by other methods.

For general weighted hypersurfaces $X \subset \mathbb{P}^4$, the full analysis of Sarkisov links from X was completed in [CPR00]. It is shown there that all such links are birational involutions and therefore that a general Fano hypersurface is birationally rigid. In this context, Type I centres are those centres which lead to quadratic involutions (Q.I.). Hypersurfaces also admit elliptic involutions (E.I.) that one might naturally define as Type II₁ centres (see Reid [Rei02] and Papadakis [Pap06]).

In codimension 2, Corti–Mella [CM04] work out examples of links from $X \subset \mathbb{P}^5$ in detail and explain how to carry out calculations of links on any example. The main part of [CM04] is devoted to the ‘exclusion’ methods that show that particular Fano 3-folds have no other links. The following result is implicit in [CM04], and we phrase it as the prototype for Theorem 1.1.

Theorem 1.3 (Corti–Mella [CM04]). *Let $X \subset \mathbb{P}^5$ be a codimension 2 Fano 3-fold and $p \in X$ be a Type I centre. Suppose that the equations defining the tangent space at p have weights $a_1 \leq a_2$. Then there is a Sarkisov link that starts with the Kawamata blowup of p followed by a flop:*

$$X \leftarrow Y \dashrightarrow Y'.$$

The link continues in one of two ways:

$$\begin{aligned} a_1 = a_2: & \quad Y' \rightarrow \mathbb{P}^1 \text{ is a del Pezzo fibration;} \\ a_1 < a_2: & \quad Y' \rightarrow X' \text{ is a divisorial contraction to a point on a Fano} \\ & \quad \text{3-fold } X'. \end{aligned}$$

In the second case, the divisor contracts to a singular point p' on a Fano 3-fold X' and the index of $p' \in X'$ is $a_2 - a_1$.

1.2. Graded rings of blowups. We construct links using variation of GIT. These methods were first discussed by Dolgachev and Hu [DH98], Reid [Rei92] and Thaddeus [Tha96]. They were used to construct Mori flips as toric hypersurfaces in [Bro99], and some of those flips appear in the links of Theorem 1.1. Hu and Keel’s more recent notion of *Mori dream space* [HK00] is very close to our needs here. The difference is that our computations are

explicit: as discussed by Corti and Reid in the introduction to [CR00], we seek descriptions of every aspect of these links by equations or something equally concrete. We work directly with graded rings associated to the problem. There are many rings one could define, and during calculations there are two main considerations: we can only work with rings that are reasonably small, and we cannot always be certain that we have computed the whole ring under consideration. So, while the theory of [HK00] is comprehensive, we need other detailed information to make our calculations.

Let X be a Fano 3-fold and $p \in X$. For an extremal contraction $f: Y \rightarrow X$ contracting a divisor $E \subset Y$ to a point $p \in X$, we follow Cox [Cox95], Hu and Keel [HK00], Kawakita [Kaw02] and others and describe a graded ring

$$\mathcal{M}(f) = \bigoplus_{n \in \mathbb{Z}} \bigoplus_{m \geq 0} H^0(Y, -(mK_Y + nE)) \quad (\text{see Definition 2.6}).$$

Theorem 1.4. *Let f be an extremal extraction from a point $p \in X$. If $\mathcal{M}(f)$ is finitely generated as a \mathbb{C} -algebra, then there is a 2-ray game (see section 2.1 below) Φ from $p \in X$ starting with f .*

Of course, the finite generation condition on \mathcal{M} is the essential point. To check it is tantamount to describing the link, and we do this for certain links from anticanonically embedded Fano 3-folds in codimension 3. We restate and prove this theorem as Theorem 5.1 below.

This method describes other phenomena, such as Ryder’s elliptic and K3 fibrations [Ryd06] and the ‘bad’ links of Corti et al. [CPR00]. The methods we describe are also used in [BCZ04] to compute links from Mori fibre spaces and in [CS05] to compute birational modifications.

Throughout this paper we work over the field $k = \mathbb{C}$.

Acknowledgments. This paper was initiated during the HDG meeting of the Newton Institute, Cambridge, 2002, and with visits of the first author to Udine in 2003 and of the second author to DPMMS, Cambridge 2002. We thank Professors A. Corti, M. Gross, M. Reid and H. Takagi for their mathematical support during that time.

2. RANK 2 LINKS

In the spirit of [CPR00], we consider explicit Fano 3-folds X embedded in weighted projective space (wps) $X \subset \mathbb{P}^n$ for small values of n : we denote the weights of the wps \mathbb{P}^n by $\mathbb{P}^n(a_0, \dots, a_n)$ when we know them, but even without this indication the symbol \mathbb{P}^n always denotes some wps. By definition, a Fano 3-fold has minimal Picard rank, $\rho_X = 1$. There are well-known lists of such Fanos: Reid’s famous 95 hypersurfaces, Iano-Fletcher’s 85 families in codimension 2 [IF00], and Altınok’s lists of 70 and 142 Fanos in codimensions 3 and 4 [Alt98]. These lists, and others in higher codimension, are in [BDK⁺].

We sometimes assume that X is quasismooth, that is, that the weighted affine cone on X is singular only at the origin. Quasi-smooth varieties in wps have only quotient singularities.

2.1. Links and Mori fibre spaces. This subject is already well discussed—by Corti [Cor00] for instance—so we are brief. We recall that a 3-fold X is \mathbb{Q} -factorial if every Weil divisor on X admits some multiple that is a Cartier divisor. When this condition fails for a divisor D , there is a projective morphism $Y \rightarrow X$ that is an isomorphism in codimension 1 so that the strict transform of D is a (relatively ample) Cartier divisor; see [Kaw88], Lemma 3.1, for more detailed explanation.

2.1.1. *The 2-ray game and rank 2 links.* The key observation is that if a projective and \mathbb{Q} -factorial variety Y has Picard rank $\rho(Y) = 2$, then up to isomorphism it admits at most two projective morphisms, each of relative Picard rank 1. There are two instant sources of such varieties: fibrations $Y \rightarrow S$ and blowups $Y \rightarrow X$, where in each case the base is assumed \mathbb{Q} -factorial with $\rho = 1$ and the map is extremal in the sense that $\rho(Y/S) = 1$ or $\rho(Y/X) = 1$. Each of these descriptions of Y accounts for one of the two possible extremal morphisms that could exist from Y . If Y admits a second morphism, $Y \rightarrow Z$ say, then one of two things may happen. If Z is \mathbb{Q} -factorial, then the game is over. On the other hand, if Z is not \mathbb{Q} -factorial, then the morphism is necessarily small and one looks to make a flip

$$Y \rightarrow Z \leftarrow Y_1,$$

meaning that the rational map $Y \dashrightarrow Y_1$ is an isomorphism in codimension 1 but is not an isomorphism, and that the flipped variety Y_1 is again \mathbb{Q} -factorial with $\rho = 2$. And so the 2-ray game proceeds: we switch attention to Y_1 , look for its second morphism $Y_1 \rightarrow Z_2$ —again one of the two possible morphisms from Y_1 is already accounted for—and terminate or flip as required. From either of our starting configurations, there is never a choice to be made as long as the necessary morphisms and flips exist. However, the game can break down either by encountering a Y_i with only one morphism or a flipping contraction $Y_i \rightarrow Z_{i+1}$ that does not flip.

The expression *2-ray game* refers to this procedure, whether or not it breaks down. Successful games are called 2-ray links.

Definition 2.1. A *2-ray link* is a diagram

$$(2.1) \quad \begin{array}{ccccccc} & & Y = Y_0 & \dashrightarrow & Y_1 & \dashrightarrow & \dots & \dashrightarrow & Y_k = Y' & & \\ & \swarrow & & \searrow & \swarrow & & & & \searrow & & \\ X & & & & Z_1 & & & & & & X' \end{array}.$$

of projective varieties and projective morphisms, in which each Y_i is \mathbb{Q} -factorial with $\rho(Y_i) = 2$, X and X' are \mathbb{Q} -factorial with $\rho = 1$, and each Z_i has $\rho(Z_i) = 1$ but is not \mathbb{Q} -factorial. Moreover, the rational maps $Y_i \dashrightarrow Y_{i+1}$ are isomorphisms in codimension 1, but are not isomorphisms.

2.1.2. *The Sarkisov program.* Now we discuss the category in which we play a 2-ray game. There are many that are appropriate, but we consider the Mori ‘category’ of projective, \mathbb{Q} -factorial, terminal 3-folds and extremal morphisms. As is familiar, such 3-folds occur naturally in the minimal model program in 3 dimensions, even if one starts with nonsingular 3-folds; see [KM98] for a general introduction. We refer to [Rei87] for the definition of *terminal* singularities, since we do not use it directly.

Definition 2.2. (1) A morphism $f: Y \rightarrow X$ is *extremal* if $f_*\mathcal{O}_Y = \mathcal{O}_X$ and $\rho(Y/X) = 1$.

(2) An *extremal contraction* is an extremal (projective) morphism $f: Y \rightarrow X$ between projective, \mathbb{Q} -factorial varieties such that $-K_Y$ is relatively ample for f . We say that f is of *divisorial type* if $\dim X = \dim Y$ and that f is of *fibre type* if $\dim X < \dim Y$.

(3) A 2-ray link—with notation as in (2.1) of Definition 2.1—is said to *take place in the Mori category* if

- (i) each Y_i is a 3-fold with terminal singularities,
- (ii) $\dim X \leq 3$, and if $\dim X = 3$ then X has terminal singularities; and the same conditions for X' ,
- (iii) $Y \rightarrow X$ and $Y' \rightarrow X'$ are extremal contractions,
- (iv) all morphisms $Y_i \rightarrow Z_j$ (for $j = i$ or $i+1$ and $Z_j \neq X, X'$) are extremal and small.

If f is an extremal contraction of divisorial type, then necessarily f contracts an irreducible divisor $E \subset Y$ to a subvariety of codimension at least 2 in X .

Definition 2.3. A *Mori fibre space* (Mfs) is an extremal contraction $V \rightarrow S$ of fibre type with $\dim V = 3$ in which V has terminal singularities.

There are three cases of Mfs $V \rightarrow S$ defined as follows:

- if $\dim S = 0$, then V is a *Fano 3-fold*;
- if $\dim S = 1$, then V is a *del Pezzo fibration*;
- if $\dim S = 2$, then V is a *conic bundle*.

Definition 2.4. A *Sarkisov link* between two Mfs $V \rightarrow S$ and $V' \rightarrow S'$ is a 2-ray link that takes place in the Mori category in which both V and V' appear.

Of course, in Definition 2.4 the two Mfs necessarily appear at the two ends of the 2-ray link. But we can be more specific. There are four configurations—named Types I–IV in [Cor00] section 2.2—in which V and V' can appear in a 2-ray link, depending on the dimensions of S and S' :

- Type I V and V' are Fano 3-folds
- Type II V is a Fano 3-fold and $V' \rightarrow S'$ is of fibre type
- Type III $V \rightarrow S$ is of fibre type and V' is a Fano 3-fold
- Type IV $V \rightarrow S$ and $V' \rightarrow S'$ are of fibre type.

There is only one way that each type can fit into a link: comparing with the link (2.1) of Definition 2.1, for Type I, $V = X$ and $V' = X'$, for Type II, $V = X$ and $(V' \rightarrow S') = (Y' \rightarrow X')$, and so on. The links in Theorem 1.1 all start with $(V \rightarrow S)$ being a Fano 3-fold ($X \rightarrow \text{pt.}$), and the first step being a blowup $Y \rightarrow X$ of a Fano 3-fold, so they are all of Type I or II.

We have given these definitions over $\text{Spec}(k)$. They also make sense over arbitrary base scheme U , with all notions replaced by their relative counterparts: $\rho(Y) = 2$ is replaced by $\rho(Y/U) = 2$, and so on. This is done carefully in [Cor00].

2.2. Finitely-generated extremal extractions. Let X be a Fano 3-fold and $p \in X$ a point.

Definition 2.5. A morphism $f: Y \rightarrow X$ is an *extremal extraction of p* if and only if it is an extremal contraction of a divisor E on Y with centre $p \in X$ and there is a member in $|-K_Y|$ that has only Du Val singularities.

We use the notation of [CPR00]: $A = -K_X$ and $B = -K_Y = f^*A - a(K_X, E)E$. The main case we consider is when $p \in X$ is a nontrivial terminal quotient singularity, in which case it is well known that there is exactly one extremal extraction of p , the Kawamata blowup [Kaw96].

We define a global version of the algebras considered by Kawakita [Kaw02].

Definition 2.6. Let $f: Y \rightarrow X$ be an extremal extraction of p and define a k -algebra $\mathcal{B} = \bigoplus_{m \geq 0} H^0(Y, mB)$ that is graded by m . The *graded ring of f* is the \mathbb{Z} -graded \mathcal{B} -algebra

$$\mathcal{M} = \mathcal{M}(f) = \bigoplus_{n \in \mathbb{Z}} \mathcal{M}_n$$

whose n th graded piece for each integer n is the graded \mathcal{B} -module

$$\mathcal{M}_n = \bigoplus_{m \geq 0} H^0(Y, mB - nE).$$

We consider only cases for which $\mathcal{B} = \bigoplus_{m \geq 0} H^0(Y, mB)$ is a finitely-generated k -algebra graded by m . The ring \mathcal{M} is defined as a \mathcal{B} -algebra to emphasise its grading by n —the order of vanishing of functions on E , and ultimately the action responsible for flips in

the links—but it is usually regarded as a k -algebra having two gradings by m and n . In particular, references to generators of \mathcal{M} mean generators as a k -algebra unless specified otherwise. The ring structure of \mathcal{M} is determined by multiplication in $k(Y)$. For fixed values of $m, n \in \mathbb{Z}$, there are k -subalgebras

$$R(Y, mB - nE) = \bigoplus_{k \geq 0} H^0(Y, k(mB - nE)) \subset \mathcal{M},$$

and these are zero if $m < 0$ or if $m = 0$ and $n > 0$. We only consider such subrings of \mathcal{M} for m and n coprime.

Definition 2.7. An extremal extraction f is *finitely generated* if and only if $\mathcal{M}(f)$ is finitely generated (as a k -algebra).

If f is finitely generated and the 2-ray game on Y plays out to a link Φ from $p \in X$, then loosely speaking we refer to $\mathcal{M}(f)$ as *the graded ring of the link* Φ . But this is an abuse of terminology: the ring $\mathcal{M}(f)$ exists even if the link does not, and when the link does exist its graded ring, even considered only up to isomorphism, depends on which end is taken to be the start of the link. In any case $\mathcal{M}(f)$ is of finite index in the Cox ring of Y .

3. THE 2-RAY GAME FOR TORIC VARIETIES

The following result is well known: it follows from [HK00] Corollary 2.10, for instance. To get detailed information about the geometry of each step, we trace a proof with an explicit commentary. Note that there are no conditions on the singularities of \mathcal{X} or \mathcal{Y} in this theorem. And although this is entirely toric geometry, we do not use [Rei83].

Theorem 3.1. *Let \mathcal{X} be a \mathbb{Q} -factorial toric variety with $\rho_{\mathcal{X}} = 1$ and $\mathcal{Y}_0 \rightarrow \mathcal{X}$ be the contraction of a divisor corresponding to the inclusion of a new element in the 1-skeleton of the fan $\Delta_{\mathcal{X}}$. Then there is a unique sequence of projective toric maps*

$$\begin{array}{ccccccc} & & \mathcal{Y}_0 & \dashrightarrow & \mathcal{Y}_1 & \dashrightarrow & \dots & \dashrightarrow & \mathcal{Y}' & & \\ & \swarrow & & \searrow & \swarrow & & & & \searrow & & \\ \mathcal{X} & & & & \mathcal{Z}_1 & & & & & & \mathcal{X}' \end{array}.$$

At each step, the map $\mathcal{Y}_i \dashrightarrow \mathcal{Y}_{i+1}$ is an isomorphism in codimension 1. The morphism $\mathcal{Y}' \rightarrow \mathcal{X}'$ is either a divisorial contraction or a fibration to a toric variety of lower dimension.

3.1. The Cox ring and the proof of Theorem 3.1. We describe the 2-ray game for a rank 2 toric variety in terms of its homogeneous coordinate ring. Compare the explicit calculations with with Cox's description of the homogeneous coordinate ring [Cox95] and Lecture 2 of [Rei97].

3.1.1. A rank 2 torus action. Let $R_0 = \mathbb{C}[x_1, \dots, x_N]$ and $\mathcal{A} = \text{Spec } R_0$. Let $G = \mathbb{C}^* \times \mathbb{C}^*$ with coordinates λ, μ . An action of G on \mathcal{A} is determined by N characters

$$\chi_i \in C_G = \text{Hom}_{\mathbb{Z}}(G, \mathbb{C}^*), \quad \text{for } i = 1, \dots, N,$$

by defining $g^*(x_i) = \chi_i(g) \cdot x_i$ for $i = 1, \dots, N$. Of course, the isomorphism $\mathbb{Z} \oplus \mathbb{Z} \rightarrow C_G$ taking (a, b) to the character $(\lambda, \mu) \mapsto \lambda^a \mu^b$ makes this explicit: if χ_i corresponds to (a_i, b_i) , then $(\lambda, \mu) \in G$ acts on \mathcal{A} by

$$(x_1, \dots, x_n) \mapsto (\lambda^{a_1} \mu^{b_1} x_1, \dots, \lambda^{a_N} \mu^{b_N} x_N).$$

We leave this isomorphism implicit from now on and write $\chi_i = (a_i, b_i)$. Once and for all, we fix such an action: we fix a choice of N characters $\chi_i = (a_i, b_i) \in \mathbb{Z} \oplus \mathbb{Z}$, and we insist that these characters rationally span $\mathbb{Z} \oplus \mathbb{Z}$.

In order to apply GIT, we consider the trivial line bundle $\mathcal{L} \rightarrow \mathcal{A}$, with fibre coordinate t , by the embedding $R_0 \subset R = \mathbb{C}[x_1, \dots, x_N, t]$. To extend the action compatibly from \mathcal{A} to \mathcal{L} is just a matter of choosing another character $\chi \in C_G$ and defining

$$g^*(t) = \chi(g)^{-1} \cdot t.$$

As usual, we regard χ as a parameter, and we emphasise this choice in the notation by writing $g^{*\chi}$ for the action of $g \in G$ on \mathcal{L} . Again, we can be more explicit: if χ corresponds to $(a, b) \in \mathbb{Z} \oplus \mathbb{Z}$, then $(\lambda, \mu) \in G$ acts on \mathcal{L} by

$$(x_1, \dots, x_n) \mapsto (\lambda^{a_1} \mu^{b_1} x_1, \dots, \lambda^{a_N} \mu^{b_N} x_N, \lambda^{-a} \mu^{-b} t).$$

We denote the invariant ring under this action on \mathcal{L} by

$$R^{G,\chi} = \{f \in R : g^{*\chi}(f) = f\}.$$

If $f \in R^{G,\chi}$, then we define $D_f = \{P \in \mathcal{L} : f(P) \neq 0\}$. A point $(p_1, \dots, p_N) \in \mathcal{A}$ is χ -semistable if and only if there exists $f \in R^{G,\chi}$, with $f \notin R_0$, and $s \in \mathbb{C}^*$ such that $(p_1, \dots, p_N, s) \in D_f$. The set of χ -semistable points is denoted \mathcal{A}_χ^{ss} .

Finally, the GIT quotient of \mathcal{A} by G (associated to the choice of χ) is denoted $\mathcal{A} //_\chi G$ and is defined to be the categorical quotient $\mathcal{A}_\chi^{ss} // G$. As explained in [Dol03], Theorem 8.1, this quotient is locally of the form $D_f // G = \text{Spec } \mathcal{O}(D_f)^G$, where f is a G -equivariant function and D_f is the corresponding principal open affine set.

3.1.2. Projective description of the quotient. Each choice of χ determines a $\mathbb{Z} \oplus \mathbb{Z}$ grading on R as described above. It is convenient to introduce another grading on R (and its subrings) to carry line bundles on the various quotients. The new \mathbb{N} grading is by powers of t , or equivalently by setting $\deg(x_i) = 0$ for each i and $\deg(t) = 1$. We denote the piece of R in degree $m \geq 0$ by R_m , notation which is consistent with the earlier use of $R_0 \subset R$. Clearly this grading survives taking invariants: $(R^{G,\chi})_m = (R_m)^{G,\chi}$ for each degree $m \geq 0$.

We can form $\text{Proj } R$ with respect to this grading, and it can be regarded as the \mathbb{C}^* quotient

$$\pi: \text{Spec } R \setminus \text{Spec } R_0 \rightarrow \text{Proj } R.$$

Of course, $\text{Proj } R = \mathcal{A} = \text{Spec } R_0$ is affine in this case, and the map π above is just the restriction away from the zero section of the projection $\mathcal{L} \rightarrow \mathcal{A}$, which is a geometric quotient. We can do the same for subrings of R , and we denote the quotient maps corresponding to any χ by

$$\pi^{G,\chi}: \text{Spec } R^{G,\chi} \setminus \text{Spec } R_0^{G,\chi} \rightarrow \text{Proj } R^{G,\chi}.$$

The degree 0 inclusion $R^{G,\chi} \rightarrow R$ induces maps ρ below:

$$\begin{array}{ccc} \text{Spec } R \setminus \text{Spec } R_0 & \xrightarrow{\rho} & \text{Spec } R^{G,\chi} \setminus \text{Spec } R_0^{G,\chi} \\ \downarrow \pi & & \downarrow \pi^{G,\chi} \\ \mathcal{A}_\chi^{ss} \xrightarrow{j} & \text{Proj } R & \xrightarrow{\rho} \text{Proj } R^{G,\chi} \end{array}.$$

Proposition 3.2. *The map $\rho \circ j = \pi_\chi: \mathcal{A}_\chi^{ss} \rightarrow \text{Proj } R^{G,\chi}$ is a surjective morphism and it induces an isomorphism $\mathcal{A} //_\chi G \cong \text{Proj } R^{G,\chi}$.*

Proof. Both the surjectivity of π_χ and the fact that it is defined at geometric points are straightforward: at a point in $\text{Proj } R^{G,\chi}$ there is a (non-trivial, invariant) function that does not vanish there, which is the same as the condition for semistability. Indeed, π_χ is expressed in coordinates as a choice of generators for $R^{G,\chi}$ as an $R_0^{G,\chi}$ -algebra, and a point lies in \mathcal{A}_χ^{ss} if and only if some polynomial in these generators does not vanish there.


To check that π_χ is set-theoretically injective, consider $p_1, p_2 \in \mathcal{A}$ lying in G -orbits whose closures are disjoint. There is an invariant function f which vanishes on the orbit $G \cdot p_1$ and is not identically zero on $G \cdot p_2$. This function then separates the image of the points on $\text{Proj } R^{G,\chi}$.

Finally, we match scheme structures by considering local patches. An affine patch on $\text{Proj } R^{G,\chi}$ is Spec of the ring of homogeneous rational functions of total degree 0 in t which have only expressions in f in the denominator. But this ring is exactly $\mathcal{O}_{D_f}^G$, and its spectrum is an affine patch on $\mathcal{A} //_\chi G$ by construction of the quotient. \square

3.1.3. Notation: the effective cone of characters. The characters $\chi_i: G \rightarrow \mathbb{C}^*$, $(\lambda, \mu) \mapsto \lambda^{a_i} \mu^{b_i}$, $i = 1, \dots, N$ span a convex cone denoted $\Sigma_{\text{eff}} \subset C_G \otimes \mathbb{R} \cong \mathbb{R}^2$. Let $r_0, \dots, r_M \subset \Sigma_{\text{eff}}$ be the distinct rays generated by the characters $\chi_i \in \mathbb{R}^2$. Reordering if necessary, we may assume that the rays r_α are ordered clockwise from one boundary of Σ_{eff} to the other. Two characters can generate the same ray, and so we may assume, again after reordering, for each r_α there are integers $l_\alpha \leq m_\alpha$ such that

$$\chi_i \in r_\alpha \quad \text{if and only if} \quad l_\alpha \leq i \leq m_\alpha.$$

The collection of rays $\{r_0, \dots, r_M\}$ divides Σ_{eff} into M chambers and we denote by \mathcal{C}_α these open chambers; in particular $\partial \overline{\mathcal{C}_\alpha} = r_\alpha \cup r_{\alpha+1}$, $\alpha = 0, \dots, M-1$.



From now on a character χ will be identified with the corresponding $(a, b) \in \mathbb{R}^2$ by the isomorphism $C_G \otimes \mathbb{R} \cong \mathbb{R}^2$ which sends $\chi \mapsto (a, b)$. For any two rays $r, r' \subset \Sigma_{\text{eff}}$ (not necessarily among the r_i) we say that $r \leq r'$ if and only if r is contained in the convex span of r_0 and r' . We extend this partial order to elements of C_G by saying $(a, b) \leq (a', b')$ when $(a, b)\mathbb{R}_{\geq 0} \leq (a', b')\mathbb{R}_{\geq 0}$. (That is, we simply put elements of Σ_{eff} in clockwise partial order.)

3.1.4. Stability and geometric quotients. Let $(a, b) \in \Sigma_{\text{eff}}$. We define a G -invariant open affine subset $\mathcal{B}_{(a,b)} = \mathcal{A} \setminus V(J_{(a,b)}) \subset \mathcal{A}$, where $J_{(a,b)} \subset R_0$ is the homogeneous ideal

$$J_{(a,b)} = (\{x_i : (a_i, b_i) \leq (a, b)\}) \cap (\{x_i : (a_i, b_i) \geq (a, b)\})$$

and x_1, \dots, x_N are the coordinates on \mathcal{A} . (Note that $(\{x_i : (a_i, b_i) \leq (a, b)\}) \subset R_0$ is the ideal generated by those x_i on the anticlockwise side of (a, b) in Σ_{eff} .)

Recall, from [Dol03] 8.1 for instance, that a point $p \in \mathcal{A}$ is χ -stable if and only if the following three conditions hold: it is χ -semistable, its stabiliser $G_p \subset G$ is finite, and if $f \in R^{G,\chi}$ is as in the definition of χ -semistable, then for every $q \in D_f$, the orbit $G \cdot q$ is closed in D_f .

Proposition 3.3. *Fix a character χ corresponding to a primitive integral vector $(a, b) \in \Sigma_{\text{eff}} \cap C_G$. Then $\mathcal{A}_\chi^{ss} = \mathcal{B}_{(a,b)}$. If, moreover, (a, b) is in the interior of some chamber \mathcal{C}_j , then $\mathcal{A}_\chi^s = \mathcal{A}_\chi^{ss}$, and so the quotient $\mathcal{A} //_\chi G$ is geometric.*

Proof. A point $p \in \mathcal{A}$ lies in $\mathcal{B}_{(a,b)}$ if there exists a variable x_i with $(a_i, b_i) \leq (a, b)$ for which $x_i(p) \neq 0$ and another x_j (possibly equal to x_i) with $(a_j, b_j) \geq (a, b)$ for which

$x_j(p) \neq 0$. Let $m, m_i, m_j \in \mathbb{N}$ such that $m(a, b) = m_i(a_i, b_i) + m_j(a_j, b_j)$. If $F = x_i^{m_i} x_j^{m_j} t^m$ then $F \in R_m^{G_\chi}$ and $F(p, 1) \neq 0$. In particular $p \in \mathcal{A}_\chi^{ss}$ and $\mathcal{B}_{(a,b)} \subset \mathcal{A}_\chi^{ss}$.

Conversely, if $p \in \mathcal{A}_\chi^{ss}$, then for some $m > 0$ there exists $F \in R_m^{G_\chi}$ and $\tau \in \mathbb{C}^*$ such that $F(p, \tau) \neq 0$. Any monomial appearing in F is of the form $x_1^{i_1} \cdots x_N^{i_N} t^m$. After evaluating t at τ , such a monomial is in $J_{(a,b)}$; so $F|_{t=\tau}$ is also in $J_{(a,b)}$. Since $F|_{t=\tau}(p) \neq 0$, so $\mathcal{A}_\chi^{ss} \subset \mathcal{B}_{(a,b)}$ and the two sets are equal.

Suppose now that $\chi = (a, b)$ is the interior of some chamber. We show that $\mathcal{A}_\chi^{ss} \subset \mathcal{A}_\chi^s$. Let $p \in \mathcal{A}_\chi^{ss} = \mathcal{B}_{(a,b)}$. Fix i, j such that the two components $x_i(p), x_j(p)$ are nonzero and

$$(a_i, b_i) < (a, b) < (a_j, b_j).$$

In particular, $D_f \subset \mathcal{A}_\chi^{ss}$ for $f = x_i x_j$. If $(\lambda, \mu) \in G_p$, then $\lambda^{a_i} \mu^{b_i} = 1$ and $\lambda^{a_j} \mu^{b_j} = 1$; these equations have only finitely many solutions, so G_p is finite. By the same token, G_q is finite for any $q \in D_f$. So all orbits in this set are closed, and $p \in \mathcal{A}_\chi^s$. \square

3.1.5. Affine patches on the quotients. It is convenient to record here the natural affine patches on these geometric quotients, although we do not need them yet. Suppose that χ corresponds to (a, b) and does not lie on any ray r_α , $\alpha = 1, \dots, M$. Then, by Proposition 3.3,

$$\mathcal{A}_{i,j} = \mathcal{A} \setminus (x_i x_j = 0),$$

for any $i < j$ with $(a_i, b_i) < (a, b) < (a_j, b_j)$, is an open affine G -invariant set contained in \mathcal{A}_χ^{ss} , and moreover \mathcal{A}_χ^{ss} is covered by such sets $\mathcal{A}_{i,j}$.

If the two characters (a_i, b_i) and (a_j, b_j) are a \mathbb{Z} -basis for $\mathbb{Z} \oplus \mathbb{Z}$, then the usual identification of affine patches on \mathbb{P}^n with \mathbb{C}^n shows that $\mathcal{A}_{i,j} //_\chi G \cong \mathbb{C}^{N-2}$ and that $x_1, \dots, \widehat{x}_i, \dots, \widehat{x}_j, \dots, x_N$ serve as coordinates on this patch: the two degrees of freedom in the group action can be used to fix values for the i th and j th coordinates of a point leaving the other coordinates free. If instead (a_i, b_i) and (a_j, b_j) span a sublattice of C_G of index r , then the patch is a $\mathbb{Z}/r\mathbb{Z}$ quotient of \mathbb{C}^{N-2} in the same way as for patches on wps.

3.1.6. Extremal morphisms between the quotients. Fix a chamber \mathcal{C}_α with its boundary rays r_α and $r_{\alpha+1}$. By Proposition 3.3, the polarisations $\chi \in \mathcal{C}_\alpha$ give isomorphic geometric quotients (by well-defined isomorphisms), so set

$$\mathcal{Y}_\alpha = \text{Proj } R^{G,\chi} \quad \text{for some } \chi \in \mathcal{C}_\alpha.$$

Equally, the strictly semistable polarisations $(a, b) \in r_\alpha$ give isomorphic categorical quotients, so set

$$\mathcal{Z}_\alpha = \text{Proj } R^{G,\chi} \quad \text{for some } \chi \in r_\alpha.$$

Lemma 3.4. *Let $\alpha \in \{0, \dots, M-1\}$.*

- (a) \mathcal{Y}_α is \mathbb{Q} -factorial of dimension $N-2$ and Picard rank $\rho \leq 2$.
- (b) \mathcal{Y}_α admits two morphisms

$$\begin{array}{ccc} & \mathcal{Y}_\alpha & \\ F_\alpha \swarrow & & \searrow G_\alpha \\ \mathcal{Z}_\alpha & & \mathcal{Z}_{\alpha+1} \end{array}$$

either of which may be an isomorphism.

The task in (b) is to describe the degree 0 homomorphism of graded rings that is induced by the inclusion map of subsets of \mathcal{A} . We take advantage of the freedom in which graded ring we actually use as the homogeneous coordinate ring of \mathcal{Y}_α .

Proof. (a) The dimension count $\dim \mathcal{Y}_\alpha = \dim \mathcal{A}_\chi^{ss} - \dim G$ holds because the quotient is geometric. The \mathbb{Q} -factoriality follows from [HK00] Lemma 2.1; to confirm the hypotheses of that Lemma, invariant Cartier divisors on \mathcal{A}_χ^s are supported on the hyperplanes $x_i = 0$ and so can be linearised as required. Again following [HK00] Lemma 2.1, the map

$$\mathrm{Pic}(\mathcal{Y}_\alpha)_\mathbb{Q} \longrightarrow \mathrm{Pic}^G(\mathcal{A}_\chi^s)_\mathbb{Q}$$

is an isomorphism. If there is exactly one character lying strictly on one side of $r_\alpha \subset \Sigma_{\mathrm{eff}}$ —that character would have to be either χ_0 or χ_N —then the codomain of this map is \mathbb{Q} . Otherwise there are at least two characters on each side of r_α , and the codomain is \mathbb{Q}^2 .

(b) We describe $G_0: \mathcal{Y}_0 \rightarrow \mathcal{Z}_1$; other cases are similar. Choose characters χ_0, χ_1 lying in the rays r_0, r_1 respectively. Pick a set of generators f_1, \dots, f_v of $R_1 = R^{G, \chi_1}$, the homogeneous coordinate ring of \mathcal{Z}_1 . Let $g \in R_m^{G, \chi_0}$ be any nonzero polynomial of positive degree $m > 0$ (in t) in the homogeneous coordinate ring of \mathcal{Z}_0 .

The elements $F_i = f_i g^{\deg_t(f_i)}$ all lie in $R = R^{G, \chi}$, where $\chi = \chi_0^m \chi_1$ lies in the interior of the chamber \mathcal{C}_1 . So we use (some truncation of) R as the homogeneous coordinate ring of \mathcal{Y}_1 , and the map $R_1 \rightarrow R$ induced by $f_i \mapsto F_i$ is the required degree 0 graded k -algebra homomorphism. \square

3.1.7. Divisors on the quotients and exceptional loci. For a homogeneous element $x \in R_0$ define the *basic divisor* \mathcal{D}_x to be the divisor $x = 0$ in each quotient. (The notation is birational, and the variety on which \mathcal{D}_x is currently be considered will be mentioned explicitly if it matters. Of course, \mathcal{D}_x only makes sense on quotients $\mathcal{A}_\chi^{ss} // G$ for which x does not generate a component of the irrelevant ideal.)

Intersections of basic divisors describe the base locus of linear systems. Define

$$\mathcal{F}_\alpha = \bigcap_{\chi_i < r_{\alpha-1}} \mathcal{D}_{x_i} \subset \mathcal{Y}_\alpha \quad \text{and} \quad \mathcal{G}_\alpha = \bigcap_{\chi_i > r_{\alpha+1}} \mathcal{D}_{x_i} \subset \mathcal{Y}_\alpha.$$

(As usual, χ_i is the character associated to variable x_i .) Also, let

$$\mathcal{B}_\alpha = \bigcap_{\chi_i \in r_\alpha} \mathcal{D}_{x_i} \subset \mathcal{Z}_\alpha.$$

The maps F_α and G_α from \mathcal{Y}_α are defined in Lemma 3.4.

Lemma 3.5. *Let $\alpha \in \{0, \dots, M-1\}$.*

(a) *The exceptional locus of the morphism F_α is \mathcal{F}_α , and the restriction of F_α to the exceptional locus $F_\alpha: \mathcal{F}_\alpha \rightarrow \mathcal{B}_\alpha$ exhibits \mathcal{F}_α as a weighted projective bundle over \mathcal{B}_α . Similarly $G_\alpha: \mathcal{G}_\alpha \rightarrow \mathcal{B}_{\alpha+1}$ is a weighted projective bundle over $\mathcal{B}_{\alpha+1}$.*

(b) *The map $\mathcal{Y}_\alpha \dashrightarrow \mathcal{Y}_{\alpha+1}$ in the diagram*

$$\begin{array}{ccc} \mathcal{Y}_\alpha & \dashrightarrow & \mathcal{Y}_{\alpha+1} \\ G_\alpha \searrow & & \swarrow F_{\alpha+1} \\ & \mathcal{Z}_{\alpha+1} & \end{array}$$

is a \mathcal{D}_x -flop for any x whose character lies in the ray r_α .

Proof. We prove the requirements for F_α . To prove (a), consider first the case $F_0: \mathcal{Y}_0 \rightarrow \mathcal{Z}_0$. In this case, \mathcal{Z}_0 coincides with \mathcal{B}_0 and is a wps (possibly just a point), and the map F_0 has effective \mathbb{C}^* quotients of affine space punctured at the origin as its fibres. The general case follows, since $F_\alpha: \mathcal{F}_\alpha \rightarrow \mathcal{B}_\alpha$ is of the same form as the F_0 just considered.

To prove (b), note that any such x has zero locus a divisor in \mathcal{B}_α and so is certainly trivial on the fibres of F_α . The exceptional loci are irreducible so the relative Picard rank is one. \square

3.1.8. *Localisation and flips.* To understand the birational map $\mathcal{Y}_\alpha \dashrightarrow \mathcal{Y}_{\alpha+1}$ better, we realise it as a variation of \mathbb{C}^* action in a neighbourhood of a point in the base \mathcal{Z}_α . If $x_l, x_{l+1}, \dots, x_{l+k}$ are the variables whose characters lie in r_α , we consider the case of the point $x_{l+1} = \dots = x_{l+k} = 0$ in $\mathcal{B}_\alpha \subset \mathcal{Z}_\alpha$ and that χ_l is a primitive vector in C_G . (This is not quite the general case, but in our applications it is enough.) The quotient map $C_G \rightarrow \mathbb{Z}$ with kernel generated by χ maps characters $\chi' < \chi$ to the negative integers and those $\chi'' > \chi$ to the positive integers. These are weights for a \mathbb{C}^* action on \mathbb{C}^{N-k} . The variation of that action, as in [Bro99] for instance, describes a (generalised) flip in an affine patch in \mathcal{Z}_α .

In more complicated cases later, when we consider equations inside the toric flip, we have to shrink this affine neighbourhood to an analytic neighbourhood of the point to describe a flip.

3.1.9. *The proof of Theorem 3.1.* Given $\mathcal{Y} \rightarrow \mathcal{X}$, let $\mathcal{N} = C(\mathcal{Y})$ be the Cox ring of \mathcal{Y} ; by construction, this is a polynomial ring (with N variables, say) with a grading by $\mathbb{Z} \oplus \mathbb{Z}$. Moreover, the left-hand ray r_0 of Σ_{eff} contains a single character, corresponding to the new element of the 1-skeleton. The variation of this quotient described above produces a link diagram as required. (We must omit the quotient by characters in the first chamber \mathcal{C}_0 ; it is isomorphic to \mathcal{X} , because the left-hand ray r_0 contains only a single character, and its two morphisms are the structure map from \mathcal{X} to a point and an isomorphism $\mathcal{X} \cong \mathcal{Z}_1$. Similarly, we omit the quotient in the final chamber if the previous map is also a divisorial contraction.) The maps are isomorphisms in codimension 1 because the contracted loci, as described by Lemma 3.4, are geometric quotients of (a stable set in) \mathbb{C}^k for $k \leq N-2$ by $\mathbb{C}^* \times \mathbb{C}^*$ so have dimension at most $N-4$, which is codimension 2 in the quotients. The possible behaviour at the two ends is completely determined by the 2-ray game.

3.2. **Examples.** For calculations, it is convenient to summarise the action by a matrix of integers

$$\begin{pmatrix} a_1 & a_2 & \cdots & a_N \\ b_1 & b_2 & \cdots & b_N \end{pmatrix}.$$

The lemmas above can then be summarised by a sketch of the layout of this matrix χ near \mathcal{Y}_α :

$$\left(\begin{array}{c|c|c|c} I_{\mathcal{F}_\alpha} & & \mathcal{Y}_\alpha & I_{G_\alpha} \\ & \mathcal{Z}_\alpha & & \\ \hline x_1 \dots x_{m_{\alpha-1}} & x_{l_\alpha} \dots x_{m_\alpha} & x_{l_{\alpha+1}} \dots x_{m_{\alpha+1}} & x_{l_{\alpha+2}} \dots x_N \end{array} \right)$$

where $x_{l_\alpha}, \dots, x_{m_\alpha}$ are the generators of \mathcal{N} whose characters lie in the ray R_α .

3.2.1. *Standard flop between two scrolls.* Consider the graded polynomial ring

$$\mathcal{N} = \mathbb{C}[x_1, x_2, y, z_1, z_2] \quad \text{with weights} \quad \begin{pmatrix} 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 \end{pmatrix}.$$

That is, x_1, \dots, z_2 are eigencoordinates on \mathbb{C}^5 for the action of $G = \mathbb{C}^* \times \mathbb{C}^*$ in which x_1 is in the $(0, 1)$ -eigenspace, y is in the $(1, 1)$ -eigenspace, etc. Setting

$$\begin{aligned} \mathcal{B}_0 &= \mathbb{C}^5 \setminus \{(x_1 = x_2 = 0) \cup (y = z_1 = z_2 = 0)\}, \\ \mathcal{A}_0 &= \mathbb{C}^5 \setminus \{(x_1 = x_2 = y = 0) \cup (y = z_1 = z_2 = 0)\}, \\ \mathcal{B}_1 &= \mathbb{C}^5 \setminus \{(x_1 = x_2 = y = 0) \cup (z_1 = z_2 = 0)\} \end{aligned}$$

write

$$\mathcal{Y} = \mathcal{B}_0/G, \quad \mathcal{Z} = \mathcal{A}_0//G, \quad \mathcal{Y}' = \mathcal{B}_1/G.$$

These three varieties fit in the rank 2 link

$$\begin{array}{ccccc} & & \mathcal{Y} & \dashrightarrow & \mathcal{Y}' & & \\ & \swarrow & & & & \searrow & \\ \mathbb{P}^1 & & & & \mathcal{Z} & & \mathbb{P}^1 \end{array}$$

in which the map $\mathcal{Y} \dashrightarrow \mathcal{Y}'$ is the standard flop. This link is one of the 3-fold analogues of $\mathbb{P}^1 \times \mathbb{P}^1$ with its two fibration structures.

A different choice of weights such as

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & -1 & -1 \end{pmatrix}$$

is more symmetric and certainly does determine the same link as that of the original. It has the advantage that one can see immediately that the flip is the \mathbb{C}^* flip by quotients of \mathbb{C}^4 by the \mathbb{C}^* action $(1, 1, -1, -1)$ —the bottom row of the matrix ‘localised’ at a point in the base. But the disadvantage is that it presents the flopping variety \mathcal{Z}' embedded with quasi-reflections for

$$\mathcal{Z}' = \text{Proj } \mathbb{C}[y, x_1 z_1, x_1 z_2, x_2 z_1, x_2 z_2] \subset \mathbb{P}(1, 2, 2, 2, 2).$$

3.2.2. *Francia antiflip.* Consider the action of G on $\mathbb{C}[x_1, x_2, y, z_1, z_2]$ given by weights

$$\begin{pmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & -1 & -2 & -3 \end{pmatrix}.$$

Calculating the quotients this time describes a Sarkisov link

$$\begin{array}{ccccc} & & \mathcal{Y} & \dashrightarrow & \mathcal{Y}' & & \\ & \swarrow & & & & \searrow & \\ \mathbb{P}^3 & & & & \mathcal{Z} & & \mathbb{P}(1, 1, 2, 3) \end{array}$$

in which one end is the $(1, 2, 3)$ -blowup of a point of \mathbb{P}^3 , the other is the $(1, 1, 2)$ -blowup of a nonsingular point of $\mathbb{P}(1, 1, 2, 3)$, and the map $\mathcal{Y} \dashrightarrow \mathcal{Y}'$ is the opposite of the Francia flip.

We describe this in terms of toric geometry. Start with the usual fan for \mathbb{P}^3 in \mathbb{Z}^3 . After introducing the ray $(1, -1, 2)$ and subdividing minimally, giving the fan $\Delta_{\mathcal{Y}}$, the convex cone

$$\langle (1, -1, 2), (0, 1, 0), (1, 0, 0), (0, 0, 1) \rangle$$

is divided into two simplexes by the face joining $(1, 0, 0)$ and $(0, 0, 1)$. The Francia antiflip is the operation on fans $\Delta_{\mathcal{Y}} \rightsquigarrow \Delta_{\mathcal{Y}'}$ that removes this edge and replaces it by the edge joining $(1, -2, 1)$ and $(0, 1, 0)$: indeed in the notation of [Rei83]

$$1 \times (1, -1, 2) + 1 \times (0, 1, 0) = 1 \times (1, 0, 0) + 2 \times (0, 0, 1).$$

Since in $\Delta_{\mathcal{Y}'}$

$$(0, 0, 1) = 1 \times (1, -1, 2) + 1 \times (-1, -1, -1) + 2 \times (0, 1, 0)$$

we can remove the ray $(0, 0, 1)$, the opposite of a $(1, 1, 2)$ -blowup of a nonsingular point. The result is $\mathbb{P}(1, 1, 2, 3)$ because

$$1 \times (1, 0, 0) + 1 \times (1, -1, 2) + 2 \times (-1, -1, -1) + 3 \times (0, 1, 0) = (0, 0, 0).$$

As another example, the action

$$\begin{pmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & -1 & -2 & -5 \end{pmatrix}$$

is similar: the corresponding link starts with the $(1, 2, 5)$ -blowup of \mathbb{P}^3 , makes the antipodal flip of type $(1, 1, -1, -4)$, and finishes with (the inverse of) the Kawamata blowup of $1/3(1, 1, 2)$ in $\mathbb{P}(1, 3, 4, 5)$. One can check that these are the only two such examples of Sarkisov links starting from \mathbb{P}^3 : more precisely, any other calculation with $N = 5$ and ordinary \mathbb{P}^3 at one end has nonterminal singularities emerging from the antipodal flip.

3.2.3. *Change of structure.* The action

$$\begin{pmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 2 & 0 & -1 & -1 \end{pmatrix}.$$

is similar to the first example, being

$$\begin{array}{ccccc} & & \mathcal{Y} & & \mathcal{Y}' & & \\ & \swarrow & & \searrow & \swarrow & \searrow & \\ \mathbb{P}^3 & & & & \mathcal{Z} & & \mathbb{P}^1 \end{array}$$

where $\mathcal{Y} \dashrightarrow \mathcal{Y}'$ is the Francia flip—the toric flip with weights $(1, 2, -1, -1)$ as [Bro99] Section 1—and $\mathcal{Y}' \rightarrow \mathbb{P}^1$ has fibres $\mathbb{P}(1, 1, 3)$. (In fact, the 3-folds here each have a line of canonical singularities, so this is not a Sarkisov link: the map $\mathcal{Y} \rightarrow \mathbb{P}^3$ is the $(2, 3, 3)$ blowup of a point as one sees immediately by localising.)

In the toric fan after this flip, the plane spanned by the new edge comprises cones of the fan, so the variety \mathcal{Y}' admits a toric morphism to \mathbb{P}^1 .

4. GENERAL FANO 3-FOLDS IN CODIMENSION 3

4.1. The equations of Fano 3-folds.

4.1.1. *Weighted Grassmannians and pfaffian equations.* We consider explicit Fano 3-folds $X \subset \mathbb{P}^6 = \mathbb{P}^6(a_0, \dots, a_6)$ other than the standard complete intersection $X_{2,2,2} \subset \mathbb{P}^6$. There are 69 families of such Fano 3-folds, and they are listed in [BDK⁺] with their degrees, baskets and Hilbert numerators. (See [ABR02] Section 4.6, or [Bro07] Section 2.1 in the context of K3 surfaces, for an account of these notions.)

A general member of each family is quasismooth and has singularities equal to those of the basket. It has equations that are the five maximal pfaffians of a skew 5×5 matrix of homogeneous forms, and it is easy to construct a general member by writing down such a matrix; this matrix is also the first syzygy matrix of the equations, so the degrees of forms appearing are determined by the Hilbert numerator. When focussing on a particular singular point of such a Fano, we can usually make coordinate changes to fix a large proportion of the free parameters, as exemplified by [BRK].

Example No. 10: $X \subset \mathbb{P}^6(1, 1, 1, 2, 2, 2, 3)$ with

$$\mathbf{Degree} = 13/6, \quad \mathbf{Basket} = \left\{ 3 \times \frac{1}{2}(1, 1, 1), \frac{1}{3}(1, 1, 2) \right\}.$$

This example has Hilbert numerator $1 - 3t^4 - 2t^5 + 2t^6 + 3t^7 - t^{11}$ so we construct the five Pfaffian equations to include three of degree 4 and two of degree 5. Let $x, x_1, x_2, y, y_1, y_2, z$ be the weighted homogeneous coordinates of \mathbb{P}^6 in that order. We consider each type of singularity in turn as a centre.

We move a singularity of type $\frac{1}{2}(1, 1, 1)$ at the coordinate point P_y by a change of coordinates. To ensure $P_y \in X$ is quasismooth, the equations near P_y must eliminate the

variables y_1, y_2 and one other. This is easily arranged by writing the syzygy matrix as

$$\begin{pmatrix} y & A_3 & B_3 & C_2 \\ & D_2 & E_2 & F_1 \\ & & z & y_2 \\ & & & y_1 \end{pmatrix}$$

where A, \dots, F are general forms of the degrees indicated. By row and column operations on the matrix, we assume further that y does not appear in A, \dots, E and that $F_1 = x$. Apart from being quasismooth, the only appeal to generality is that the 3–4 entry of the syzygy matrix is z rather than xy . (In [CPR00], this kind of assumption that a particular monomial appears in the equations is called ‘starred monomial’. It is needed for the vanishing calculation of Lemma 4.6 below and omitting the starred monomial will typically change the link.)

The link from the $\frac{1}{3}(1, 1, 2)$ point can be best computed by presenting the equations of a general X by syzygy matrix

$$\begin{pmatrix} z & A'_3 & B'_2 & C'_2 \\ & D'_3 & E'_2 & F'_2 \\ & & y_1 & y \\ & & & x \end{pmatrix}$$

for general forms A', \dots, F' . It is possible to write the syzygy matrix so that both of the links above can be ‘seen’ in its format, but we do not attempt to do this.

4.1.2. *Type I singularities.* If $X \subset \mathbb{P}^n$ is quasismooth at $p \in X$, then there must be $n - 3$ polynomials in the ideal of $\mathcal{O}_{X,p}$ that vanish and are independently linear near p . If, as is typical, $p = p_w$ is at a coordinate point of \mathbb{P}^n , then the ideal of X must contain $n - 3$ equations of the form

$$w^k \ell = w^{k-1} m_{k-1} + \dots + m_0$$

with ℓ containing at least one term that is a linear variable—that is, ℓ is a *quasilinear* form—and polynomials m_i in the other variables. The $n - 3$ homogeneous quasilinear polynomials like ℓ are called the *tangent polynomials* at p .

Definition 4.1 ([ABR02] Section 5.5). Let $X \subset \mathbb{P}^n$ be a Fano 3-fold, and $p \in X$ be a quasismooth point of X ; in particular, $p \in X$ is a quotient singularity of type $\frac{1}{r}(1, a, r - a)$ for some coprime $0 < a < r$.

Then $p \in X$ is a *Type I point* if and only if p lies at one of the coordinate points $p = p_w$ of \mathbb{P}^n and there are coordinate functions x, y, z on \mathbb{P}^n that vanish at p , have weights $1, a, r - a$ respectively and are independent in m_p/m_p^2 modulo the tangent polynomials.

A Type I point $p \in X$ is a *Type I centre* if and only if $K_X^3 > \frac{1}{ra(r-a)}$.

This definition is perfect for our calculations, although it would be slightly awkward to use more generally: the term *centre* usually applies to a component of the base locus of a linear system which is determined by a birational map, and this definition is anticipating such a map.

We illustrate the definition with a non-example. The Fano 3-fold

$$X_{66}: (x^{66} + xy^{13} + z^{11} + t^3 - u^2 = 0) \subset \mathbb{P}(1, 5, 6, 22, 33)$$

(with variables x, y, z, t, u in that order) is quasismooth and contains the point $p = (0, 0, 0, 1, 1)$ as a singularity of type $\frac{1}{11}(1, 5, 6)$. But $p \in X$ is *not* a Type I point according to the definition, because it does not (and cannot even after coordinate changes) lie at a coordinate point.

Proposition 4.2. *Let $X \subset \mathbb{P}^6$ be a general member of the 69 codimension 3 families of Fano 3-folds and $p \in X$ be a singularity of type $\frac{1}{r}(a, b, c)$ which is a Type I centre. We may choose coordinates so that p lies at a coordinate point $p_t \in \mathbb{P}^6$. Then the syzygy matrix of $X \subset \mathbb{P}^6$ can be written as*

$$\begin{pmatrix} t & A & B & C \\ & D & E & F \\ & & y_3 & y_2 \\ & & & y_1 \end{pmatrix}$$

where y_1, y_2, y_3 are the tangent polynomials at p and A, \dots, F are forms (of degree determined by the Hilbert numerator of X) in all the variables. Without loss of generality, y_1, y_2, y_3 are three distinct coordinate functions on \mathbb{P}^6 .

Proof. All claims in this proposition come from direct observation of the list of codimension 3 Fanos. The form of the syzygy matrix follows from the definition of Type I point after one has checked that the equations are of the form $ty_i = \dots$ rather than $t^2y_i = \dots$ or even higher powers of t in the tangent monomial. \square

Notation 4.3. Let $p \in X$ be Type I centre on a Fano as in Proposition 4.2. We choose coordinates so that

- $X \subset \mathbb{P}^6(a, b, c, r, a_1, a_2, a_3)$ with coordinates $x, y, z, t, y_1, y_2, y_3$ in that order,
- $p = p_t \in X$ is the t -coordinate point,
- x, y, z are the eigencoordinates of the \mathbb{Z}/r stabiliser at p ,
- y_1, y_2, y_3 are the tangent polynomials at p ,
- the weights of the tangent polynomials satisfy $a_1 \leq a_2 \leq a_3$.

The variables x, y, z are also called the *polarising variables* at p .

Corollary 4.4 (of Proposition 4.2). *Using Notation 4.3, let Π be the locus $t = y_1 = y_2 = y_3 = 0$ in \mathbb{P}^6 . Then $\Pi \cong \mathbb{P}^2(a, b, c)$ and $X \cap \Pi$ is the locus defined by the three 2×2 minors*

$$\left(\bigwedge^2 \begin{pmatrix} A & B & C \\ D & E & F \end{pmatrix} = 0 \right) \subset \Pi = \mathbb{P}^2(a, b, c)$$

which is a finite reduced set of nonsingular points of $\mathbb{P}^2(a, b, c)$.

In fact, setting $d = \deg A - \deg D$ (which equals $\deg B - \deg E$ and $\deg C - \deg F$) and

$$d_i = \frac{1}{2} \left((-1)^i d + \sum_{j=1}^3 (a_j + r) \right) \quad \text{for } i = 1, 2,$$

the number of points of $X \cap \Pi$ is $N = \sum_{i \neq j} (a_i + r)(a_j + r) - d_1 d_2$.

Proof. The finiteness of $X \cap \Pi$ is automatic since otherwise the displayed minors would define a divisor in X contradicting $\rho_X = 1$. The extra claim is that the degrees of these minors are sufficiently high and divisible in each case so that they determine free linear systems on $\mathbb{P}^2(a, b, c)$, from which it follows that for general X these minors have common solutions only away from the toric strata of $\mathbb{P}^2(a, b, c)$. This is checked for each of the 69 cases in turn.

The counting argument is standard use of the Hilbert–Burch theorem. If $Z = X \cap \Pi$, then there is an exact sequence of $\mathcal{O} = \mathcal{O}_\Pi$ -modules

$$0 \rightarrow \mathcal{O}(-d_1) \oplus \mathcal{O}(-d_2) \rightarrow \bigoplus_{i=1}^3 \mathcal{O}(-(a_i + r)) \rightarrow \mathcal{O} \rightarrow \mathcal{O}_Z \rightarrow 0$$

(for suitable d_1, d_2) where the two nontrivial maps are given by the matrix

$$M = \begin{pmatrix} A & B & C \\ D & E & F \end{pmatrix}$$

and the vector of its (signed) 2×2 minors respectively. We compute chern classes to conclude. The equality of c_1 for the third and fourth terms gives the formula for d_i ; that is, comparison of degrees in the exact sequence. Then computing the Hilbert polynomial of Z , or equivalently c_2 , gives the required number. \square

4.2. The proof of Theorem 1.1. Let $Y \rightarrow X$ be the Kawamata blowup of p with exceptional divisor $E \subset Y$; note that $\rho_Y = 2$. Define $F_i \subset Y$ to be the birational transform on Y of $(y_i = 0) \subset X$.

Lemma 4.5. *The intersection $\Gamma = F_1 \cap F_2 \cap F_3$ is a reduced union of disjoint rational curves $\Gamma = \cup \Gamma_i$. For each i ,*

$$E\Gamma_i = 1 \quad \text{and} \quad A\Gamma_i = 1/r,$$

and so $B\Gamma_i = 0$ and $\Gamma_i \stackrel{\text{num}}{\sim} \Gamma_j$ for all i, j . (The number of components $\Gamma_i \subset \Gamma$ is the number of points, N , computed in Corollary 4.4.)

Proof. By Corollary 4.4, $\Gamma_X = (y_1 = y_2 = y_3 = 0) \subset X$ is the cone on a finite set of reduced points with vertex at p —in particular, its components are irreducible rational curves. The Kawamata blowup is the blowup of the ideal at p generated by the polarising variables, so $E \subset Y$, the preimage of p , is isomorphic to $\mathbb{P}(a, b, c)$ and the birational transform of Γ_X intersects E in the same Hilbert–Burch locus as $X \cap \Pi$.

Thus $E\Gamma_i = 1$, and $A\Gamma_i = 1/r$ is because $rA \cap \Gamma$ computed on X is the same as $\Pi \cap \Gamma$. The numerical equivalence of the Γ_i follows since $\rho_Y = 2$ implies that pair A, E span $\overline{NE}^1(Y)$. \square

We denote $A = -K_X$ and $B = -K_Y$, and record that on Y

$$B = A - (1/r)E \quad \text{and} \quad E^3 = r^2/(abc),$$

so Lemma 4.5 shows that $B\Gamma_i = 0$.

Any function f of weight k on X with $f(p) = 0$ vanishes to order at least k/r on E when pulled back to Y . For the polarising variables x, y, z , as functions on Y we have

$$(4.1) \quad x \in \mathcal{O}_Y(aA - \frac{a}{r}E), \quad y \in \mathcal{O}_Y(bA - \frac{b}{r}E), \quad z \in \mathcal{O}_Y(cA - \frac{c}{r}E).$$

(See [CPR00], Proposition 3.4.6, for instance.) By the next lemma, the tangent polynomials y_i vanish on E exactly once more than generic functions of their degree.

Lemma 4.6. *For each $i = 1, 2, 3$,*

$$F_i \sim a_i A - ((a_i + r)/r)E = a_i B - E.$$


where \sim denotes linear equivalence (and so a fortiori numerical equivalence).

Proof. By Proposition 4.2, t does not vanish near $p \in X$ and

$$y_i = \frac{\textit{ith minor}}{t} \in \mathcal{O}(a_i A).$$

Pulling this back to Y , the minor has degree $a_i + r$ so it vanishes to order at least $(a_i + r)/r$ along E . But in each case, the minor includes a nontrivial monomial in the polarising variables x, y, z —by generality of X in its family—so that by (4.1) the vanishing is exactly $(a_i + r)/r$ along E as claimed. \square

According to this lemma, we can sketch in $N^1(Y) \cong \mathbb{R}^2$ the rays through the various divisors as follows (noting that rays through the F_i may coincide).



Corollary 4.7. *B is nef on Y and it supports an extremal ray of $\overline{\text{NE}}(Y)$; in the notation of Lemma 4.5, that ray is $\mathbb{R}_+[\Gamma]$ and it is a flopping ray.*

Proof. Lemma 4.6 implies that $(a_1 + r)B = F_1 + rA$, so if $B\gamma < 0$ then $\gamma \subset \cap F_i$. So B is nef since each $B\Gamma_i = 0$, and moreover it supports the ray $\mathbb{R}_+[\Gamma]$. \square

Let $Y \rightarrow Z$ be the contraction of the ray $\mathbb{R}_+[\Gamma]$. Then Z has Picard rank 1 and B is the ample generator, so in particular Z is a (non \mathbb{Q} -factorial) Fano 3-fold.

Lemma 4.8. *Let $Z \leftarrow Y_1$ be the flop of $Y \rightarrow Z$. Then on Y_1*

(a) *the intersection $F_1 \cap F_2 \cap F_3$ is empty.*

(b) *F_3 is nef.*

(c) *$F_1 \cap F_2$ is a non-empty union of curves and $F_3C = 0$ for every curve $C \subset F_1 \cap F_2$.*

(d) *the second extremal ray of the Mori cone $\overline{\text{NE}}(Y_1)$ equals $\mathbb{R}_+[C]$, where C is any reduced and irreducible component of $F_1 \cap F_2$. This ray is supported by F_3 and it is contracted by the extremal morphism φ_{F_3} .*

Proof. (a) The flop $Z \leftarrow Y_1$ restricted to $E \cong \mathbb{P}(a, b, c)$ is the blowup of the locus of points $(y_1 = y_2 = y_3 = 0) \subset E \subset Z$. This is the same locus as computed in Corollary 4.4 since the y_i are equal to the minors on E . To make the flop is to blowup this locus; and by construction, the generators of the blowup ideal have no common solutions on the blowup.

(b) If on Y_1 a curve γ has $F_3\gamma < 0$ then also $F_i\gamma < 0$ for $i = 2, 3$, since $a_3 \geq a_i$, so $\gamma \subset F_1 \cap F_2 \cap F_3$. But there are no such curves, so F_3 is nef on Y_1 .

(c) The intersection of any pair $F_i \cap F_j$ does not contain a surface (since $Y \dashrightarrow Y_1$ is an isomorphism in codimension 1), so is at most one dimensional. We compute $F_1 \cap F_2$ on X —the map to Y_1 is a blowup of $p \in X$ followed by a flop, so the result follows. Setting $y_1 = y_2 = 0$ in the equations of X cuts out the flopping curves Γ_X together with the distinct locus

$$(C = F = y_3t - AE + BD = 0) \subset \mathbb{P}^4(r, a, b, c, a_3).$$

This latter locus is not empty and is at least one dimensional. Its birational transform on Y_1 is the locus $F_1 \cap F_2$ in question, and its intersection with F_3 is trivial by (a).

(d) Follows immediately from (a–c). \square

The continuation of the link from Y_1 depends on strictness of the inequalities $a_1 \leq a_2 \leq a_3$.

Proposition 4.9. *Suppose that $a_2 = a_3$. Then the contraction of the extremal ray by $\varphi_{F_3}: Y_1 \rightarrow X'$ determines a morphism to a \mathbb{Q} -factorial variety X' . There are two subcases:*

- *if $a_1 = a_2$ then $\dim X' = 2$ and φ_{F_3} is a conic bundle.*
- *if $a_1 < a_2$ then $\dim X' = 3$ and φ_{F_3} is a divisorial contraction that contracts an irreducible divisor to a line in X' .*

Proof. By assumption $F_2 \sim F_3$. So by Lemma 4.8(a) the restriction of F_3 to F_1 contains a free pencil on F_1 , and so it is in fact a multiple of a free pencil. In particular, there are effective curves γ with $F_3\gamma = 0$. Since F_3 is nef, any such curve generates an extremal ray, and so the map φ_{F_3} contracts F_1 to a line.

Now the classification of extremal contractions completes the proof. First, φ_{F_3} is not small so its image is \mathbb{Q} -factorial. If $a_1 = a_2$ then φ_{F_3} also contracts each of F_2 and F_3 to a line and so φ_{F_3} can only be a conic bundle. If $a_1 < a_2$ then on Y_1

$$F_3^3 = F_3F_2(F_1 + (a_3 - a_1)B) = (a_3 - a_1)BF_3F_2 > 0,$$

the positivity holding because $F_2 \cap F_3$ contains non-flopping curves. So the image of φ_{F_3} is a 3-fold and the only possibility is that φ_{F_3} is a divisorial contraction to a line. \square

Proposition 4.10. *Suppose that $a_2 < a_3$. Then the contraction of the extremal ray by $\varphi_{F_3}: Y_1 \rightarrow Z_2$ is a flipping contraction and the flip $Z_2 \leftarrow Y_2$ exists.*

The second extremal ray of the Mori cone $\overline{NE}^1(Y_2)$ is supported by the divisor F_2 , and the contraction of this extremal ray $\varphi_{F_2}: Y_2 \rightarrow X'$ determines a morphism to a \mathbb{Q} -factorial variety X' . There are two subcases:

- if $a_1 = a_2$ then $\dim X' = 1$ and φ_{F_2} is a del Pezzo fibration.
- if $a_1 < a_2$ then $\dim X' = 3$ and φ_{F_2} is a divisorial contraction that contracts an irreducible divisor to a point in X' .

Proof. Since $a_2 < a_3$, any curve γ with $F_3\gamma = 0$ has $F_2\gamma < 0$ and $F_1\gamma < 0$ so lies in $F_1 \cap F_2$. There are only finitely many such curves and so the extremal neighbourhood supported by F_3 is isolated; it is flipping because B is nef on Y_1 so strictly positive on the ray.

The flip exists because the exceptional locus is cut out by two divisors and Mori's easy flip theorem, [Kol92] Theorem 20.11, applies. Furthermore, by Mori's theorem, the intersection $F_1 \cap F_2$ is empty on the flipped variety Y_2 . (Of course, because it resolves the 'weighted pencil' $F_1: F_2$, Mori's Theorem requires the two divisors to be proportional in $\text{Pic}(Y_1)$, whereas F_1 and F_2 may not be. But they are proportional in a neighbourhood of the flipping locus. For instance, if we set $F'_1 = (a_3 - a_2)F_1 + (a_2 - a_1)F_3$, which is linearly equivalent to $(a_3 - a_1)F_2$, then $F'_1 \cap F_2$ equals $F_1 \cap F_2$ set-theoretically away from F_3 because the flop cleared that triple intersection. So on Y_2 we have that $F_1 \cap F_2 = \emptyset$ off F_3 ; but, again because of the flop, this is also true on F_3 .)

Let Y_2 be the flipped variety. As for F_3 in Lemma 4.8, F_2 is nef. In fact, F_2 supports the second extremal ray on Y_2 , because it is trivial on any curve contained in F_1 . Moreover, the extremal contraction φ_{F_2} contracts F_1 so is not small. Once again the classification of extremal contractions finishes the proof. First, the image of φ_{F_2} is \mathbb{Q} -factorial. If $a_1 = a_2$, then φ_{F_2} contracts any surface in the pencil $|F_1|$ to a point and so must be a del Pezzo fibration. If $a_1 < a_2$ then

$$F_2^3 = F_2(F_1 + (a_2 - a_1)B)^2 = (a_2 - a_1)^2 F_2 B^2 > 0,$$

the positivity holding because B is ample. So the image of φ_{F_2} is a 3-fold and the only possibility is that φ_{F_2} is a divisorial contraction to a point. \square

4.3. Supplements to Theorem 1.1.

4.3.1. *Calculations in the ambient toric 2-ray link.* The key point in all of the cases here is that by Lemma 4.6 the generators of the unprojection ideal all vanish to order 1 along the unprojection divisor. This is not a consequence of the Type I definition, but of the

A row operation on the character matrix renders it as

$$\begin{pmatrix} 3 & 5 & 1 & 1 & 1 & 0 & -1 & -1 \\ -1 & -1 & 0 & 0 & 0 & 1 & 1 & 1 \end{pmatrix}$$

and equations for y_1z and y_1x_i for some i eliminate two of the variables in a neighbourhood of P_{y_1} , so that after the usual flop we see a toric hypersurface flip of type $(3, 1, 1, -1, -1; 2)$, where the 2 indicates the weight of the hypersurface. The resulting 3-fold has a fibration to \mathbb{P}^1 , given by the ratio of the last two variables, with cubic surfaces as the fibres—these can already be seen in the syzygy matrix.

4.3.2. *End of the link I: equations from unprojection.* The calculation uses on Reid and Takagi's example of a so-called Type III unprojection [Rei00], 9.16. We sketch the general approach and then give a typical example of the method.

4.3.3. *A special case of Type III unprojection.* Consider \mathbb{C}^{10} with coordinates $t, x, x_1, x_2, A, B, C, D, E, F$. Let $V \subset \mathbb{C}^{10}$ be the 7-fold defined by the five maximal Pfaffians of the following 5×5 skew matrix (where, as usual, we omit the diagonal of zeros and the skew lower triangular half):

$$\begin{pmatrix} t & A & B & C \\ & D & E & F \\ & & x_2 & x_1 \\ & & & x \end{pmatrix}.$$

Of course, $V \cong \mathcal{C} \text{Grass}(2, 5)$, the cone on the Grassmannian in its Plücker embedding, and the equations are neatly expressed in terms of matrix entries m_{ij} by

$$\varphi_\ell = m_{hi}m_{jk} - m_{hj}m_{ik} + m_{hk}m_{ij}$$

for the five ways of choosing $\{h, i, j, k, \ell\} = \{1, 2, 3, 4, 5\}$ with h, i, j, k in increasing order.

The elimination of t (which appears only in Pfaffians $\varphi_3, \varphi_4, \varphi_5$) from these equations determines a birational map from V to

$$W: (\varphi_1 = \varphi_2 = 0) \subset \mathbb{C}^9.$$

The locus $(x = 0)$ in W splits into two components $E \cup F$, where the ideals of these loci in \mathbb{C}^9 are

$$I_E = (x, x_1, x_2) \quad \text{and} \quad I_F = \left(x, \bigwedge^2 \begin{pmatrix} B & C \\ E & F \\ x_2 & -x_1 \end{pmatrix} \right)$$

and $\wedge^2 M$ denotes the three 2×2 minors of a matrix M . In fact, E is the unique exceptional divisor of $W \dashrightarrow V$, and the unprojection construction we describe next will produce another birational map $W \dashrightarrow U$ —the so-called *Type III* unprojection—for which F is the unique exceptional divisor.

Unprojections can be computed by comparing free resolutions, so setting $\mathcal{O} = \mathcal{O}_{\mathbb{C}^9}$ we write

$$\begin{array}{ccccccccccc} 0 & \leftarrow & \mathcal{O}_F & \leftarrow & \mathcal{O} & \xleftarrow{M_0} & 4\mathcal{O} & \xleftarrow{M_1} & 5\mathcal{O} & \xleftarrow{M_2} & 2\mathcal{O} & \leftarrow & 0 \\ & & \uparrow & & \parallel & & J_1 \uparrow & & J_2 \uparrow & & & & \\ 0 & \leftarrow & \mathcal{O}_X & \leftarrow & \mathcal{O} & \xleftarrow{(\varphi_2, -\varphi_1)} & 2\mathcal{O} & \xleftarrow{t(\varphi_1, \varphi_2)} & \mathcal{O} & \leftarrow & & & 0 \end{array}$$

where the free resolution of \mathcal{O}_F is computed by $M_0 = (-x, \varphi_1, \varphi_2, \varphi_3)$ and

$$M_1 = \begin{pmatrix} \varphi_1 & \varphi_2 & \varphi_3 & \varphi_4 & \varphi_5 \\ x & 0 & 0 & -C & B \\ 0 & x & 0 & -F & E \\ 0 & 0 & x & -x_1 & x_2 \end{pmatrix}, \quad M_2 = \begin{pmatrix} B & C \\ E & F \\ x_2 & x_1 \\ 0 & x \\ -x & 0 \end{pmatrix}$$

and the comparison maps are

$$J_1 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad J_2 = \begin{pmatrix} A \\ D \\ 0 \\ -x_2 \\ -x_1 \end{pmatrix}.$$

The unprojection is then the solution of the equation

$$(4.2) \quad J_2(-1) = M_2 \begin{pmatrix} s_1 \\ s_2 \end{pmatrix},$$

where s_1, s_2 is the standard basis of $2\mathcal{O}$, the domain of M_2 , and -1 is the basis of \mathcal{O} , the domain of J_2 . (Note that s_1, s_2 both have a simple pole along the divisor F , as usual for unprojection variables.) That is, we consider \mathbb{C}^{11} with new variables s_1, s_2 and define $U \subset \mathbb{C}^{11}$ by the five equations (4.2). Of course, two of the equations are $x_1 = xs_1$ and $x_2 = xs_2$, so one would usually eliminate these two variables to get

$$U: (A + Bs_1 + Cs_2 = D + Es_1 + Fs_2 = 0) \subset \mathbb{C}^9.$$

There are two twists that make this applicable for us. First, the whole construction can be graded with weights chosen on the variables for which the Pfaffians are homogeneous. In that case, the resolutions are also graded, and one can take Proj of the rings appearing to give birational maps between varieties in wps. Second, the variables can be specialised to give calculations in lower dimensions; typical for us is to set A, \dots, F to be functions of 3 new variables together with the x variables so that we work with projective 3-folds.

Example No. 9: $X \subset \mathbb{P}^6(1, 1, 1, 3, 4, 5, 6)$ with

Degree = $5/6$, **Basket** = $\{\frac{1}{6}(1, 1, 5)\}$.

The link from the $\frac{1}{6}(1, 1, 5)$ point will be a Kawamata blowup, a flop, a hypersurface flip of type $(4, 1, 1, -3, -1; 2)$ followed by the contraction of a divisor to a point on a Fano 3-fold X' . The unprojection method computes the equations of X' from the syzygy matrix of X . In variables x, x_1, x_2, y, z, t, u , this matrix is

$$\begin{pmatrix} u & A_7 & B_5 & C_4 \\ & t & E_3 & F_2 \\ & & z & y \\ & & & x \end{pmatrix}$$

where A, \dots, F are general forms of the indicated degrees.

The projection $X \dashrightarrow Z$ from P_u works by eliminating u as usual. To continue the link, we introduce two new variables, s_1 and s_2 of weights 2 and 3 respectively, subject

to relations

$$\begin{pmatrix} A_7 & B_5 & C_4 \\ t & E_3 & F_2 \\ 0 & z & y \\ -z & 0 & x \\ -y & -x & 0 \end{pmatrix} \begin{pmatrix} 1 \\ s_1 \\ s_2 \end{pmatrix} = 0,$$

where the big matrix is the final 3 columns of the syzygy matrix. We use these equations to eliminate y , z and also t , so the resulting variety is

$$X': (A_7 + s_1 B_5 + s_2 C_4 = 0) \subset \mathbb{P}^4(1, 1, 1, 2, 3)$$

given by an equation of degree 7 in variables x, x_1, x_2, s_1, s_2 (after substituting for y, z, t).

Notice that X' contains the $s_1 s_2$ line $\mathbb{P}(2, 3)$ and that for general A it will have a single node in the interior of this line—this point is the image of the contracted divisor. Thus X' is very far from being a general element of the family of all such hypersurfaces. This is typical and already observed in [CM04]. Another point is that the model case of Type III unprojection in [Rei00] is a (rational) divisorial contraction to a line, so that this example exhibits something new.

4.3.4. End of the link II: numerical properties. We use the previous calculations to compute detailed information about the varieties at the end of the link in each of the four cases in Theorem 1.1. We use the notation of Theorem 1.1 throughout.

Proposition 4.11 (Divisorial contraction to a point). *If $a_1 < a_2 < a_3$ then the image of the contracted divisor is a singular point p' on a Fano 3-fold X' and the index of $p' \in X'$ is an integer multiple of $a_2 - a_1$.*

Proof. The map $\varphi: Y' \rightarrow X'$ is given by (multiples of) the linear system of the divisor F_2 and it contracts the divisor F_1 to a point $q \in X'$. Moreover, X' is a Fano 3-fold so denoting $A' = -K_{X'}$,

$$\varphi^*(A') = B + \frac{a'}{r'} F_1$$

where r' is the index of $q \in X'$ and a' is some positive integer. Since $\rho_{X'} = 1$, the divisor $\varphi^*(A')$ must be some positive multiple, possibly rational, of F_2 , say $\varphi^*(A') = \lambda F_2$. So since $F_2 = F_1 + (a_2 - a_1)B$ on Y' ,

$$\lambda F_1 + \lambda(a_2 - a_1)B = B + \frac{a'}{r'} F_1.$$

But B and F_1 are linearly independent in $\text{Pic}(Y')$, so $r' = a'(a_2 - a_1)$ is a positive integral multiple of $a_2 - a_1$. \square

Notice that, in the notation of the proof, if $F_1 \subset Y'$ is Cartier, then the index is exactly $a_2 - a_1$: in that case, we can use an argument of Kawamata [Kaw96] to show that $a' = 1$. Let $V \xrightarrow{g} Y' \rightarrow X'$ be a resolution of singularities, with composition denoted f . If $a' \neq 1$, then there is a divisor $G \subset V$ with discrepancy $1/r'$ over X' . But then, away from all exceptional divisors other than G and F_1 , $f^*(-K_{X'}) = -K_V - (a'/r')F_1 - (1/r')G$ while also

$$f^*(-K_{X'}) = g^*(-K_{Y'} - (a'/r')F_1) = -K_V - (a'/r')F_1 - mG$$

where $m \geq a'/r'$. Equating these two expressions shows that Y' does not have terminal singularities, which is not true. Kawakita's analysis of divisorial contractions [Kaw05] distinguishes between cases when F_1 is Cartier or not, but we have not carried out the additional calculations to see which cases occur.

Proposition 4.12 (Conic bundle). *If $a_1 = a_2 = a_3$ then the conic bundle $Y' \rightarrow \mathbb{P}^2$ has discriminant $\Delta \subset \mathbb{P}^2$ of degree*

$$\deg \Delta = 12 - a_1 \deg(X) + \left(\frac{a_1 + r}{r} \right) \left(\frac{1}{a(r-a)} \right).$$

Proof. By [Tak06], Lemma 3.1, the degree of the discriminant is equal to $12 - B^2F_1$ computed on Y' . We can equally well compute on Y , since the map $Y \dashrightarrow Y'$ is a B -flop. Computing on Y , we see that $AE^2 = A^2E = 0$ so

$$\begin{aligned} B^2F_1 &= \left(A - \frac{1}{r}E \right)^2 \left(a_1A - \left(\frac{a_1 + r}{r} \right) E \right) \\ &= a_1A^3 - \left(\frac{a_1 + r}{r^3} \right) E^3 \\ &= a_1A^3 - \left(\frac{a_1 + r}{r} \right) \left(\frac{1}{a(r-a)} \right) \end{aligned}$$

as required. \square

In fact, only the following three links from the 69 families result in a conic bundle:

No.2: $X \subset \mathbb{P}^6(1, 1, 1, 1, 1, 2)$ of degree $13/2$ and basket $\{\frac{1}{2}(1, 1, 1)\}$. So $a_1 = a_2 = a_3 = 1$ and the discriminant has degree $\deg \Delta = 12 - 13/2 + 3/2 = 7$. (The equations of X have degrees $2, 3, 3, 3, 3$ and $Y \dashrightarrow Y'$ flops 7 irreducible rational curves.)

No.3: $X \subset \mathbb{P}^6(1, 1, 1, 1, 1, 2, 3)$ of degree $14/3$ and basket $\{\frac{1}{3}(1, 1, 2)\}$. So $a_1 = a_2 = a_3 = 1$ and the discriminant has degree $\deg \Delta = 12 - 14/3 + 4/3 \times 1/2 = 8$. (The equations of X have degrees $3, 3, 4, 4, 4$ and $Y \dashrightarrow Y'$ flops 6 irreducible rational curves.)

No.8: $X \subset \mathbb{P}^6(1, 1, 1, 2, 2, 2, 2)$ of degree $5/2$ and basket $\{5 \times \frac{1}{2}(1, 1, 1)\}$. So $a_1 = a_2 = a_3 = 2$ and the discriminant has degree $\deg \Delta = 12 - 2 \times 5/2 + 4/2 = 9$. (The equations of X have degrees $4, 4, 4, 4, 4$ and $Y \dashrightarrow Y'$ flops 12 irreducible rational curves.)

Proposition 4.13 (Del Pezzo fibration). *If $a_1 = a_2 < a_3$ then the degree of fibres of the del Pezzo fibration $Y' \rightarrow \mathbb{P}^1$ is*

$$(-K_{F_1})^2 = \frac{1}{(a_3 - a_1)^2} \left(a_1 a_3^2 \deg(X) - \frac{(a_1 + r)(a_3 + r)^2}{ra(r-a)} + N \right)$$

where N is the number of curves flopped by $Y_1 \dashrightarrow Y'$, which is the same number N computed in Corollary 4.4.

Proof. Computing on Y_2 , we choose F_1 as a fibre to work on and find its degree as

$$(-K_{F_1})^2 = (B - F_1)|_{F_1}^2 = B^2F_1 = \frac{1}{(a_3 - a_1)^2} (F_3 - F_1)^2 F_1 = \frac{1}{(a_1 - a_3)^2} F_1 F_3^2,$$

where equalities come from $F_1^2 = 0$ on Y_2 . We compute this number on X .

The flip $Y_1 \dashrightarrow Y_2$ is an F_3 -flop, so the expression $F_1 F_3^2$ is the same whether computed on Y_1 or Y_2 . The flop $Y \dashrightarrow Y_1$ is a true B -flop, so the expression $F_1 F_3^2 = F_1(F_2 + (a_3 - a_2)B)F_3$ differs only in the term $F_1 F_2 F_3$ when computed on Y rather than Y_1 . This intersection defines exactly the reduced flopping curves, which are contracted to nonsingular points on X . So $F_1 F_3^2$ computed on Y_1 equals $F_1 F_3^2 + N$ computed on X .

Now $F_i = a_i B - E = a_i A - ((a_i + r)/r)E$, and so $AE^2 = A^2E = 0$ and $E^3 = r^2/(a(r-a))$ completes the calculation. \square

In fact, 11 of the links from the 69 families result in del Pezzo fibrations. We list them in Table 1.

No.	Fano 3-fold X	$p \in X$	A^3	a_3, a_2, a_1	#	$[n]$	d
4	$X_{3^4,4} \subset \mathbb{P}(1^5, 2^2)$	$\frac{1}{2}(1, 1, 1)$	5	2, 1, 1	8	[2]	4
6	$X_{4^2,5^2,6} \subset \mathbb{P}(1^4, 2, 3, 4)$	$\frac{1}{4}(1, 1, 3)$	11/4	2, 1, 1	7	[3]	3
7	$X_{3,4^3,5} \subset \mathbb{P}(1^4, 2, 2, 3)$	$\frac{1}{3}(1, 1, 2)$	19/6	2, 1, 1	7	[2]	3
10	$X_{4^3,5^2} \subset \mathbb{P}(1^3, 2^3, 3)$	$\frac{1}{2}(1, 1, 1)$	13/6	3, 2, 2	14	[3]	3
12	$X_{4,5,6^2,7} \subset \mathbb{P}(1^3, 2^2, 3, 5)$	$\frac{1}{5}(1, 2, 3)$	17/10	2, 1, 1	5	[2]	2
16	$X_{4,5^2,6^2} \subset \mathbb{P}(1^2, 2^3, 3^2)$	$\frac{1}{3}(1, 1, 2)$	7/6	3, 2, 2	11	[2]	2
17	$X_{6^2,7^2,8} \subset \mathbb{P}(1^2, 2^2, 3, 4, 5)$	$\frac{1}{5}(1, 1, 4)$	4/5	3, 2, 2	10	[4]	2
21	$X_{6^3,7^2} \subset \mathbb{P}(1^2, 2, 3^3, 4)$	$\frac{1}{5}(1, 1, 2)$	3/4	4, 3, 3	15	[4]	2
43	$X_{5,6^3,7} \subset \mathbb{P}(1^2, 2^2, 3^2, 4)$	$\frac{1}{3}(1, 1, 3)$	11/12	3, 2, 2	10	[3]	2
55	$X_{8,9,10^2,11} \subset \mathbb{P}(1, 2, 3^2, 4, 5, 7)$	$\frac{1}{7}(1, 2, 5)$	3/14	4, 3, 3	8	[2]	1
56	$X_{8,9^2,10^2} \subset \mathbb{P}(1, 2, 3, 4^2, 5^2)$	$\frac{1}{5}(1, 2, 3)$	1/5	5, 4, 4	11	[2]	1

TABLE 1. Sarkisov links that result in a del Pezzo fibration. The link comprises the Kawamata blowup of the Type I centre $p \in X$ followed by the standard flop of the indicated number of curves, completed by a flip to the total space of the fibration.

Notation: # denotes the number of flopped curves, A^3 is the degree of X , d is the degree of the del Pezzo fibre. The column labelled $[n]$ describes the flip: $[n]$ indicates the toric hypersurface flip of $n - 1$ irreducible \mathbb{P}^1 s meeting in the quotient singularity $\frac{1}{n}(1, 1, n - 1)$; in each case there is a single curve after the flip that passes through no singularities—compare the classification in [Bro99].

5. FINITE GENERATION AND RANK 2 LINKS

5.1. Finitely-generated extremal extractions. We give an elementary proof of

Theorem 5.1. *Let $f: Y \rightarrow X$ be a finitely-generated extremal extraction. Then the two ray game on Y determines a rank 2 link from X .*

In fact, this follows from [HK00], Proposition 2.9. Our proof is suited to making calculations in explicit situations, because we do not apply the variation of GIT directly to Y , but simply follow the birational transforms of Y through an ambient toric link. The main point is that the link continues to an end and doesn't fail because some contraction or flip doesn't exist. We use the ambient toric link to guarantee that such maps exist. A less important point, but one that is interesting from the point of view of calculation, is that the toric link contains only one link from X and not a sequence of links.

Proof. Let $f: X \rightarrow Y$ be a finitely-generated extremal extraction with exceptional divisor $E \subset Y$ from a Fano 3-fold X . Let $\mathcal{M} = \mathcal{M}(f)$ be the graded ring of f . By assumption, one can choose generators for \mathcal{M} and write $\mathcal{M} = \mathcal{N}/I$ for a doubly graded polynomial ring \mathcal{N} and a homogeneous ideal $I \subset \mathcal{N}$.

Claim 5.2. *The generators of \mathcal{M} can be chosen so that none lie in $H^0(Y, aA + bE)$ with both $a, b > 0$.*

First, the generating set must include a basis of $H^0(Y, E) \cong k$: for any $c, d > 0$, $H^0(Y, -cB + dE) = 0$ (since $(-cB + dE)C < 0$ for all curves $C \subset Y$ that miss E), so $H^0(Y, E)$ lies on a boundary edge of the cone Σ_{eff} . Now let $H = H^0(Y, aA + bE)$ be a graded piece of \mathcal{M} with $a, b > 0$. Since A is numerically trivial on E , $aA + bE$ is negative and the cohomology of

$$0 \rightarrow \mathcal{O}_Y(aA + (b-1)E) \rightarrow \mathcal{O}_Y(aA + bE) \rightarrow \mathcal{O}_E(aA + bE) \rightarrow 0$$

shows that any element of H is divisible by the generator of $H^0(Y, E)$. So elements of H are generated by those of $H^0(Y, E)$ and $H^0(Y, aA + (b-1)E)$, and repeating this process proves the claim.

For such a choice of generators, let

$$(5.1) \quad \begin{array}{ccccccc} & & \mathcal{Y}_0 & \dashrightarrow & \mathcal{Y}_1 & \dashrightarrow & \dots & \dashrightarrow & \mathcal{Y}_k & & \\ & \swarrow & & & \swarrow & & & & & \searrow & \\ \mathcal{Z}_0 & & & & \mathcal{Z}_1 & & & & & & \mathcal{Z}_{k+1} \end{array}$$

be the toric link arising from the grading on \mathcal{N} ; assume that the trivial left-hand map (that would come from the basis of $H^0(Y, E)$) is removed, so that $\mathcal{Z}_0 \leftarrow \mathcal{Y}_0$ is a birational toric blowup. Furthermore, all \mathcal{Y}_i have the same dimension, and we may assume that $\mathcal{Y}_k \rightarrow \mathcal{Z}_{k+1}$ is either a (birational) divisorial contraction or a map to a toric variety of lower dimension but not a point (again by omitting a trivial right-hand map if necessary).

Claim 5.3. *Without loss of generality, X embeds in \mathcal{Z}_0 and intersects the open toric stratum of \mathcal{Z}_0 nontrivially.*

Certainly X embeds into *some* quotient of $\mathcal{A} = \text{Spec } \mathcal{N}$, and the choice of generators mean that both X and \mathcal{Z}_0 are the quotients polarised by a character in the first nontrivial ray in Σ_{eff} , which is along multiples of A in each case. So $X \hookrightarrow \mathcal{Z}_0$.

If the image of this embedding misses the open stratum, then there are generators of \mathcal{N} that map to sections in \mathcal{M} that vanish on X . That is, we have chosen a generator $z_i \in \mathcal{N}$ that is contained in the ideal I . Such generators are redundant and can be omitted (although we must recompute the sequence (5.1)).

Claim 5.4. *The map $X \leftarrow Y$ embeds in the map $\mathcal{Z}_0 \leftarrow \mathcal{Y}_0$.*

Similarly, using (a suitable multiple of) $A - \varepsilon E$ for small $\varepsilon > 0$ shows that $Y \hookrightarrow \mathcal{Y}_0$. The maps between quotients are the same since they are defined by (the restriction of) the same linear system. (There may be many generators lying between A and B in Σ_{eff} , and some choices of polarisation defining Y will embed it naturally in other \mathcal{Y}_i ; nevertheless characters very close to A polarise the quotients of both Y and \mathcal{Y}_0 .)

Claim 5.5. *The birational images $Y_i \subset \mathcal{Y}_i$ of Y in each \mathcal{Y}_i are equal to $\text{Proj } R_i$, where $R_i = \bigoplus_{j \geq 0} H^0(Y, j(m_i B + n_i E))$ for some integers $m_i > 0$ and $n_i \in \mathbb{Z}$. Moreover, each of these varieties Y_i is normal.*

Each \mathcal{Y}_i is Proj of some \mathbb{N} -graded subring of \mathcal{N} given by the positive multiples of polynomials in \mathcal{N} of bidegree (m_i, n_i) , for any fixed integer pair (m_i, n_i) lying in the chamber of Σ_{eff} corresponding to \mathcal{Y}_i . The graded subring of \mathcal{M} comprising the summands of these same degrees defines the image $Y_i \subset \mathcal{Y}_i$ of Y . Since Y meets the open torus of \mathcal{Y}_0 , and since all transformations occurring in (5.1) take place on toric strata, each Y_i is birational to Y . (We also denote $Y_0 = Y$.) The Y_i are normal because they are Proj of the ring of all multiples of an ample divisor; this is Zariski projective normalisation, as in [Rei87], for example.

Similarly, one can define varieties $Z_i \subset \mathcal{Z}_i$. Again, these will be 3-folds birational to Y , with the possible exception of Z_{k+1} —in fact, Z_{k+1} is birational to Y if and only if $\mathcal{Y}_k \rightarrow \mathcal{Z}_{k+1}$ is birational, again because birationality is detected on the intersection with the open toric stratum. The result is an array of varieties and morphisms similar to (5.1) but composed of the Y_i and Z_i . The theorem then follows by showing that this array is a complete 2-ray link from X .

Inductively, we assume that Y_i is \mathbb{Q} -factorial with $\rho_{Y_i} = 2$; these properties hold for $Y = Y_0$.

Claim 5.6. *If $i < k-1$ and the morphism $Y_i \rightarrow Z_{i+1}$ is an isomorphism, then $Z_{i+1} \leftarrow Y_{i+1}$ is also an isomorphism.*

Since Z_{i+1} is isomorphic to Y_i , $aB - bE$ is an ample \mathbb{Q} -Cartier divisor on Z_{i+1} for some $a, b > 0$. So the same is true for $aB - (b + \varepsilon)E$ for sufficiently small and rational $\varepsilon > 0$. Multiples of this divisor embed $Z_{i+1} \hookrightarrow \mathcal{Y}_{i+1}$. In particular, this map is an isomorphism to its image when restricted to the open toric stratum, and so it is the birational map $Z_{i+1} \dashrightarrow Y_{i+1}$; but this is an isomorphism, which is the claim.

Claim 5.7. *Suppose that $Y_i \rightarrow Z_{i+1}$ is not an isomorphism and is small—that is, no divisor is contracted. Then $i < k$, $Z_{i+1} \leftarrow Y_{i+1}$ is small, and the birational map $Y_i \dashrightarrow Y_{i+1}$ is a flip, flop or antiflop that corresponds to a step in a 2-ray link.*

Let $D = m_i B + n_i E$ be an ample divisor on Z_{i+1} . (This divisor will polarise the resulting flip.) Then on Y_i , the divisor $D + \varepsilon E$ is positive on the contracted locus for small $\varepsilon > 0$ (because it is at the level of toric varieties in (5.1)) and

$$Y_i \cong \mathrm{Proj}_{Z_{i+1}} \bigoplus_{n \geq 0} H^0(Z_{i+1}, nN(D + \varepsilon E))$$

for suitable fixed (and divisible) $N > 0$. Since it is a quotient of some subring of \mathcal{N} , the $\mathcal{O}_{Z_{i+1}}$ -algebra $\bigoplus_{n \geq 0} H^0(Z_{i+1}, nN(D - \varepsilon E))$ is finitely generated, and its $\mathrm{Proj}_{Z_{i+1}}$ is the required flip and it embeds the flip in \mathcal{Y}_{i+1} . In particular, Y_{i+1} is also \mathbb{Q} -factorial with $\rho_{Y_{i+1}} = 2$.

Claim 5.8. *Suppose that $Y_i \rightarrow Z_{i+1}$ is birational and contracts a divisor. Then this map is an extremal divisorial contraction that completes the link and all further maps $Y_j \rightarrow Z_{j+1}$ are isomorphisms.*

Since $\rho_{Y_i} = 2$ and Z_{i+1} is projective, $\rho(Z_{i+1}) = 1$ and this contraction is extremal and Z_{i+1} is also \mathbb{Q} -factorial. So if $i < k$, then $Z_{i+1} \leftarrow Y_{i+1}$ must be an isomorphism, as in Claim 5.6 above. Now, inductively for any $j > i$, $Z_{i+1} \cong Y_j$ and $Y_j \rightarrow Z_{j+1}$ is an isomorphism (because Y_j has $\rho = 1$); and again, as in Claim 5.6 above, the ‘flip’ of this is an isomorphism.

Claim 5.9. *Suppose that $Y_i \rightarrow Z_{i+1}$ is not birational. Then this map is an extremal fibration that completes the link and $i = k$.*

The map is certainly extremal, as usual, since $\rho_{Y_i} = 2$, and so it does complete the link. Since Y_i meets the big torus of \mathcal{Y}_i , this map can only fail to be birational if the toric map $\mathcal{Y}_i \rightarrow \mathcal{Z}_{i+1}$ is not birational. This can only occur if $i = k$. \square

5.2. Missing generators. The calculations above apply to many other situations. It can happen that one does not compute the full graded ring \mathcal{M} at first attempt. For example, in the Del Pezzo fibrations of Brown–Corti–Zucconi [BCZ04], links are started without enough generators of $\mathcal{M}(f)$ to realise them. The typical indication of missing generators is when a flip of Y_i is not made by the ambient toric flip—unprojection is the key to identifying the missing generator. This is explained in [BCZ04] section 4.4.3.

5.2.1. *Elliptic and K3 fibrations of Ryder.* Ryder [Ryd02] (and similar results in [Ryd06]) shows that the general Fano 3-fold $X = X_{22} \subset \mathbb{P}(1, 1, 3, 7, 11)$ is birational to variety fibred over \mathbb{P}^1 in K3 surfaces by computing a 2-ray link.

Given the matrix of weights of a $G = \mathbb{C}^* \times \mathbb{C}^*$ action on \mathbb{C}^6

$$\begin{pmatrix} 0 & 3 & 11 & 7 & 1 & 1 \\ 1 & 1 & 3 & 1 & 0 & 0 \end{pmatrix},$$

consider the general hypersurface of bidegree $(22, 6)$. One can check that the quotient linearised by $\chi = (3, 1)$ is the Fano X . The link proceeds by blowing up the $\frac{1}{3}(1, 1, 2)$ quotient singularity $X \leftarrow Y$. The first toric modification (a toric flip with base the quotient \mathbb{C}^6/G linearised by $(11, 3)$) happens away from Y , because, if z is the variable of weight $(11, 3)$, the equation of Y includes a z^2 term. But the second toric modification induces an antiflip of hypersurface type $(-7, -4, -10, 1, 1; -20)$. Writing down equations shows that the flipping curve passes through the two singularities $\frac{1}{7}(1, 3, 4), \frac{1}{2}(1, 1, 1) \in Y$. The antiflip $Y \dashrightarrow Y'$ results in a nonsingular variety that has a morphism to \mathbb{P}^1 . Again, the equations show that the fibre is a hypersurface of degree 6 in $\mathbb{P}(1, 1, 1, 3)$; the general fibre is nonsingular so is a K3 surface.

The weight matrix used here could be deduced using calculations similar to [CPR00] section 4.10, computing the vanishing of the coordinates on $\mathbb{P}(1, 1, 3, 7, 11)$ on the exceptional divisor of $X \leftarrow Y$. However, in this case it is likely that we have not calculated the full graded ring \mathcal{M} , since there are presumably other generators coming from the linear systems of $10B + 2E, 13B + 3E, 16B + 4E, 19B + 5E$. But we already have enough sections to compute the link, so we do not look for these extra generators.

REFERENCES

- [ABR02] Selma Altınok, Gavin Brown, and Miles Reid. Fano 3-folds, K3 surfaces and graded rings. In *Topology and geometry: commemorating SISTAG*, volume 314 of *Contemp. Math.*, pages 25–53. Amer. Math. Soc., Providence, RI, 2002.
- [AKMW02] Dan Abramovich, Kalle Karu, Kenji Matsuki, and Jarosław Włodarczyk. Torification and factorization of birational maps. *J. Amer. Math. Soc.*, 15(3):531–572 (electronic), 2002.
- [Alt98] Selma Altınok. *Graded rings corresponding to polarised K3 surfaces and \mathbb{Q} -Fano 3-folds*. PhD thesis, University of Warwick, 1998.
- [BCZ04] Gavin Brown, Alessio Corti, and Francesco Zucconi. Birational geometry of 3-fold Mori fibre spaces. In *The Fano Conference*, pages 235–275. Univ. Torino, Turin, 2004.
- [BDK⁺] Gavin Brown, Sarah Davis, Alexander Kasprzyk, Michael Kerber, Olof Sisask, and Stephen Tawn. The graded ring database webpage. www.kent.ac.uk/ims/grdb/, 2004–6.
- [BRK] Gavin Brown, Miles Reid, and Michael Kerber. K3 surfaces and Fano 3-folds, Tom and Jerry. In preparation.
- [Bro99] Gavin Brown. Flips arising as quotients of hypersurfaces. *Math. Proc. Cambridge Philos. Soc.*, 127(1):13–31, 1999.
- [Bro07] Gavin Brown. A database of polarised K3 surfaces. *Exp. Math.*, 2007. 28pp. To appear.
- [CM04] Alessio Corti and Massimiliano Mella. Birational geometry of terminal quartic 3-folds. I. *Amer. J. Math.*, 126(4):739–761, 2004.
- [Cor00] Alessio Corti. Singularities of linear systems and 3-fold birational geometry. In *Explicit birational geometry of 3-folds*, volume 281 of *London Math. Soc. Lecture Note Ser.*, pages 259–312. Cambridge Univ. Press, Cambridge, 2000.
- [Cox95] David A. Cox. The homogeneous coordinate ring of a toric variety. *J. Algebraic Geom.*, 4(1):17–50, 1995.
- [CPR00] Alessio Corti, Aleksandr Pukhlikov, and Miles Reid. Fano 3-fold hypersurfaces. In *Explicit birational geometry of 3-folds*, volume 281 of *London Math. Soc. Lecture Note Ser.*, pages 175–258. Cambridge Univ. Press, Cambridge, 2000.

- [CR00] Alessio Corti and Miles Reid, editors. *Explicit birational geometry of 3-folds*, volume 281 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2000.
- [CR02] Alessio Corti and Miles Reid. Weighted Grassmannians. In *Algebraic geometry*, pages 141–163. de Gruyter, Berlin, 2002.
- [CS05] Alessio Corti and Ivan Smith. Conifold transitions and Mori theory. *Math. Res. Lett.*, 12(5-6):767–778, 2005.
- [DH98] Igor V. Dolgachev and Yi Hu. Variation of geometric invariant theory quotients. *Inst. Hautes Études Sci. Publ. Math.*, (87):5–56, 1998.
- [Dol03] Igor Dolgachev. *Lectures on invariant theory*, volume 296 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2003.
- [Gri01] M. M. Grinenko. On fibrations into del Pezzo surfaces. *Mat. Zametki*, 69(4):550–565, 2001.
- [HK00] Yi Hu and Sean Keel. Mori dream spaces and GIT. *Michigan Math. J.*, 48:331–348, 2000.
- [IF00] A. R. Iano-Fletcher. Working with weighted complete intersections. In *Explicit birational geometry of 3-folds*, volume 281 of *London Math. Soc. Lecture Note Ser.*, pages 101–173. Cambridge Univ. Press, Cambridge, 2000.
- [Kaw88] Yujiro Kawamata. Crepant blowing-up of 3-dimensional canonical singularities and its application to degenerations of surfaces. *Ann. of Math. (2)*, 127(1):93–163, 1988.
- [Kaw96] Yujiro Kawamata. Divisorial contractions to 3-dimensional terminal quotient singularities. In *Higher-dimensional complex varieties (Trento, 1994)*, pages 241–246. de Gruyter, Berlin, 1996.
- [Kaw02] Masayuki Kawakita. Divisorial contractions in dimension three which contract divisors to compound A_1 points. *Compositio Math.*, 133(1):95–116, 2002.
- [Kaw05] Masayuki Kawakita. Three-fold divisorial contractions to singularities of higher indices. *Duke Math. J.*, 130(1):57–126, 2005.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998.
- [Kol92] János Kollár, editor. *Flips and abundance for algebraic threefolds*. Société Mathématique de France, Paris, 1992.
- [Pap06] Stavros Argyrios Papadakis. Type II unprojection. *J. Algebraic Geom.*, 15(3):399–414, 2006.
- [Rei83] Miles Reid. Decomposition of toric morphisms. In *Arithmetic and geometry, Vol. II*, volume 36 of *Progr. Math.*, pages 395–418. Birkhäuser Boston, Boston, MA, 1983.
- [Rei87] Miles Reid. Young person’s guide to canonical singularities. In *Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985)*, volume 46 of *Proc. Sympos. Pure Math.*, pages 345–414. Amer. Math. Soc., Providence, RI, 1987.
- [Rei92] Miles Reid. What is a flip? Unpublished manuscript of Utah seminar, 1992.
- [Rei97] Miles Reid. Chapters on algebraic surfaces. In *Complex algebraic geometry (Park City, UT, 1993)*, volume 3 of *IAS/Park City Math. Ser.*, pages 3–159. Amer. Math. Soc., Providence, RI, 1997.
- [Rei00] Miles Reid. Graded rings and birational geometry. Proc. of algebraic geometry symposium (Kinosaki), pages 1–72. 2000.
- [Rei02] Miles Reid. Quasi-Gorenstein unprojection. Unpublished manuscript, 2002.
- [Ryd02] Daniel Ryder. *Elliptic and K3 fibrations birational to Fano 3-fold weighted hypersurfaces*. PhD thesis, University of Warwick, 2002.
- [Ryd06] Daniel Ryder. Classification of elliptic and K3 fibrations birational to some \mathbb{Q} -Fano 3-folds. *J. Math. Sci. Univ. Tokyo*, 13(1):13–42, 2006.
- [Tak06] Hiromichi Takagi. Classification of primary \mathbb{Q} -Fano threefolds with anti-canonical Du Val K3 surfaces. I. *J. Algebraic Geom.*, 15(1):31–85, 2006.
- [Tha96] Michael Thaddeus. Geometric invariant theory and flips. *J. Amer. Math. Soc.*, 9(3):691–723, 1996.