

Edges not in any monochromatic copy of a fixed graph

Hong Liu

Oleg Pikhurko

Maryam Sharifzadeh *

July 10, 2018

Abstract

For a sequence $(H_i)_{i=1}^k$ of graphs, let $\text{nim}(n; H_1, \dots, H_k)$ denote the maximum number of edges not contained in any monochromatic copy of H_i in colour i , for any colour i , over all k -edge-colourings of K_n .

When each H_i is connected and non-bipartite, we introduce a variant of Ramsey number that determines the limit of $\text{nim}(n; H_1, \dots, H_k) / \binom{n}{2}$ as $n \rightarrow \infty$ and prove the corresponding stability result. Furthermore, if each H_i is what we call *homomorphism-critical* (in particular if each H_i is a clique), then we determine $\text{nim}(n; H_1, \dots, H_k)$ exactly for all sufficiently large n . The special case $\text{nim}(n; K_3, K_3, K_3)$ of our result answers a question of Ma.

For bipartite graphs, we mainly concentrate on the two-colour symmetric case (i.e., when $k = 2$ and $H_1 = H_2$). It is trivial to see that $\text{nim}(n; H, H)$ is at least $\text{ex}(n, H)$, the maximum size of an H -free graph on n vertices. Keevash and Sudakov showed that equality holds if H is the 4-cycle and n is large; recently Ma extended their result to an infinite family of bipartite graphs. We provide a larger family of bipartite graphs for which $\text{nim}(n; H, H) = \text{ex}(n, H)$. For a general bipartite graph H , we show that $\text{nim}(n; H, H)$ is always within a constant additive error from $\text{ex}(n, H)$, i.e., $\text{nim}(n; H, H) = \text{ex}(n, H) + O_H(1)$.

1 Introduction

Many problems of extremal graph theory ask for (best possible) conditions that guarantee the existence of a given ‘forbidden’ graph. Two prominent examples of this kind are the Turán function and Ramsey numbers. Recall that, for a graph H and an integer n , the *Turán function* $\text{ex}(n, H)$ is the maximum size of an n -vertex H -free graph. Let K_t denote the complete graph on t vertices. The famous theorem of Turán [48] states that the unique maximum K_{r+1} -free graph of order n is the *Turán graph* $T(n, r)$, the complete balanced r -partite graph. Thus $\text{ex}(n, K_{k+1}) = t(n, r)$, where we denote $t(n, r) := e(T(n, r))$. For a sequence a_1, \dots, a_k of integers, the *Ramsey number* $R(a_1, \dots, a_k)$ is the minimum R such that for every edge-colouring of K_R with colours from $[k] := \{1, \dots, k\}$, there is a colour- i copy of K_{a_i} for some $i \in [k]$. The fact that R exists (i.e., is finite) was first established by Ramsey [39] and then independently rediscovered by Erdős and Szekeres [16]. Both of these problems motivated a tremendous amount of research, see e.g. the recent surveys by Conlon, Fox and Sudakov [4], Füredi and Simonovits [20], Keevash [28], Radziszowski [38] and Sudakov [44].

*Mathematics Institute and DIMAP, University of Warwick, Coventry, CV4 7AL, UK. Email addresses: {h.liu.9, o.pikhurko, m.sharifzadeh}@warwick.ac.uk. H.L. was supported by the Leverhulme Trust Early Career Fellowship ECF-2016-523. O.P. was supported by ERC grant 306493 and EPSRC grant EP/K012045/1. M.Sh. was supported by ERC grant 306493 and Marie Curie Individual Fellowship 752426.

A far-reaching generalisation is to ask for the number of guaranteed forbidden subgraphs. For the Turán function this gives the famous *Erdős-Rademacher problem* that goes back to Rademacher (1941; unpublished): what is the minimum number of copies of H in a graph of given order n and size $m > \text{ex}(n, H)$? This problem was revived by Erdős [8, 9] in the 1950–60s. Since then it continues to be a very active area of research, for some recent results see e.g. [3, 5, 6, 26, 30, 32, 34, 35, 36, 40, 41, 45]. The analogous question for Ramsey numbers, known as the *Ramsey multiplicity problem*, was introduced by Erdős [10] in 1962 and is wide open, see e.g. [2, 7, 17, 21, 27, 43, 46, 47].

A less studied but still quite natural question is to maximise the number of edges that do not belong to any forbidden subgraph. Such problems in the Turán context (where we are given the order n and the size $m > \text{ex}(n, H)$ of a graph G) were studied in [13, 19, 22, 23]. In the Ramsey context, a problem of this type seems to have been first posed by Erdős, Rousseau, and Schelp (see [12, Page 84]). Namely, they considered the maximum number of edges not contained in any monochromatic triangle in a 2-edge-colouring of K_n . Also, Erdős [12, Page 84] wrote that “*many further related questions can be asked*”. Such questions will be the focus of this paper.

Let us provide a rather general definition. Suppose that we have fixed a sequence of graphs H_1, \dots, H_k . For a k -edge-colouring ϕ of K_n , let $\text{NIM}(\phi)$ consist of all *NIM-edges*, that is, those edges of K_n that are not contained in any colour- i copy of H_i for any $i \in [k]$. In other words, $\text{NIM}(\phi)$ is the complement (with respect to $E(K_n)$) of the union over $i \in [k]$ of the edge-sets of H_i -subgraphs of colour- i . Define

$$\text{nim}(n; H_1, \dots, H_k) := \max_{\phi: E(K_n) \rightarrow [k]} |\text{NIM}(\phi)|,$$

to be the maximum possible number of NIM-edges in a k -edge-colouring of K_n . If all H_i 's are the same graph H , we will write $\text{nim}_k(n; H)$ instead. Note that for $k = 2$ by taking one colour-class to be a maximum H -free graph, we have $\text{nim}_2(n; H) \geq \text{ex}(n, H)$. In ([12, Page 84]), Erdős mentioned that together with Rousseau and Schelp, they showed that in fact

$$\text{nim}_2(n; H) = \text{ex}(n, H), \quad \text{for all } n \geq n_0(H), \tag{1}$$

when $H = K_3$ is the triangle. As observed by Alon (see [29, Page 42]), this also follows from an earlier paper of Pyber [37]. Keevash and Sudakov [29] showed that (1) holds when H is an arbitrary clique K_t (or, more generally, when H is *edge-colour-critical*, that is, the removal of some edge $e \in E(H)$ decreases the chromatic number) as well as when H is the 4-cycle C_4 (and $n \geq 7$). They [29, Problem 5.1] also posed the following problem.

Problem 1.1 (Keevash and Sudakov [29]). *Does (1) hold for every graph H ?*

In a recent paper, Ma [33] answered Problem 1.1 in the affirmative for the infinite family of reducible bipartite graphs, where a bipartite graph H is called *reducible* if it contains a vertex $v \in V(H)$ such that $H - v$ is connected and $\text{ex}(n, H - v) = o(\text{ex}(n, H))$ as $n \rightarrow \infty$. Ma [33] also studied the case of $k \geq 3$ colours and raised the following question.

Question 1.2 (Ma [33]). *Is it true that $\text{nim}_3(n; K_3) = t(n, 5)$?*

The lower bound in Question 1.2 follows by taking a blow-up of a 2-edge-colouring of K_5 without a monochromatic triangle, and assigning the third colour to all pairs inside a part.

1.1 Non-bipartite case

In order to state some of our results, we have to introduce the following variant of Ramsey number. Given a set X , denote by $\binom{X}{i}$ (resp. $\binom{X}{\leq i}$), the set of all subsets of X of size i (resp. at most i). Given two graphs H and G , a (not necessarily injective) map $\phi : V(H) \rightarrow V(G)$ is a *homomorphism* if it preserves all adjacencies, i.e. $\phi(u)\phi(v) \in E(G)$ for every $uv \in E(H)$, and we say that G is a *homomorphic copy* of H .

Definition 1.3. Given a sequence of graphs $\mathcal{H} = (H_1, \dots, H_k)$, denote by $r^*(H_1, \dots, H_k)$ the maximum integer r^* such that there exists a colouring $\xi : \binom{[r^*]}{\leq 2} \rightarrow [k]$ such that

(P1) the restriction of ξ to $\binom{[r^*]}{2}$ is (H_1, \dots, H_k) -homomorphic-free (that is, for each $i \in [k]$ there is no edge-monochromatic homomorphic copy of H_i in the i -th colour);

(P2) for every distinct $i, j \in [r^*]$ we have $\xi(\{i, j\}) \neq \xi(\{i\})$, that is, we forbid a pair having the same colour as one of its points.

For any $r' \leq r^*$, we will call a colouring $\xi : \binom{[r']}{\leq 2} \rightarrow [k]$ feasible (with respect to (H_1, \dots, H_k)) if it satisfies both (P1) and (P2). We say that (H_1, \dots, H_k) is nice if every feasible colouring $\xi : \binom{[r^*(H_1, \dots, H_k)]}{\leq 2} \rightarrow [k]$ assigns the same colour to all singletons.

Note that the colour assigned by ξ to the empty set $\emptyset \in \binom{[r^*]}{\leq 2}$ does not matter. Note also that when $k = 2$, due to (P2), a feasible colouring should use the same colour on all singletons. Thus, $r^*(H_1, H_2) = \max\{\chi(H_1), \chi(H_2)\} - 1$. If we ignore (P2), then we obtain the following variant of Ramsey number that was introduced by Burr, Erdős and Lovász [1]. Let $r_{\text{hom}}(H_1, \dots, H_k)$ be the *homomorphic-Ramsey number*, that is the maximum integer r such that there exists an (H_1, \dots, H_k) -homomorphic-free colouring $\xi : \binom{[r]}{2} \rightarrow [k]$. We remark that for the homomorphic-Ramsey number, the colours of vertices do not play a role. When all H_i 's are cliques, this Ramsey variant reduces to the classical graph Ramsey problem:

$$r_{\text{hom}}(K_{a_1}, \dots, K_{a_k}) = R(a_1, \dots, a_k) - 1. \quad (2)$$

There are some further relations to r^* . For example, by assigning the same colour i to all singletons and using the remaining $k - 1$ colours on pairs, one can see that

$$r^*(H_1, \dots, H_k) \geq \max_{i \in [k]} r_{\text{hom}}(H_1, \dots, H_{i-1}, H_{i+1}, \dots, H_k). \quad (3)$$

If some H_i is bipartite, then the problem of r^* reduces to r_{hom} . Indeed, as K_2 is a homomorphic copy of any bipartite graph, when some H_i is bipartite, no feasible colouring ξ can use colour i on any pair. Consequently, we have equality in (3). This is one of the reasons why we restrict to non-bipartite H_i in this section.

It would be interesting to know if (3) can be strict. We conjecture that if all H_i 's are cliques then there is equality in (3) and, furthermore, every optimal colouring uses the same colour on all singletons:

Conjecture 1.4. For any integers $3 \leq a_1 \leq \dots \leq a_k$, $(K_{a_1}, \dots, K_{a_k})$ is nice. In particular, $r^*(K_{a_1}, \dots, K_{a_k}) = R(K_{a_2}, \dots, K_{a_k}) - 1$.




Figure 1: A feasible colouring of K_4 with respect to (C_5, C_5, C_5) , with two different colours on vertices.

It is worth noting that not all k -tuples are nice. For example, it is easy to show that $r^*(C_5, C_5, C_5) = r_{\text{hom}}(C_5, C_5) = 4$, where C_i denotes the cycle of length i , while Figure 1 shows a feasible colouring of $\binom{[4]}{\leq 2}$ assigning two different colours to singletons.

Our first result shows that this new variant plays a similar role for the function $\text{nim}(\cdot)$ as the chromatic number in the Erdős-Simonovits-Stone Theorem [15, 14].

Theorem 1.5. *Let H_i be a non-bipartite graph, $i \in [k]$, and let $r^* := r^*(H_1, \dots, H_k)$. For every $\varepsilon > 0$, we have that, for all large n ,*

$$\text{nim}(n; H_1, \dots, H_k) \leq \left(1 - \frac{1}{r^*}\right) \frac{n^2}{2} + \varepsilon n^2. \quad (4)$$

Furthermore, if each H_i is connected or there exists a feasible colouring of $\binom{[r^*]}{\leq 2}$ with k colours such that all singletons have the same colour, then we have $\text{nim}(n; H_1, \dots, H_k) \geq t(n, r^*)$.

We also obtain the following stability result stating that if the number of NIM-edges is close to the bound in (4), then the NIM-graph is close to a Turán graph. Let the *edit distance* between graphs G and H of the same order be

$$\delta_{\text{edit}}(G, H) := \min_{\sigma} |E(G) \Delta \sigma(E(H))|, \quad (5)$$

where the minimum is taken over all bijections $\sigma : V(H) \rightarrow V(G)$. In other words, $\delta_{\text{edit}}(G, H)$ is the minimum number of adjacency edits needed to make G and H isomorphic.

Theorem 1.6. *For any non-bipartite graphs H_i , $i \in [k]$, and any constant $\varepsilon > 0$, there exists $\delta > 0$ such that the following holds for sufficiently large n . If the number of NIM-edges of some $\phi : \binom{[n]}{2} \rightarrow [k]$ satisfies*

$$\text{nim}(\phi; H_1, \dots, H_k) \geq \left(1 - \frac{1}{r^*}\right) \frac{n^2}{2} - \delta n^2,$$

then $\delta_{\text{edit}}(G^{\text{nim}}, T(n, r^*)) \leq \varepsilon n^2$, where $r^* := r^*(H_1, \dots, H_k)$ and G^{nim} is the NIM-graph of ϕ , i.e., the spanning subgraph with edge set $\text{NIM}(\phi)$.

Our next theorem shows that if Conjecture 1.4 is true, then this would determine the exact value of $\text{nim}(\cdot)$ for a rather large family of graphs, including cliques. We call a graph H *homomorphism-critical* if it satisfies the following. If F is a minimal homomorphic copy of H , i.e. no proper subgraph of F is a homomorphic copy of H , then for any edge $uv \in E(F)$, there exists a homomorphism $g : V(H) \rightarrow V(F)$ such that $|g^{-1}(u)| = |g^{-1}(v)| = 1$, i.e. the pre-image sets of u and v are singletons.

For example, all complete multipartite graphs with at least two parts of size 1 are homomorphism-critical. A simple consequence of this property is the following. As F is minimal, it does not have any isolated vertices. Therefore, for any vertex $v \in V(F)$, there exists a homomorphism $g : V(H) \rightarrow V(F)$ such that $|g^{-1}(v)| = 1$.

Theorem 1.7. *Let (H_1, \dots, H_k) be a nice sequence of non-bipartite graphs such that each H_i is homomorphism-critical. Then for sufficiently large n ,*

$$\text{nim}(n; H_1, \dots, H_k) = t(n, r^*),$$

where $r^* := r^*(H_1, \dots, H_k)$. Additionally, the NIM-graph of every extremal colouring is isomorphic to $T(n, r^*)$.

In the following theorems, we prove Conjecture 1.4 for $k = 3$, and for $a_1 = \dots = a_4 = 3$ when $k = 4$.

Theorem 1.8. *For all integers $3 \leq a_1 \leq a_2 \leq a_3$, $(K_{a_1}, K_{a_2}, K_{a_3})$ is nice. In particular,*

$$r^*(K_{a_1}, K_{a_2}, K_{a_3}) = R(a_2, a_3) - 1.$$

Theorem 1.9. *We have that (K_3, K_3, K_3, K_3) is nice. In particular,*

$$r^*(K_3, K_3, K_3, K_3) = R(3, 3, 3) - 1 = 16.$$

The following is an immediate corollary of Theorems 1.7, 1.8 and 1.9. In particular, the special case $a_1 = a_2 = a_3 = 3$ answers Question 1.2 affirmatively.

Corollary 1.10. *Let $3 \leq a_1 \leq a_2 \leq a_3$ be integers. Then for sufficiently large n ,*

$$\text{nim}(n; K_{a_1}, K_{a_2}, K_{a_3}) = t(n, R(a_2, a_3) - 1),$$

$\text{nim}_4(n; K_3) = t(n, 16)$, and the NIM-graph of every extremal colouring is the corresponding Turán graph. \square

1.2 Bipartite graphs

For bipartite graphs, we will provide a new family for which Problem 1.1 has a positive answer. Let us call an h -vertex graph H *weakly-reducible* if there exist $n_0 \in \mathbb{N}$ and a vertex $v \in V(H)$ such that $\text{ex}(n, H - v) < \text{ex}(n, H) - 2^{2h^2}n$ for all $n \geq n_0$. (The function 2^{2h^2} comes from the proof and we make no attempt to optimise it.) Note that the family of weakly-reducible graphs includes all reducible graphs except the path of length 2 and this inclusion is strict. For example, for integers $t > s \geq 2$, the disjoint union of the complete bipartite graphs $K_{2,t}$ and $K_{2,s}$ is weakly-reducible but not reducible; this can be easily deduced from the result of Füredi [18] that $\text{ex}(n, K_{2,k}) = (\sqrt{k}/2 + o(1))n^{3/2}$ for any fixed $k \geq 2$ as $n \rightarrow \infty$.

Theorem 1.11. *Let H be a weakly-reducible bipartite graph and n be sufficiently large. Then*

$$\text{nim}_2(n; H) = \text{ex}(n, H).$$

Furthermore, every extremal colouring has one of its colour classes isomorphic to a maximum H -free graph of order n .

For a general bipartite graph H , we give in the following two theorems a weaker bound with an additive constant error term, namely,

$$\text{nim}_2(n; H) \leq \text{ex}(n, H) + O_H(1).$$

This provides more evidence towards Problem 1.1.

Theorem 1.12. *Let H be a bipartite graph on at most h vertices containing at least one cycle. Then for sufficiently large n ,*

$$\text{nim}_2(n; H) \leq \text{ex}(n, H) + h^2.$$

For more than 2 colours, we obtain an asymptotic result for trees. Fix a tree T , by taking random overlays of $k-1$ copies of extremal T -free graphs, we see that $\text{nim}_k(n; T) \geq (k-1)\text{ex}(n, T) - O_{k,|T|}(1)$ (this construction is from Ma [33]). We prove that this lower bound is asymptotically true.

Theorem 1.13. *Let T be a forest with h vertices. If $k = 2$ or if T is a tree, then there exists a constant $C := C(k, h)$ such that, for all sufficiently large n ,*

$$|\text{nim}_k(n; T) - (k-1)\text{ex}(n, T)| \leq C(k, h).$$

Organisation of the paper. We first introduce some tools in Section 2. Then in Section 3, we will prove Theorems 1.11, 1.12, and 1.13. In Section 4, we will prove Theorems 1.5 and 1.6. We will present the proof for Theorem 1.7 in Section 5 and the proofs of Theorems 1.8 and 1.9 in Section 6. Finally, in Section 7 we give some concluding remarks.

2 Preliminaries

In this section, we recall and introduce some notation and tools. Recall that $[m] := \{1, 2, \dots, m\}$ and $\binom{X}{i}$ (resp. $\binom{X}{\leq i}$) denotes the set of all subsets of a set X of size i (resp. at most i). We also use the term i -set for a set of size i . We may abbreviate a singleton $\{x\}$ (resp. a pair $\{x, y\}$) as x (resp. xy). If we claim, for example, that a result holds whenever $1 \gg a \gg b > 0$, this means that there are a constant $a_0 \in (0, 1)$ and a non-decreasing function $f : (0, 1) \rightarrow (0, 1)$ (that may depend on any previously defined constants or functions) such that the result holds for all $a, b \in (0, 1)$ with $a \leq a_0$ and $b \leq f(a)$. We may omit floors and ceilings when they are not essential.

Let $G = (V, E)$ be a graph. Its *order* is $v(G) := |V|$ and its *size* is $e(G) := |E|$. The *complement* of G is $\overline{G} := (V, \binom{V}{2} \setminus E)$. The chromatic number of G is denoted by $\chi(G)$. For $U \subseteq V$, let $G[U] := (U, \{xy \in E : x, y \in U\})$ denote the subgraph of G induced by U . Also, denote

$$\begin{aligned} N_G(v, U) &:= \{u \in U \mid uv \in E\}, \\ d_G(v, U) &:= |N_G(v, U)|, \end{aligned}$$

and abbreviate $N_G(v) := N_G(v, V)$ and $d_G(v) := d_G(v, V)$. Let $\delta(G) := \min\{d_G(v) : v \in V\}$ denote the minimum degree of G .

Let $\mathcal{U} = \{U_1, U_2, \dots, U_k\}$ be a collection of disjoint subsets of V . We write $G[U_1, \dots, U_k]$ or $G[\mathcal{U}]$ for the multipartite subgraph of G with vertex set $U := \cup_{i \in [k]} U_i$ where we keep the *cross-edges* of G (i.e. edges that connect two parts); equivalently, we remove all edges from $G[U]$ that lie inside a

part $U_i \in \mathcal{U}$. In these shorthands, we may omit G whenever it is clear from the context, e.g. writing $[U_1, \dots, U_k]$ for $G[U_1, \dots, U_k]$. We say that \mathcal{U} is a *max-cut k -partition* of G if $e(G[U_1, \dots, U_k])$ is maximised over all k -partitions of $V(G)$.

For disjoint sets V_1, \dots, V_t with $t \geq 2$, let $K[V_1, \dots, V_t]$ denote the complete t -partite graph with parts V_1, \dots, V_t . Its isomorphism class is denoted by $K_{|V_1|, \dots, |V_t|}$. For example, if part sizes differ by at most 1, then we get the Turán graph $T(|V_1| + \dots + |V_t|, t)$. Let M_h denote the matching with h edges.

Definition 2.1. For an edge-colouring $\phi : \binom{[n]}{2} \rightarrow [k]$ of $G := K_n$, define $\text{NIM}(\phi; H_1, \dots, H_k)$ to be the set of all edges not contained in any monochromatic copy of H_i in colour i , and let $\text{nim}(\phi; H_1, \dots, H_k) := |\text{NIM}(\phi; H_1, \dots, H_k)|$. Thus

$$\text{nim}(n; H_1, \dots, H_k) = \max_{\phi: E(K_n) \rightarrow [k]} \text{nim}(\phi; H_1, \dots, H_k).$$

If the H_i 's are all the same graph H , then we will use the shorthands $\text{NIM}_k(\phi; H)$, $\text{nim}_k(\phi; H)$ and $\text{nim}_k(n; H)$ respectively. Also, we may drop k when $k = 2$ and omit the graphs H_i when these are understood. Let G^{nim} be the spanning subgraph of G with $E(G^{\text{nim}}) = \text{NIM}(\phi; H_1, \dots, H_k)$. For $i \in [k]$, denote by G_i and G_i^{nim} the spanning subgraphs of G with edge-sets $E(G_i) = \{e \in E(G) : \phi(e) = i\}$ and $E(G_i^{\text{nim}}) = \{e \in E(G^{\text{nim}}) : \phi(e) = i\}$. We call an edge $e \in E(G^{\text{nim}})$ (respectively, $e \in E(G_i^{\text{nim}})$) a NIM-edge (resp. a NIM- i -edge).

Definition 2.2. For $\xi : \binom{[t]}{\leq 2} \rightarrow [k]$ and disjoint sets V_1, \dots, V_t , the blow-up colouring $\xi(V_1, \dots, V_t) : \binom{V_1 \cup \dots \cup V_t}{2} \rightarrow [k]$ is defined by

$$\xi(V_1, \dots, V_t)(xy) := \begin{cases} \xi(ij), & \text{if } xy \in E(K[V_i, V_j]), \\ \xi(i), & \text{if } x, y \in V_i. \end{cases}$$

If $|V_i| = N$ for every $i \in [t]$, then we say that $\xi(V_1, \dots, V_t)$ is an N -blow-up of ξ .

We say that a colouring ϕ contains another colouring ψ and denote this by $\phi \supseteq \psi$ if ψ is a restriction of ϕ . In particular, $\phi \supseteq \xi(V_1, \dots, V_t)$ means that ϕ is defined on every pair inside $V_1 \cup \dots \cup V_t$ and coincides with $\xi(V_1, \dots, V_t)$ there.

We will make use of the multicolour version of the Szemerédi Regularity Lemma (see, for example, [31, Theorem 1.18]). Let us recall first the relevant definitions. Let $X, Y \subseteq V(G)$ be disjoint non-empty sets of vertices in a graph G . The *density* of (X, Y) is

$$d(X, Y) := \frac{e(G[X, Y])}{|X||Y|}.$$

For $\varepsilon > 0$, the pair (X, Y) is ε -regular if for every pair of subsets $X' \subseteq X$ and $Y' \subseteq Y$ with $|X'| \geq \varepsilon|X|$ and $|Y'| \geq \varepsilon|Y|$, we have $|d(X, Y) - d(X', Y')| \leq \varepsilon$. Additionally, if $d(X, Y) \geq \gamma$, for some $\gamma > 0$, we say that (X, Y) is (ε, γ) -regular. A partition $\mathcal{P} = \{V_1, \dots, V_m\}$ of $V(G)$ is an ε -regular partition of a k -edge-coloured graph G if

1. for all $ij \in \binom{[m]}{2}$, $||V_i| - |V_j|| \leq 1$;
2. for all but at most $\varepsilon \binom{m}{2}$ choices of $ij \in \binom{[m]}{2}$, the pair (V_i, V_j) is ε -regular in each colour.

Lemma 2.3 (Multicolour Regularity Lemma). *For every real $\varepsilon > 0$ and integers $k \geq 1$ and M , there exists M' such that every k -edge-coloured graph G with $n \geq M$ vertices admits an ε -regular partition $V(G) = V_1 \cup \dots \cup V_r$ with $M \leq r \leq M'$.*

Given $\varepsilon, \gamma > 0$, a graph G , a colouring $\phi : E(G) \rightarrow [k]$ and a partition $V(G) = V_1 \cup \dots \cup V_r$, define the *reduced graph*

$$R := R(\varepsilon, \gamma, \phi, (V_i)_{i=1}^r) \tag{6}$$

as follows: $V(R) = \{V_1, \dots, V_r\}$ and V_i and V_j are adjacent in R if (V_i, V_j) is ε -regular with respect to the colour- ℓ subgraph of G for every $\ell \in [k]$ and the colour- m density of (V_i, V_j) is at least γ for some $m \in [k]$. For brevity, we may omit ϕ or $(V_i)_{i=1}^r$ in (6) when these are clear. The graph R comes with the *majority edge-colouring* which assigns to each edge $V_i V_j \in E(R)$ the colour that is the most common one among the edges in $G[V_i, V_j]$ under the colouring ϕ . In particular, the majority colour has density at least γ in $G[V_i, V_j]$. We will use the following consequence of the Embedding Lemma (see e.g. [31, Theorem 2.1]).

Lemma 2.4 (Embedding Lemma). *Let H and R be graphs and let $1 \geq \gamma \gg \varepsilon \gg 1/m > 0$. Let G be a graph obtained by replacing every vertex of R by m vertices, and replacing the edges of R with ε -regular pairs of density at least γ . If R contains a homomorphic copy of H , then $H \subseteq G$.*

We will also need the Slicing Lemma (see e.g. [31, Fact 1.5]).

Lemma 2.5 (Slicing Lemma). *Let $\varepsilon, \alpha, \gamma \in (0, 1)$ satisfy $\varepsilon \leq \min\{\gamma, \alpha, 1/2\}$. If (A, B) is an (ε, γ) -regular pair, then for any $A' \subseteq A$ and $B' \subseteq B$ with $|A'| \geq \alpha|A|$ and $|B'| \geq \alpha|B|$, we have that (A', B') is an $(\varepsilon', \gamma - \varepsilon)$ -regular pair, where $\varepsilon' := \max\{\varepsilon/\alpha, 2\varepsilon\}$.*

Conventions: Throughout the rest of this paper, we will use G as an edge-coloured K_n . For a given number of colours k and a sequence of graphs $(H_i)_{i=1}^k$, we will always write $\psi : \binom{[n]}{2} \rightarrow [k]$ for an extremal colouring realising $\text{nim}(n; H_1, \dots, H_k)$. We do not try to optimise the constants nor prove most general results, instead aiming for the clarify of exposition.

3 Proofs of Theorems 1.11, 1.12 and 1.13

By adding isolated vertices, we can assume that each graph H_i has even order. The following proposition will be frequently used. It basically says that there are no monochromatic copies of $K_{v(H_i), v(H_i)/2}$ in colour i that contains a NIM- i -edge. Its proof follows from the fact that every edge of $K_{v(H_i), v(H_i)/2}$ is in an H_i -subgraph.

Proposition 3.1. *For every graph G , fixed bipartite graphs H_1, \dots, H_k , and a k -edge-colouring $\phi : E(G) \rightarrow [k]$, we have the following for every vertex $v \in V(G)$ and $i \in [k]$. Let $U_i := \{v' \in V(G) : vv' \in G_i^{\text{nim}}\}$.*

(i) *For every vertex $u \in U_i$, the graph $G_i[N_{G_i}(v) \setminus \{u\}, N_{G_i}(u) \setminus \{v\}]$ is $K_{v(H_i), v(H_i)/2}$ -free.*

(ii) *The graph $G_i[U_i, V \setminus (U_i \cup \{v\})]$ is $K_{v(H_i), v(H_i)/2}$ -free. □*

One of the key ingredients for the 2-colour case for bipartite graphs is the following lemma, which is proved by extending an averaging argument of Ma [33]. It states that any 2-edge-colouring of K_n has only linearly many NIM-edges, or there is neither large NIM star nor matching in one of the colours.

Lemma 3.2. *For any h -vertex bipartite H with h even and any 2-edge-colouring ϕ of $G := K_n$ with $\text{nim}(\phi; H) > 2^{2h^2}n$, there exists $i \in [2]$ such that G_i^{nim} is $\{K_{1,h}, M_{h/2}\}$ -free.*

Proof. We may assume, without loss of generality, that G_1^{nim} contains $K_{1,h}$, since otherwise

$$\text{nim}(\phi; H) \leq 2 \cdot \text{ex}(n, K_{1,h}) \leq (h-1)n,$$

contradicting $\text{nim}(\phi; H) > 2^{2h^2}n$. Let S_v be an h -star in G_1^{nim} centred at v . We will show that if G_2^{nim} contains the star $K_{1,h}$ (Case 1) or the matching $M_{h/2}$ (Case 2), then it follows that $\text{nim}(\phi; H) \leq 2^{2h^2}n$, which is a contradiction. In each case, we will define a set $S \subseteq V(G)$, with $h+1 \leq |S| \leq h^2$, containing S_v as follows. In Case 1, let S_u be an h -star centred at u in G_2^{nim} (u and v are not necessarily distinct). Define $S = V(S_v) \cup V(S_u)$ with $h+1 \leq |S| \leq 2h+2$. In Case 2, let $M \subseteq G_2^{\text{nim}}$ be a matching with edge set $\{e_1, \dots, e_{h/2}\}$, where $e_i = z_{i,1}z_{i,2}$ for every $1 \leq i \leq h/2$. Denote $Z := \cup_{i=1}^{h/2} \{z_{i,1}, z_{i,2}\}$. For each edge $e_i \in E(M)$, without loss of generality, assume that $d_{G_2}(z_{i,1}) \geq d_{G_2}(z_{i,2})$. Define iteratively for every $i = 1, \dots, h/2$ a set U'_i as follows,

$$\begin{cases} U'_i \subseteq W_i, & |U'_i| = \frac{h}{2}, & \text{if } |W_i| \geq h/2, \\ U'_i = W_i, & & \text{otherwise,} \end{cases}$$

where $W_i := N_{G_2}(z_{i,1}) \setminus \left(Z \cup \left(\cup_{j=1}^{i-1} U'_j \right) \right)$; further define $U_i := U'_i \cup \{z_{i,1}, z_{i,2}\}$. Finally, set $S := \left(\cup_{i=1}^{h/2} U_i \right) \cup V(S_v)$. So $h+1 \leq |S| \leq h+1 + (h/2+2) \cdot h/2 \leq h^2$.

We now define a partition of $V(G) \setminus S$ that will be used in both Case 1 and Case 2. For each vertex $w \in V(G) \setminus S$, denote by f_w the function $S \rightarrow [2]$ whose value on $s \in S$ is $f_w(s) = \phi(sw)$. In other words, f_w encodes the colours of the edges from w to S . Define

$$\begin{aligned} Y_1 &:= \{v \in V(G) \setminus S : |f_v^{-1}(2)| < h/2\}, \\ Y_2 &:= \{v \in V(G) \setminus S : |f_v^{-1}(1)| < h/2\}, \\ X &:= V(G) \setminus (S \cup Y_1 \cup Y_2). \end{aligned}$$

Thus X consists of those $v \in V(G) \setminus S$ that send at least $h/2$ edges of each colour to S .

We will show in the following claims that, for each class in this partition, there are few vertices in that class or the number of NIM-edges incident to it is linear.

Claim 3.3. $e(G_i^{\text{nim}}[X]) \leq h \binom{|S|}{h/2} n$.

Proof of Claim. Assume to the contrary that $e(G_i^{\text{nim}}[X]) \geq h \binom{|S|}{h/2} n/2$, for some $i \in [2]$. Then there exists a vertex $x \in X$ with $d_{G_i^{\text{nim}}[X]}(x) \geq h \binom{|S|}{h/2}$. By the definition of X , each vertex in $N_{G_i^{\text{nim}}[X]}(x)$ has at least $h/2$ G_i -neighbours in S . By the Pigeonhole Principle, there exists a copy of $K_{h,h/2} \subseteq G_i[N_{G_i^{\text{nim}}[X]}(x), S]$, which is a contradiction by Proposition 3.1(ii). \square

Claim 3.4. $|Y_1| < h \cdot 2^{|S|}$.

Proof of Claim. Assume to the contrary that $|Y_1| \geq h \cdot 2^{|S|}$. Since the total number of functions $S \rightarrow [2]$ is $2^{|S|}$, by averaging, there exists a function f and a subset $Y_f \subseteq Y_1$ with $|Y_f| \geq h$ such that for all vertices $y \in Y_f$, the functions f and f_y are the same. By the definition of Y_1 , there is a

subset $I \subseteq V(S_v) \setminus \{v\}$ with $|I| \geq h/2$ such that for all $s \in I$, $f(s) = 1$, i.e., all pairs between Y_f and I are of colour 1. Recall that S_v is the h -star consisting of NIM- i -edges, thus, there exists a copy of $K_{h/2,h} \subseteq G_1[N_{G_1^{\text{nim}}}(v), Y_f]$, which contradicts Proposition 3.1(ii). \square

We now show that Y_2 has also to be small (given that G_2^{nim} contains a large star or matching), otherwise $\text{nim}(\phi, H)$ is linear.

Case 1: G_2^{nim} has the star $K_{1,h}$.

A similar argument as in Claim 3.4 (with S_u playing the role of S_v) shows that $|Y_2| < h \cdot 2^{|S|}$.

Case 2: G_2^{nim} has the matching $M_{h/2}$.

By the definition of S , all the U_i 's are pairwise disjoint and $h+1 \leq |S| \leq h^2$, see Figure 2. Suppose




Figure 2: Case 2 of Lemma 3.2

that $|Y_2| \geq h \cdot 2^{|S|}$. Again there exists a function $f : S \rightarrow [2]$ with $|f^{-1}(1)| < h/2$ and a subset $Y_f \subseteq Y_2$ with $|Y_f| \geq h$, such that for all vertices $y \in Y_f$, f_y is the same as f . We will use the following claim.

Claim 3.5. For every $1 \leq i \leq h/2$, there exists $w \in U_i$ such that $f(w) = 1$.

Proof of Claim. For a fixed $1 \leq i \leq h/2$, assume to the contrary that for all $s \in U_i$, we have $f(s) = 2$, i.e., $E(G[U_i, Y_f]) \subseteq E(G_2)$. Thus, $|N_{G_2}(z_{i,1}) \setminus (\cup_{j=1}^{i-1} U'_j \cup Z)| \geq |Y_f|$. Consequently, $|U'_i| = h/2$. Therefore, there exists $K_{h/2,h} \subseteq G_2[U_i \setminus \{z_{i,1}\}, Y_f]$, which contradicts Proposition 3.1(i) with $z_{i,1}$ and $z_{i,2}$ playing the roles of v and u respectively. \square

By Claim 3.5 together with the fact that the U_i 's are pairwise disjoint, f assumes value 1 at least $h/2$ times, which contradicts $Y_f \subseteq Y_2$. Therefore, in both cases, $|Y_2| < h \cdot 2^{|S|}$.

Let $Y := Y_1 \cup Y_2$. Since $|S| \leq h^2$, by Claims 3.3 and 3.4, we get that

$$\begin{aligned} \text{nim}(\phi; H) &\leq e(G^{\text{nim}}[S]) + e(G^{\text{nim}}[S, V \setminus S]) + e(G^{\text{nim}}[Y]) + e(G^{\text{nim}}[Y, X]) + e(G^{\text{nim}}[X]) \\ &\leq |S| \cdot n + |Y| \cdot n + e(G^{\text{nim}}[X]) \\ &\leq (h^2 + 2 \cdot h2^{h^2})n + h \binom{h^2}{h/2} n < 2^{2h^2} n, \end{aligned}$$

a contradiction.

This completes the proof of Lemma 3.2. \square

3.1 Weakly-reducible bipartite graphs

Proof of Theorem 1.11. Let H be a weakly-reducible bipartite graph. Let $h = v(H)$ and $w \in V(H)$ be a vertex such that $\text{ex}(n, H - w) < \text{ex}(n, H) - 2^{2h^2}n$ for $n \geq n_0$. In particular, we have that $\text{ex}(n, H) > 2^{2h^2}n$ for $n \geq n_0$. Thus by Lemma 3.2, we may assume that there is $i \in [2]$ such that $e(G_i^{\text{nim}}) \leq \text{ex}(n, \{K_{1,h}, M_{h/2}\}) \leq h^2$. By the symmetry between the two colours, let us assume that $i = 1$. Suppose that $E(G_1^{\text{nim}}) \neq \emptyset$ as otherwise we are trivially done. We now distinguish the following two cases.

Case 1: For every edge $e = uv \in E(G_1^{\text{nim}})$, $d_{G_1}(u) \leq 10h$ and $d_{G_1}(v) \leq 10h$.

In this case, pick one such edge, $e = uv$, and define $V_1 = (N_{G_1}(u) \cup N_{G_1}(v)) \setminus \{u, v\}$. So $|V_1| \leq d_{G_1}(u) + d_{G_1}(v) \leq 20h$. Let

$$V_2 := V(G) \setminus (V_1 \cup \{u, v\}) = N_{G_2}(u) \cap N_{G_2}(v).$$

Note that the subgraph of G_2^{nim} induced on vertex set V_2 satisfies $e(G_2^{\text{nim}}[V_2]) \leq \text{ex}(n, H - w)$. Otherwise, a copy of $H - w$ in $G_2^{\text{nim}}[V_2]$ together with u forms a copy of H in colour 2. Recall that $|V_1| \leq 20h$, $e(G_1^{\text{nim}}) \leq h^2$ and $V_1 \cup V_2 \cup \{u, v\}$ is a partition of $V(G)$. Therefore for large n , we have

$$\begin{aligned} \text{nim}(\psi; H) &\leq e(G_2^{\text{nim}}) + e(G_1^{\text{nim}}) \leq e(G_2^{\text{nim}}[V_2]) + (|V_1| + 2)n + h^2 \\ &\leq \text{ex}(n, H - w) + 30hn \leq \text{ex}(n, H) - 2^{2h^2}n + 30hn < \text{ex}(n, H). \end{aligned} \quad (7)$$

Case 2: There exists an edge $e = uv \in E(G_1^{\text{nim}})$ such that $d_{G_1}(u) \geq 10h$.

Pick $A \subseteq N_{G_1}(u)$ with $|A| = 10h$, and denote

$$\begin{aligned} X &:= \{z \in V(G) \setminus (A \cup \{u, v\}) : d_{G_2}(z, A) \geq h\}, \\ Y &:= \{z \in V(G) \setminus (A \cup \{u, v\}) : d_{G_1}(z, A) \geq h\} \setminus X. \end{aligned}$$

Note that $X \cup Y \cup A \cup \{u, v\}$ is a partition of $V(G)$. We will use the following claims.

Claim 3.6. For every vertex $w \in X \cup Y$, $d_{G_2^{\text{nim}}}(w, X) < h \binom{10h}{h}$.

Proof of Claim. Assume to the contrary that there exists a vertex $w \in X \cup Y$ with $d_{G_2^{\text{nim}}}(w, X) \geq h \binom{10h}{h}$, and define $S := N_{G_2^{\text{nim}}}(w, X)$. Since $|A| = 10h$ and vertices in S all have G_2 -degree at least h in A , there exists a subset of S of size at least h such that its vertices are connected in G_2 to the same h vertices in A , i.e., $K_{h,h} \subseteq G_2[S, A]$, which contradicts Proposition 3.1(ii). \square

Define $Y' = Y \cap N_{G_1}(v)$ to be the set of all vertices in Y that are adjacent to v with a 1-coloured edge, and $Y'' = Y \setminus Y'$.

Claim 3.7. $|Y'| < \binom{10h}{h}h$.

Proof of Claim. Assume to the contrary that $|Y'| \geq \binom{10h}{h}h$. Since all vertices in Y' have at least h G_1 -neighbours in A , there exists a copy of $K_{h,h} \subseteq G_1[Y', A]$, which extends to a copy of $K_{h+1, h+1} \supseteq H$ containing the edge $uv \in E(G_1^{\text{nim}})$, a contradiction. \square

By Claims 3.6, 3.7 and since $|A| = 10h$, the number of edges in G_2^{nim} with at least one end point in the set $A \cup Y' \cup X \cup \{u, v\}$ is at most $3h \binom{10h}{h} n$. It remains to estimate $e(G_2^{\text{nim}}[Y''])$. We claim that $e(G_2^{\text{nim}}[Y'']) \leq \text{ex}(n, H - w)$. Otherwise, since all the edges connecting v to Y'' have colour 2, we can extend the copy of $H - w \subseteq G_2^{\text{nim}}[Y'']$ to a copy of H by adding v . This contradicts the definition of G_2^{nim} . Hence,

$$\begin{aligned} \text{nim}(\psi; H) &= e(G_2^{\text{nim}}) + e(G_1^{\text{nim}}) \leq 3h \binom{10h}{h} n + \text{ex}(n, H - w) + h^2 \\ &< \text{ex}(n, H) - 2^{2h^2} n + 4h \binom{10h}{h} n < \text{ex}(n, H). \end{aligned}$$

Thus, any colouring with NIM-edges of two different colours is not extremal. \square

3.2 General bipartite graphs

In this subsection, we will prove Theorems 1.12 and 1.13.

Proof of Theorem 1.12. As H contains a cycle, $\text{ex}(n, H)/n \rightarrow \infty$ as $n \rightarrow \infty$. Then by Lemma 3.2, we may assume that, for example, G_1^{nim} is $\{K_{1,h}, M_{h/2}\}$ -free. Since G_2^{nim} is H -free, we immediately get that

$$\text{nim}_2(n, H) \leq \text{ex}(n, H) + \text{ex}(n, \{K_{1,h}, M_{h/2}\}) \leq \text{ex}(n, H) + h^2,$$

as desired. \square

Proof of Theorem 1.13. Let us first present the part of the proof which works for an arbitrary number of colours k and any forest T . Let $h = v(T)$.

The stated lower bound on $\text{nim}_k(n; T)$ can be obtained by using the argument of Ma [33]. Fix some maximum T -free graph H on $[n]$ and take uniform independent permutations $\sigma_1, \dots, \sigma_{k-1}$ of $[n]$. Iteratively, for $i = 1, \dots, k-1$, let the colour- i graph G_i consists of those pairs $\{\sigma_i(x), \sigma_i(y)\}$, $xy \in E(H)$, that are still uncoloured. Finally, colour all remaining edges with colour k . Clearly, all edges of colours between 1 and $k-1$ are NIM-edges. Since $e(H) \leq hn = O(n)$, the expected size of $\sum_{i=1}^{k-1} e(G_i)$ is at least

$$(k-1)e(H) - \binom{k-1}{2} e(H)^2 \binom{n}{2}^{-1} \geq (k-1)\text{ex}(n, T) - k^2 h^2.$$

By choosing the permutations for which $\sum_{i=1}^{k-1} e(G_i)$ is at least its expectation, we obtain the required bound.

Let us turn to the upper bound. Fix an extremal G with colouring $\phi : \binom{[n]}{2} \rightarrow [k]$, so $\text{nim}_k(\phi; T) = \text{nim}_k(n; T)$. For every $1 \leq i \leq k$, denote

$$A_i := \{v \in V(G) : \exists u \in V(G), uv \in E(G_i^{\text{nim}})\} \quad \text{and} \quad a_i := |A_i|.$$

In other words, A_i is the set of all vertices incident with at least one i -coloured NIM-edge. Note that

$$\text{nim}_k(\phi; T) \leq \sum_{i=1}^k \text{ex}(a_i, T). \tag{8}$$

Also, for every $X \subseteq [k]$, define

$$B_X := \{v \in V(G) : v \in A_i \Leftrightarrow i \in X\} = \cap_{i \in X} A_i \setminus (\cup_{j \notin X} A_j) \quad \text{and} \quad b_X := |B_X|.$$

In other words, B_X is the set of vertices which are incident with edges in G_i^{nim} if and only if $i \in X$. By definition, for two distinct subsets $X, Y \subseteq [k]$, $B_X \cap B_Y = \emptyset$.

Claim 3.8. For every two subsets $X, Y \subseteq [k]$ with $X \cup Y = [k]$, $\min\{b_X, b_Y\} < 6kh$.

Proof of Claim. Assume on the contrary that there exist two subsets $X, Y \subseteq [k]$ such that $X \cup Y = [k]$ and $b_X, b_Y \geq 6kh$. Let $B'_X \subseteq B_X$ and $B'_Y \subseteq B_Y$ be such that $|B'_X| = |B'_Y| = 6kh$. By averaging, some colour, say colour 1, contains at least $1/k$ proportion of edges in $G[B'_X, B'_Y]$. Set $F = G_1[B'_X, B'_Y]$. Then there exists $F' \subseteq F$ on vertex set $B''_X \cup B''_Y$, where $B''_X \subseteq B'_X$ and $B''_Y \subseteq B'_Y$, such that the minimum degree of F' is at least half of the average degree of F , that is,

$$\delta(F') \geq \frac{e(F)}{|V(F)|} \geq \frac{|B'_X| \cdot |B'_Y|}{k \cdot (|B'_X| + |B'_Y|)} = \frac{(6kh)^2}{k \cdot 12kh} = 3h.$$

Let $v \in V(T)$ be a leaf, u be its only neighbour, and $T' := T - v$, where $T - v$ is the forest obtained from deleting the leaf v from T . Since $X \cup Y = [k]$, without loss of generality, we can assume that $1 \in X$. Fix an arbitrary vertex $x \in B''_X$ and let w be a G_1^{nim} -neighbour of x . (Such a vertex exists as $x \in B''_X \subseteq B_X$ and $1 \in X$.) Then $\delta(F' - w) \geq \delta(F') - 1 \geq 2h$. We can then greedily embed T' in $F' - w$ with x playing the role of u . As this copy of T' is in $F' - w \subseteq G_1$, together with $xw \in G_1^{\text{nim}}$, we get a monochromatic copy of T with an edge in G^{nim} , a contradiction (see Figure 3). \square




Figure 3: Finding a copy of $T \in G_1$.

We will divide the rest of the proof into two cases.

Case 1: There exists a subset $X \subset [k]$ such that $|X| = k - 1$ and $b_X \geq 6kh$.

Let $\{j\} = [k] \setminus X$, and \mathcal{Y} be the collection of all subsets of $[k]$ containing j . By Claim 3.8, $b_Y < 6kh$, for every set $Y \in \mathcal{Y}$, implying that $a_j = \sum_{Y \in \mathcal{Y}} b_Y < 2^k \cdot 6kh$. Hence, by (8),

$$\begin{aligned} \text{nim}_k(\phi; T) &\leq \sum_{i \in [k]} \text{ex}(a_i, T) \leq \sum_{i \in [k] \setminus \{j\}} \text{ex}(a_i, T) + \text{ex}(2^k \cdot 6kh, T) \\ &\leq (k - 1)\text{ex}(n, T) + 2^k \cdot 6kh^2. \end{aligned}$$

Thus the theorem holds in this case.

Case 2: For all subsets $X \subset [k]$ with $|X| = k - 1$, we have $b_X < 6kh$.

By Claim 3.8, we have $b_{[k]} \leq 2 \cdot 6kh$. Hence, all but at most $(k+2)6kh$ vertices are adjacent to NIM-edges with at most $k-2$ different colours, which implies that they are in at most $k-2$ different sets A_i . Therefore,

$$a_1 + \cdots + a_k \leq (k-2)n + 12(k+2)kh. \quad (9)$$

Now our analysis splits further, depending on the cases of Theorem 1.13. If $k=2$, then we are done by (8) and (9):

$$\text{nim}_2(\phi; T) \leq h(a_1 + a_2) \leq 96h^2 \leq \text{ex}(n, T).$$

Thus it remains to consider the case when $k \geq 3$ and T is a tree. By taking the disjoint union of two maximum T -free graphs, we see that the Turán function of T is superadditive, that is,

$$\text{ex}(\ell, T) + \text{ex}(m, T) \leq \text{ex}(\ell + m, T), \quad \text{for any } \ell, m \in \mathbb{N}. \quad (10)$$

The Fekete Lemma implies that $\text{ex}(m, T)/m$ tends to a limit τ . Since, for example, $\text{ex}(m, T) \leq hm$, we have that $\tau \leq h$, in particular, τ is finite. Also, excluding the case $T = K_2$ when the theorem trivially holds, we have $\tau > 0$. In particular, $|\text{ex}(m, T)/m - \tau| < c$ for all large m , where $c := \tau/(3k-4) > 0$ satisfies $(\tau+c)(k-2) = (k-1)(\tau-2c)$.

Thus (8), (9), (10) and the fact that n is sufficiently large give that

$$\begin{aligned} \text{nim}_k(\phi; T) &\leq \text{ex}(a_1 + \cdots + a_k, T) \leq \text{ex}((k-2)n + 12(k+2)kh, T) \\ &\leq (\tau+c)((k-2)n + 12(k+2)kh) \leq (k-1)(\tau-2c)n + 24(k+2)kh^2 \\ &\leq (k-1)(\tau-c)n \leq (k-1)\text{ex}(n, T). \end{aligned}$$

This finishes the proof of Theorem 1.13. □

4 Proofs of Theorems 1.5 and 1.6

We need the following lemma, which states that the reduced graph of the NIM-graph cannot have a large clique, linking the nim function to the new Ramsey variant r^* .

Lemma 4.1. *For $i \in [k]$, let H_i be a non-bipartite graph, and let $1/k, 1/r \geq \gamma \gg \varepsilon \gg 1/N > 0$, where $r := R(a_1 - 1, \dots, a_k - 1)$ and $a_i := \chi(H_i)$. Let V_1, \dots, V_m be disjoint sets, each of size at least N . Take any $\phi : \binom{V}{2} \rightarrow [k]$, where $V := V_1 \cup \cdots \cup V_m$, and let G^{nim} be the NIM-graph of ϕ . Then the graph $R := R(\varepsilon, \gamma, \phi|_{E(G^{\text{nim}})}, (V_i)_{i=1}^m)$ is K_{r^*+1} -free, where $r^* := r^*(H_1, \dots, H_k)$.*

Proof. Given the graphs H_i with $a_i = \chi(H_i)$, and $r = R(a_1, \dots, a_k)$, choose additional constants so that the following hierarchy holds:

$$\frac{1}{r} \gg \gamma \gg \varepsilon_1 \gg \frac{1}{M} \gg \varepsilon \gg \frac{1}{N} > 0.$$

Let the V_i 's and ϕ be as in the statement of the lemma. For each $i \in [m]$, apply the Multicolour Regularity Lemma (Lemma 2.3) with constants ε_1 and $1/\varepsilon_1$ to the k -coloured complete graph on V_i to obtain an ε_1 -regular partition $V_i = U_{i,1} \cup \cdots \cup U_{i,m_i}$ with $1/\varepsilon_1 \leq m_i \leq M$. Let $R_i := R(\varepsilon_1, \gamma, \phi|_{\binom{V_i}{2}}, (U_{i,j})_{j=1}^{m_i})$ be the associated reduced graph.

Note that the fraction of the elements $xy \in \binom{V_i}{2}$ with $x \in U_{i,a}$ and $y \in U_{i,b}$ such that the pair $(U_{i,a}, U_{i,b})$ is not ε_1 -regular in some colour or satisfies $a = b$ is at most $\varepsilon_1 + 1/m_i$. Since $\gamma \leq 1/k$, the remaining elements of $\binom{V_i}{2}$ come from edges of R_i . Recall that $m_i = v(R_i)$. Thus, we have that

$$e(R_i) \geq \frac{(1 - \varepsilon_1 - 1/m_i) \binom{|V_i|}{2}}{\lceil |V_i|/m_i \rceil^2} \geq (1 - 2\varepsilon_1) \binom{m_i}{2}. \quad (11)$$

Let $\xi : E(R) \rightarrow [k]$ be the colouring of R . We extend it to the vertices of R as follows. Take $i \in [m]$. Let $\xi_i : E(R_i) \rightarrow [k]$ be the colouring of R_i . By (11) and since $v(R_i) \geq 1/\varepsilon_1$ and $\varepsilon_1 \ll 1/r$, we have that $e(R_i) > \text{ex}(m_i, K_r)$. By Turán's theorem [48], the graph R_i contains an r -clique. By the definition of r , the restriction of the k -edge-colouring ξ_i to this r -clique contains a colour- p copy of K_{a_p-1} for some $p \in [k]$. Let ξ assign the colour p to V_i .

Suppose to the contrary that some $(r^* + 1)$ -set A spans a clique in R . The restriction of ξ to $\binom{A}{\leq 2}$ violates either (P1) or (P2) from Definition 1.3. We will derive contradictions in both cases, thus finishing the proof. If ξ contains an edge-monochromatic homomorphic copy of some H_i in colour $i \in [k]$, then by the Embedding Lemma (Lemma 2.4) the colour- i subgraph of G^{nim} contains a copy of H_i , a contradiction to G^{nim} consisting of the NIM-edges. So suppose that (P2) fails, say, some pair $V_i V_j \in \binom{A}{2}$ satisfies $\xi(V_i V_j) = \xi(V_i)$, call this colour p . By the definition of $\xi(V_i)$, R_i contains an $(a_p - 1)$ -clique of colour p under ξ_i , say with vertices $U_1, \dots, U_{a_p-1} \in V(R_i)$. Observe that $\varepsilon_1 \geq \max\{2\varepsilon, \varepsilon M\} \geq \max\{2\varepsilon, \varepsilon \cdot v(R_i)\}$ and p is the majority colour on edges in $G^{\text{nim}}[V_i, V_j]$. The Slicing Lemma (Lemma 2.5) with e.g. $\alpha := 1/M$ gives that each pair (V_j, U_h) with $h \in [a_p - 1]$ is $(\varepsilon_1, \gamma/2)$ -regular in G_p^{nim} . The Embedding Lemma (Lemma 2.4) gives a copy of H_p in G containing at least one (in fact, at least $\delta(H_p)$) edges of $G^{\text{nim}}[V_i, V_j]$, a contradiction. \square

Proof of Theorem 1.5. For the upper bound, let $1 \gg \varepsilon \gg \gamma \gg \varepsilon_1 > 0$. Let n be large and suppose to the contrary that there exists some colouring $\phi : E(K_n) \rightarrow [k]$ that violates (4). Apply the Multicolour Regularity Lemma (Lemma 2.3) to the NIM-graph G^{nim} of ϕ with parameters ε_1 and $1/\varepsilon_1$. A calculation similar to the one in (11) applies here, where additionally one has to discard at most $k\gamma \binom{n}{2}$ edges in $\text{NIM}(\phi)$ coming from pairs that have density less than γ in each colour. By $\gamma \ll \varepsilon$, we conclude that the reduced graph $R = R(\varepsilon_1, \gamma, \phi|_{E(G^{\text{nim}})})$ of G^{nim} has at least $(1 - 1/r^* + \varepsilon/2) \frac{v(R)^2}{2}$ edges. By Turán's theorem, $K_{r^*+1} \subseteq R$, contradicting Lemma 4.1.

For the lower bound, take a feasible k -colouring ξ of $\binom{[r^*]}{\leq 2}$, where $r^* := r^*(H_1, \dots, H_k)$. If possible, among all such colourings take one such that all singletons have the same colour. Consider the blow-up colouring $\phi := \xi(X_1, \dots, X_{r^*})$ where the sets X_i form an equipartition of $[n]$.

Let us show that every edge of $K[X_1, \dots, X_{r^*}]$ is a NIM-edge. Take any copy F of H_i which is i -monochromatic in ϕ . Since the restriction of ξ to $\binom{[r^*]}{2}$ has no homomorphic copy of H_i by (P1), the graph F must use at least one edge that is inside some V_j . If ξ assigns the value i only to singletons, then no edge of the colour- i graph F can be a cross-edge. Otherwise, if F is connected, then $E(F) \subseteq \binom{V_j}{2}$ because no edge between V_j and its complement can have ϕ -colour i by (P2). We conclude that every cross-edge is a NIM-edge, giving the required lower bound. \square

Proof of Theorem 1.6. Choose $\varepsilon \gg \varepsilon_1 \gg \delta \gg \gamma \gg \varepsilon_2 \gg 1/n > 0$. Let ϕ be as in the theorem. Apply the Regularity Lemma (Lemma 2.3) to NIM-graph G^{nim} with parameters ε_2 and $1/\varepsilon_2$ to get an ε_2 -regular partition $V(G^{\text{nim}}) = V_1 \cup \dots \cup V_m$. Let $R = R(\varepsilon_2, \gamma, \phi|_{E(G^{\text{nim}})}, (V_i)_{i=1}^m)$ be the reduced graph. A similar calculation as in (11) yields that $e(R) \geq (1 - \frac{1}{r^*} - 2\delta) \frac{m^2}{2}$. On the other hand, by Lemma 4.1, R is K_{r^*+1} -free. Thus, the Erdős-Simonovits Stability Theorem [11, 42]

implies that $\delta_{\text{edit}}(R, T(m, r^*)) \leq \varepsilon_1 m^2/2$. Let a partition $V(R) = \mathcal{U}_1 \cup \dots \cup \mathcal{U}_{r^*}$ minimise $|E(R) \Delta E(K[\mathcal{U}_1, \dots, \mathcal{U}_{r^*}])|$. We know that the minimum is at most $\varepsilon_1 m^2/2$. Let $V(G^{\text{nim}}) = W_1 \cup \dots \cup W_{r^*}$ be the partition induced by \mathcal{U}_i 's, i.e., $W_i := \cup_{V_j \in \mathcal{U}_i} V_j$ for $i \in [r^*]$. Let G' be the graph obtained from G^{nim} by removing all edges that lie in any cluster V_i ; or between those parts V_i and V_j such that $V_i V_j$ is not an ε_2 -regular pair or belongs to $E(R) \Delta E(K[\mathcal{U}_1, \dots, \mathcal{U}_{r^*}])$. We have

$$|E(G^{\text{nim}}) \Delta E(G')| = |E(G^{\text{nim}}) \setminus E(G')| \leq m \cdot \frac{(n/m)^2}{2} + \varepsilon_2 m^2 \cdot \frac{n^2}{m^2} + |E(R) \Delta E(T(m, r^*))| \cdot \frac{n^2}{m^2} \leq \varepsilon_1 n^2.$$

As $e(G^{\text{nim}}) \geq (1 - 1/r^*)n^2/2 - \delta n^2$, we have $e(G') \geq (1 - 1/r^*)n^2/2 - 2\varepsilon_1 n^2$. Since G' is r^* -partite (with parts W_1, \dots, W_{r^*}), a direct calculation gives that $\delta_{\text{edit}}(G', T(n, r^*)) \leq \varepsilon n^2/2$. Finally, we obtain

$$\delta_{\text{edit}}(G^{\text{nim}}, T(n, r^*)) \leq |E(G^{\text{nim}}) \Delta E(G')| + \delta_{\text{edit}}(G', T(n, r^*)) \leq \varepsilon_1 n^2 + \frac{\varepsilon n^2}{2} \leq \varepsilon n^2,$$

as desired. \square

5 Proof of Theorem 1.7

The following lemma will be useful in the forthcoming proof of Theorem 1.7. It is proved by an easy modification of the standard proof of Ramsey's theorem.

Lemma 5.1 (Partite Ramsey Lemma). *For every triple of integers $k, r, u \in \mathbb{N}$ there is $\rho = \rho(k, r, u)$ such that if ϕ is a k -edge-colouring of the complete graph on $Y_1 \cup \dots \cup Y_r$, where Y_1, \dots, Y_r are disjoint ρ -sets, then there are u -sets $U_i \subseteq Y_i$, $i \in [r]$, and $\xi : \binom{[r]}{\leq 2} \rightarrow [k]$ such that $\xi(U_1, \dots, U_r) \subseteq \phi$. (In other words, we require that each $\binom{U_i}{2}$ and each bipartite graph $[U_i, U_j]$ is monochromatic.)*

Proof. We use induction on r with the case $r = 1$ being the classical Ramsey theorem. Let $r \geq 2$ and set $N := (u - 1)k^r + 1$. We claim that $\rho := (2k)^N \rho(k, r - 1, u)$ suffices here. Let ξ be an arbitrary k -edge-colouring of the complete graph on $Y_1 \cup \dots \cup Y_r$ where each $|Y_i| = \rho$.

Informally speaking, we iteratively pick vertices x_1, \dots, x_N in Y_r shrinking the parts so that each new vertex x_i is monochromatic to each part. Namely, we initially let $U_i^0 := Y_i$ for $i \in [r]$. Then for $i = 1, \dots, N$ we repeat the following step. Given vertices x_1, \dots, x_{i-1} and sets $U_r^{i-1} \subseteq Y_r \setminus \{x_1, \dots, x_{i-1}\}$ and $U_j^{i-1} \subseteq Y_j$ for $j \in [r - 1]$, we let x_i be an arbitrary vertex of U_r^{i-1} and, for $j \in [r]$, let U_j^i be a maximum subset of U_j^{i-1} such that all pairs between x_i and U_j^i have the same colour, which we denote by $c_j^i \in [k]$. Clearly, $|U_j^i| \geq (|U_j^{i-1}| - 1)/k$ (the -1 term is need for $j = r$), which is at least $(2k)^{N-i}$ by a simple induction on i . Thus we can carry out all N steps. Moreover, each of the the final sets U_1^N, \dots, U_{r-1}^N has size at least $\rho/(2k)^N = \rho(k, r - 1, u)$. By the induction assumption, we can find u -sets $U_j \subseteq U_j^N$, $j \in [r - 1]$, and $\xi : \binom{[r-1]}{\leq 2} \rightarrow [k]$ with $\xi(U_1, \dots, U_{r-1}) \subseteq \phi$.

Each selected vertex x_i comes with a colour sequence $(c_1^i, \dots, c_r^i) \in [k]^r$. So we can find a set $U_r \subseteq \{x_1, \dots, x_N\}$ of $\lceil N/k^r \rceil = u$ vertices that have the same colour sequence (c_1, \dots, c_r) . Clearly, all pairs in $\binom{U_r}{2}$ (resp. $[U_r, U_j]$ for $j \in [r - 1]$) have the same colour c_r (resp. c_j). Thus if we extend the colouring ξ to $\binom{[r]}{\leq 2}$ by letting $\xi(i, r) := c_i$ for $i \in [r - 1]$ and $\xi(r) := c_r$, then $\xi(U_1, \dots, U_r) \subseteq \phi$, as required. \square

The main step in proving Theorem 1.7 is given by the following lemma.

Lemma 5.2. *Under the assumptions of Theorem 1.7, there is n_0 such that if ϕ is an arbitrary k -edge-colouring of $G := K_n$ with $n \geq n_0$, $e(G^{\text{nim}}) \geq t(n, r^*)$ and*

$$\delta(G^{\text{nim}}) \geq \delta(T(n, r^*)), \quad (12)$$

where δ denotes the minimum degree, then $G^{\text{nim}} \cong T(n, r^*)$ (in particular, $e(G^{\text{nim}}) = t(n, r^*)$).

Proof. Let H_i , $i \in [k]$, and r^* be as in Theorem 1.7. So $r^* = r^*(H_1, \dots, H_k)$. Let

$$N := \max_{i \in [k]} v(H_i) - 1 \geq 1 \quad \text{and} \quad 1 \gg \varepsilon \gg \varepsilon_1 \gg 1/n_0 > 0.$$

Let $n \geq n_0$ and let ϕ be an arbitrary k -edge-colouring of $G := K_n$.

Let $\mathcal{P} = \{V_1, \dots, V_{r^*}\}$ be a max-cut r^* -partition of G^{nim} . In particular, for every $i, j \in [r^*]$ and every $v \in V_i$, we have $d_{G^{\text{nim}}}(v, V_j) \geq d_{G^{\text{nim}}}(v, V_i)$. By applying Theorem 1.6 to G^{nim} , we have

$$e(G^{\text{nim}}[\mathcal{P}]) \geq t(n, r^*) - \varepsilon_1 n^2. \quad (13)$$

A simple calculation shows that $|V_i| = \frac{n}{r^*} \pm \sqrt[3]{\varepsilon_1} n$ for all $i \in [r^*]$.

Claim 5.3. For every $i \in [r^*]$ and $v \in V_i$, $d_{G^{\text{nim}}}(v, V_i) \leq \varepsilon n$.

Proof of Claim. Assume to the contrary that there exist $i \in [r^*]$ and $v \in V_i$ such that $d_{G^{\text{nim}}}(v, V_i) > \varepsilon n$. For each $j \in [r^*]$, as \mathcal{P} is a max-cut, there exists a colour $\ell \in [k]$ such that

$$d_{G_\ell^{\text{nim}}}(v, V_j) \geq d_{G^{\text{nim}}}(v, V_j)/k \geq d_{G^{\text{nim}}}(v, V_i)/k \geq \varepsilon n/k =: m.$$

So, for $j \in [r^*]$, let $Z_j \subseteq N_{G_\ell^{\text{nim}}}(v, V_j)$ be any subset of size m . We have

$$e(\overline{G^{\text{nim}}}[Z_1, \dots, Z_{r^*}]) \leq e(\overline{G^{\text{nim}}}[V_1, \dots, V_{r^*}]) \stackrel{(13)}{\leq} \varepsilon_1 n^2. \quad (14)$$

Let $\rho := \rho(k, r^*, N)$, where ρ is the function from the Partite Ramsey Lemma (Lemma 5.1). For $i \in [r^*]$, let Y_i be a random ρ -subset of Z_i , chosen uniformly and independently at random. By (14), the expected number of missing cross-edges in $G^{\text{nim}}[Y_1, \dots, Y_{r^*}]$ is at most

$$\varepsilon_1 n^2 \left(\binom{m-1}{\rho-1} / \binom{m}{\rho} \right)^2 = \varepsilon_1 \left(\frac{\rho k}{\varepsilon} \right)^2 < 1.$$

Thus there is a choice of the ρ -sets Y_i 's such that $G^{\text{nim}}[Y_1, \dots, Y_{r^*}]$ has no missing cross-edges. By the definition of ρ , there are N -sets $U_1 \subseteq Y_1, \dots, U_{r^*} \subseteq Y_{r^*}$ and a colouring $\xi : \binom{[r^*]}{\leq 2} \rightarrow [k]$ such that $\xi(U_1, \dots, U_{r^*}) \subseteq \phi$.

Note that ξ is feasible. Indeed, if we have, for example, $\xi(ij) = \xi(i) =: c$, then by taking one vertex of U_j and all N vertices of U_i we get a colour- c copy of K_{N+1} . However, since $N+1 \geq v(H_c)$, every edge of this clique is in an H_c -subgraph, contradicting the fact that all pairs in the complete bipartite graph $K[U_i, U_j]$ are NIM-edges.

Consequently, as (H_1, \dots, H_k) is nice, ξ must assign the same colour to all singletons, say colour 1. By construction, the vertex v is monochromatic into each $Z_i \supseteq U_i$. So we can take $\xi' : \binom{[r^*+1]}{\leq 2} \rightarrow [k]$ such that $\xi'(U_1, \dots, U_{r^*}, \{v\}) \subseteq \phi$, where we additionally let $\xi'(r^*+1) := 1$. As $r^* = r^*(H_1, \dots, H_k)$, the colouring ξ' violates (P1) or (P2). This violation has to include

the vertex $r^* + 1$ since the restriction of ξ' to $\binom{[r^*]}{\leq 2}$ is the feasible colouring ξ . We cannot have $i \in [r^*]$ with $\xi'(i, r^* + 1) = 1$ because otherwise $U_i \cup \{v\}$ is an $(N + 1)$ -clique coloured 1 under ϕ , a contradiction to all pairs between $Z_i \supseteq U_i$ and v being NIM-edges. Therefore, there exists an edge-monochromatic homomorphic copy of H_j of colour j , say F , with $r^* + 1 \in V(F)$. By the definition of homomorphism-criticality, there exists a homomorphism $g : V(H_j) \rightarrow V(F)$ such that $|g^{-1}(r^* + 1)| = 1$. Therefore, we can find an edge-monochromatic copy of H_j in colour j , with $g^{-1}(r^* + 1)$ mapped to v , and all the other vertices of H_j mapped to vertices in $U_1 \cup \dots \cup U_{r^*}$, a contradiction to all pairs between this set and v being NIM-edges. \square

We next show that all pairs inside a part get the same colour under ϕ .

Claim 5.4. For any $p \in [r^*]$ and any $u_1 u_2, u_3 u_4 \in \binom{V_p}{2}$, we have $\phi(u_1 u_2) = \phi(u_3 u_4)$.

Proof of Claim. Suppose on the contrary that $u_1, \dots, u_4 \in V_p$ violate the claim. Without loss of generality, let $p = r^*$. Let $U := \{u_1, \dots, u_4\}$. By (12), Claim 5.3 and the fact that $|V_r| = n/r \pm \sqrt[3]{\varepsilon_1} n$, all but at most $2\varepsilon n$ edges from any $u \in V_{r^*}$ to $V \setminus V_{r^*}$ are NIM-edges. For $i \in [r^* - 1]$ (resp. $i = r^*$), define $Z_i \subseteq V_i$ to be a largest subset of $\bigcap_{j=1}^4 N_{G^{\text{nim}}}(u_j, V_i)$ (resp. $V_{r^*} \setminus U$) with the same colour pattern to U , i.e., for all $x, x' \in Z_i$ and $j \in [4]$ we have $\phi(u_j x) = \phi(u_j x')$. By the Pigeonhole Principle, we have for $i \in [r^* - 1]$ that

$$|Z_i| \geq \frac{|\bigcap_{j=1}^4 N_{G^{\text{nim}}}(u_j, V_i)|}{k^4} \geq \frac{|V_i| - 4 \cdot 2\varepsilon n}{k^4} \geq \frac{n}{2r^* k^4}.$$

Also, $|Z_{r^*}| \geq (|V_{r^*}| - 4)/k^4 \geq n/(2r^* k^4)$.

Similarly to the calculation after (14), there are N -subsets $U_i \subseteq Z_i$, $i \in [r^*]$, such that ϕ contains the blow-up $\xi(U_1, \dots, U_{r^*})$ of some $\xi : \binom{[r^*]}{\leq 2} \rightarrow [k]$. As in the proof of Claim 5.3, ξ is feasible and assigns the same colour, say 1, to all singletons. Since $\phi(u_1 u_2) \neq \phi(u_3 u_4)$, assume that e.g. $\phi(u_1 u_2) \neq 1$.

We define the colouring $\xi' : \binom{[r^*+1]}{\leq 2} \rightarrow [k]$ so that $\xi'(U_1, \dots, U_{r^*-1}, \{u_1\}, \{u_2\}) \subseteq \phi$, where additionally we let both $\xi'(r^*)$ and $\xi'(r^* + 1)$ be 1. Note that $\xi'(r^*, r^* + 1) = \phi(u_1 u_2)$. Also, observe that we do not directly use the part U_{r^*} when defining ξ' : the role of this part was to guarantee that ξ is monochromatic on all singletons. By the definition of r^* , the colouring ξ' violates (P1) or (P2).

Suppose first that ξ' violates (P2), that is there is a pair $ij \in \binom{[r^*+1]}{2}$ with $\xi'(ij) = 1$. Since $\xi'(r^*, r^* + 1) = \phi(u_1 u_2) \neq 1$, we have $\{i, j\} \neq \{r^*, r^* + 1\}$. Also, we cannot have $i, j \in [r^* - 1]$, because ξ' coincides on $\binom{[r^*-1]}{\leq 2}$ with the feasible colouring ξ . So we can assume by symmetry that $i \in [r^* - 1]$ and $j = r^*$. However, then the vertex u_1 is connected by NIM-1-edges to the colour-1 clique on the N -set U_i , a contradiction.

We may now assume that the colouring ξ' violates (P1). Let this be witnessed by an edge-monochromatic homomorphic copy of H_j of colour j , say F . If F contains exactly one vertex from $\{r^*, r^* + 1\}$, then by an argument similar to the last part of the proof of Claim 5.3 we get a contradiction. Otherwise, if $\{r^*, r^* + 1\} \subseteq V(F)$, then, by the definition of homomorphism-critical, there exists a homomorphism $g : V(H_j) \rightarrow V(F)$ such that $|g^{-1}(r^*)| = |g^{-1}(r^* + 1)| = 1$. Therefore, we can find an edge-monochromatic copy of H_j in colour j , with $g^{-1}(r^*)$ (resp. $g^{-1}(r^* + 1)$) mapped to u_1 (resp. u_2), and all the other vertices of H_j mapped to vertices in $U_1 \cup \dots \cup U_{r^*-1}$, a contradiction to all pairs between this set and $\{u_1, u_2\}$ being NIM-edges. \square

Let $i \in [r^*]$. By Claim 5.4 we know that $G[V_i]$ is a monochromatic clique. Since $|V_i| \geq \max_{j \in [k]} v(H_j)$, no pair inside V_i is a NIM-edge. Thus G^{nim} is r^* -partite. Our assumption $e(G^{\text{nim}}) \geq t(n, r^*)$ implies that G^{nim} is isomorphic to $T(n, r^*)$, as desired. \square

We are now ready to prove the desired exact result.

Proof of Theorem 1.7. We know by Theorem 1.5 that $\text{nim}(n; H_1, \dots, H_k) \geq t(n, r^*)$ for all n .

On the other hand, let n_0 be the constant returned by Lemma 5.2. Let $n \geq n_0^2$ and let ψ be an extremal colouring of $G := K_n$. In order to finish the proof of the theorem it is enough to show that necessarily $G^{\text{nim}} \cong T(n, r^*)$.

Initially, let $i = n$, $G_n := G$ and $\phi_n := \psi$. Iteratively repeat the following step as long as possible: if the NIM-graph of ϕ_i has a vertex x_i of degree smaller than $\delta(T(i, r^*))$, let ϕ_{i-1} be the restriction of ϕ_i to the edge-set of $G_{i-1} := G_i - x_i$ and decrease i by 1. Suppose that this procedure ends with G_m and ϕ_m .

Note that, for every $i \in \{m+1, \dots, n\}$, we have that

$$\begin{aligned} t(i-1, r^*) &= t(i, r^*) - \delta(T(i, r^*)), \\ \text{nim}(\phi_{i-1}) &\geq \text{nim}(\phi_i) - \delta(T(i, r^*)) + 1, \end{aligned}$$

the latter inequality following from the fact that every NIM-edge of ϕ_i not incident to x_i is necessarily a NIM-edge of ϕ_{i-1} . These two relations imply by induction that

$$\text{nim}(\phi_i) \geq t(i, r^*) + n - i, \quad \text{for } i = n, n-1, \dots, m. \quad (15)$$

In particular, it follows that $m > n_0$ for otherwise $\text{NIM}(\phi_{n_0})$ is a graph of order n_0 with at least $n - n_0 > \binom{n_0}{2}$ edges, which is impossible. Thus Lemma 5.2 applies to ϕ_m and gives that $\text{NIM}(\phi_m) \cong T(m, r^*)$. By (15) we conclude that $m = n$, finishing the proof of Theorem 1.7. \square

6 Proofs of Theorems 1.8 and 1.9

Next we will show that Conjecture 1.4 holds for the 3-colour case.

Proof of Theorem 1.8. Take an arbitrary feasible 3-colouring ξ of $\binom{[r]}{\leq 2}$, where $r = R(a_2, a_3) - 1$. It suffices to show that ξ assigns the same colour to all the singletons in $[r]$. Indeed, suppose that $(K_{a_1}, K_{a_2}, K_{a_3})$ is not nice. Then there exists a feasible 3-colouring ξ^* of $\binom{[r^*]}{\leq 2}$ that is not monochromatic on the singletons in $[r^*]$, where $r^* := r^*(K_{a_1}, K_{a_2}, K_{a_3}) \geq r$. Up to relabeling, we may assume that $[r]$ contains two singletons of different colours in ξ^* . We then arrive to a contradiction, as the restriction of ξ^* on $[r]$ is also feasible. Fix now an arbitrary feasible 3-colouring of $\binom{[r^*]}{\leq 2}$, which assigns the same colour, say colour i , to all the singletons in $[r^*]$. Then due to (P2), colour i cannot appear on $\binom{[r^*]}{2}$, and so $r^* \leq R(a_j, a_k) - 1 \leq r$, where $\{j, k\} = [3] \setminus \{i\}$.

For $i \in [3]$, let V_i be the set of vertices with colour i . Thus we have a partition $[r] = V_1 \cup V_2 \cup V_3$. For $i, j \in [3]$, let $\omega_j(V_i)$ be the size of the largest edge-monochromatic clique of colour j in V_i .

Observe the following properties that hold for every triple $i, j, \ell \in [3]$ of distinct indices, i.e., for $\{i, j, \ell\} = [3]$. By (P2), the colour of every edge inside V_i is either j or ℓ while all the edges going between V_j and V_ℓ have colour i . By the latter property and (P1), we have

$$\omega_i(V_\ell) + \omega_i(V_j) \leq a_i - 1 \quad \text{and} \quad V_j \neq \emptyset \Rightarrow \omega_i(V_\ell) \leq a_i - 2. \quad (16)$$

For notational convenience, define $r(n_1, \dots, n_k) := R(n_1, \dots, n_k) - 1$ to be one less than the Ramsey number (i.e. it is the maximum order of a clique admitting a $(K_{n_1}, \dots, K_{n_k})$ -free edge-colouring). By the definition of $\omega_j(V_i)$, we also have

$$|V_i| \leq r(\omega_j(V_i) + 1, \omega_\ell(V_i) + 1). \quad (17)$$

Also, we will use the following trivial inequalities involving Ramsey numbers that hold for arbitrary integers $a, b, c \geq 2$: $r(a, b) + r(a, c) \leq r(a, b + c - 1)$ and $r(a, b) < r(a + 1, b)$.

First, let us derive the contradiction from assuming that each colour $i \in [3]$ appears on at least one singleton, that is, each V_i is non-empty. In order to reduce the number of cases, we allow to swap colours 1 and 2 to ensure that $\omega_1(V_2) \geq \omega_2(V_1)$. Thus we do not stipulate now which of a_1 and a_2 is larger. Observe that

$$\begin{aligned} |V_1| + |V_2| &\stackrel{(17)}{\leq} r(\omega_2(V_1) + 1, \omega_3(V_1) + 1) + r(\omega_1(V_2) + 1, \omega_3(V_2) + 1) \\ &\leq r(\omega_1(V_2) + 1, \omega_3(V_1) + 1) + r(\omega_1(V_2) + 1, \omega_3(V_2) + 1) \\ &\leq r(\omega_1(V_2) + 1, \omega_3(V_1) + \omega_3(V_2) + 1) \stackrel{(16)}{\leq} r(\omega_1(V_2) + 1, a_3). \end{aligned} \quad (18)$$

Hence, we get

$$\begin{aligned} r = |V_1| + |V_2| + |V_3| &\stackrel{(17),(18)}{\leq} r(\omega_1(V_2) + 1, a_3) + r(\omega_1(V_3) + 1, \omega_2(V_3) + 1) \\ &\stackrel{(16)}{\leq} r(\omega_1(V_2) + 1, a_3) + r(\omega_1(V_3) + 1, a_2 - 1) \\ &< r(\omega_1(V_2) + 1, a_3) + r(\omega_1(V_3) + 1, a_3) \\ &\leq r(\omega_1(V_2) + \omega_1(V_3) + 1, a_3) \stackrel{(16)}{\leq} r(a_1, a_3) \leq r. \end{aligned}$$

The above contradiction shows that, for some $\ell \in [3]$, the part V_ℓ is empty. Let $\{i, j, \ell\} = [3]$; thus $[r] = V_i \cup V_j$. It remains to derive a contradiction by assuming that each of V_i and V_j is non-empty. By the symmetry between i and j , we can assume that $\omega_j(V_i) \geq \omega_i(V_j)$. Then we have

$$\begin{aligned} r = |V_i| + |V_j| &\stackrel{(17)}{\leq} r(\omega_j(V_i) + 1, \omega_\ell(V_i) + 1) + r(\omega_i(V_j) + 1, \omega_\ell(V_j) + 1) \\ &\leq r(\omega_j(V_i) + 1, \omega_\ell(V_i) + 1) + r(\omega_j(V_i) + 1, \omega_\ell(V_j) + 1) \\ &\leq r(\omega_j(V_i) + 1, \omega_\ell(V_i) + \omega_\ell(V_j) + 1) \\ &\stackrel{(16)}{\leq} r(a_j - 1, a_\ell) < r(a_j, a_\ell) \leq r, \end{aligned}$$

which is the desired contradiction that finishes the proof of Theorem 1.8. \square

Next, let us present the proof that $r^*(3, 3, 3, 3) = 16$, the only non-trivial 4-colour case that we can solve.

Proof of Theorem 1.9. Let $\xi : \binom{[16]}{\leq 2} \rightarrow [4]$ be an arbitrary feasible colouring. It is enough to show that all singletons in $[16]$ get the same colour. For every $i \in [4]$, let V_i denote the set of vertices of colour i . Suppose there are at least two different colours on the vertices, say $V_3, V_4 \neq \emptyset$. As 5 does not divide 16, there exists at least one class, say V_3 , of size not divisible by 5, i.e., $|V_3| \not\equiv 0 \pmod{5}$. Choose an arbitrary vertex $v \in V_4$. Since ξ is a feasible colouring, by (P2) the edges

incident to v cannot have colour $\xi(v) = 4$. We can then partition $[16] \setminus \{v\} = \cup_{j \in [3]} W_j$, where $W_j := \{u : \xi(uv) = j\}$. Let $j \in [3]$. By (P1) and (P2), colour j is forbidden in $\binom{W_j}{\leq 2}$. Then by Theorem 1.8, $|W_j| \leq r^*(K_3, K_3, K_3) = R(3, 3) - 1 = 5$. Since $\sum_{j \in [3]} |W_j| = 15$, we have that $|W_j| = 5$ for every $j \in [3]$. Again by Theorem 1.8, all vertices in W_j should have the same colour. Recall that $v \in V_4$, so $V_3 \subseteq \cup_{j \in [3]} W_j$ and consequently V_3 is the union of some W_j 's. This contradicts $|V_3| \not\equiv 0 \pmod{5}$. \square

7 Concluding remarks

- As pointed out by a referee, the function $\text{nim}(n; H_1, \dots, H_k)$ is related to that of $\text{ex}^r(n; H_1, \dots, H_r)$, which is the maximum size of an n -vertex graph G that can be r -edge-coloured so that the i -th colour is H_i -free for all $i \in [r]$. Indeed, we have the following lower bound:

$$\text{nim}(n; H_1, \dots, H_k) \geq \max_{i \in [k]} \text{ex}^{k-1}(H_1, \dots, H_{i-1}, H_{i+1}, \dots, H_k).$$

It is not inconceivable that the equality holds above if $n \geq n_0(H_1, \dots, H_k)$. Theorems 1.7 and 1.11 give classes of instances, when we have equality above. We refer the readers to Section 5.3 of [25] for more on the function ex^r .

- The Ramsey variant r^* introduced here is related to the version of Ramsey numbers studied by Gyárfás, Lehel, Schelp and Tuza [24]. In particular, Proposition 5 in [24] states that $r^*(K_3, K_3, K_3, K_3) = 16$, which is the consequence of the fact that (K_3, K_3, K_3, K_3) is nice from Theorem 1.9.
- We prove in Theorem 1.13 that for any tree T , $\text{nim}_k(n; T) = (k-1)\text{ex}(n, T) + O_T(1)$. Let T be an $(h+1)$ -vertex tree and suppose that the Erdős-Sós conjecture holds, i.e. $\text{ex}(n, T) \leq (h-1)n/2$. Then for each $n \geq k^2 h^2$ with $h|n$, we can get rid of the additive error term in the lower bound, namely, it holds that $\text{nim}_k(n; T) \geq (k-1)\text{ex}(n, T)$. This directly follows from known results on graph packings. We present here a short self-contained proof (with a worse bound on n). Let F be the disjoint union of n/h copies of K_h . Let $f_i : V(F) \rightarrow [n]$, $i \in [k-1]$, be $k-1$ arbitrary injective maps and let F_i be the graph obtained by mapping F on $[n]$ via f_i . It suffices to show that we can modify f_i 's to have $E(F_i) \cap E(F_j) = \emptyset$ for any $ij \in \binom{[k-1]}{2}$. Indeed, then the lower bound is witnessed by colouring $e \in E(K_n)$ with colour- i if $e \in E(F_i)$, for each $i \in [k-1]$, and with colour- k otherwise. Suppose that there is a ‘‘conflict’’ $uv \in E(F_i) \cap E(F_j)$. Let $F^* := \cup_{i \in [k-1]} F_i$. Note that $\Delta(F^*) \leq (k-1)(h-1)$. As $n > \Delta(F^*)^2 + 1$, there exists a vertex w that is at distance at least 3 from v . We claim that switching v and w in f_i will remove all conflicts at v and w . If true, one can then repeat this process till all conflicts are removed to get the desired f_i 's. Indeed, suppose that after switching v and w , there is a conflict $wz \in E(F_i) \cap E(F_\ell)$ for some $z \in N_{F_i}(v)$ and $\ell \in [k-1] \setminus \{i\}$. Then w, z, v form a path of length 2 in F^* , contradicting the choice of w .

It would be interesting to prove a matching upper bound, i.e. to show that

$$\text{nim}_k(n; T) = (k-1)\text{ex}(n, T)$$

for every tree T and sufficiently large n . Note that equality above need not be true when T is a forest. Indeed, consider M_2 , the disjoint union of two edges. Recall that $\text{ex}(n, M_2) = n-1$.

For any $k \geq 3$ and $n \geq 4k$, we have that $\text{nim}_k(n; M_2) = (k-1)\text{ex}(n, M_2) - \frac{1}{2}(k-1)(k-2)$. Indeed, for any k -edge-colouring ϕ of K_n , one colour class, say colour-1, has size at least $\binom{n}{2}/k \geq 2(n-1)$. As every edge share endpoints with at most $2n-4$ other edges, we see that every colour-1 edge is in a copy of M_2 . Thus, $\text{nim}(\phi) \leq \text{ex}^{k-1}(n, M_2, \dots, M_2) = \sum_{i=0}^{k-2} (n-1-i) = (k-1)\text{ex}(n, M_2) - \frac{1}{2}(k-1)(k-2)$, as desired.

Acknowledgement

We would like to thank the referees for pointing out a flaw in the original draft, and for their careful reading, which has greatly improved the presentation of this paper.

References

- [1] BURR, S. A., ERDŐS, P., AND LOVÁSZ, L. On graphs of Ramsey type. *Ars Combinatoria* 1 (1976), 167–190.
- [2] CONLON, D. On the Ramsey multiplicity of complete graphs. *Combinatorica* 32 (2012), 171–186.
- [3] CONLON, D., FOX, J., AND SUDAKOV, B. An approximate version of Sidorenko’s conjecture. *Geom. Func. Analysis* 20 (2010), 1354–1366.
- [4] CONLON, D., FOX, J., AND SUDAKOV, B. Recent developments in graph Ramsey theory. In *Surveys in combinatorics 2015*, vol. 424 of *London Math. Soc. Lecture Note Ser.* Cambridge Univ. Press, Cambridge, 2015, pp. 49–118.
- [5] CONLON, D., KIM, J. H., LEE, C., AND LEE, J. Some advances on Sidorenko’s conjecture. E-print arxiv:1510.06533, 2015.
- [6] CONLON, D., AND LEE, J. Finite reflection groups and graph norms. *Adv. Math.* 315 (2017), 130–165.
- [7] CUMMINGS, J., KRÁL’, D., PFENDER, F., SPERFELD, K., TREGLOWN, A., AND YOUNG, M. Monochromatic triangles in three-coloured graphs. *J. Combin. Theory (B)* 103 (2013), 489–503.
- [8] ERDŐS, P. Some theorems on graphs. *Riveon Lematematika* 9 (1955), 13–17.
- [9] ERDŐS, P. On a theorem of Rademacher-Turán. *Illinois J. Math.* 6 (1962), 122–127.
- [10] ERDŐS, P. On the number of complete subgraphs contained in certain graphs. *Magyar Tud. Akad. Mat. Kutató Int. Közl.* 7 (1962), 459–464.
- [11] ERDŐS, P. Some recent results on extremal problems in graph theory. Results. In *Theory of Graphs (Internat. Sympos., Rome, 1966)*. Gordon and Breach, New York, 1967, pp. 117–123 (English); pp. 124–130 (French).
- [12] ERDŐS, P. Some recent problems and results in graph theory. *Discrete Math.* 164 (1997), 81–85.

- [13] ERDŐS, P., FAUDREE, R. J., AND ROUSSEAU, C. C. Extremal problems involving vertices and edges on odd cycles. *Discrete Math.* 101 (1992), 23–31.
- [14] ERDŐS, P., AND SIMONOVITS, M. A limit theorem in graph theory. *Stud. Sci. Math. Hungar.* 1 (1966), 51–57.
- [15] ERDŐS, P., AND STONE, A. H. On the structure of linear graphs. *Bull. Amer. Math. Soc.* 52 (1946), 1087–1091.
- [16] ERDŐS, P., AND SZEKERES, G. A combinatorial problem in geometry. *Compositio Math* 2 (1935), 463–470.
- [17] FRANEK, F., AND RÖDL, V. Ramsey problem on multiplicities of complete subgraphs in nearly quasirandom graphs. *Graphs Combin.* 8 (1992), 299–308.
- [18] FÜREDI, Z. New asymptotics for bipartite Turán numbers. *J. Combin. Theory (A)* 75 (1996), 141–144.
- [19] FÜREDI, Z., AND MALEKI, Z. The minimum number of triangular edges and a symmetrization method for multiple graphs. *Combin. Probab. Comput.* 26 (2017), 525–535.
- [20] FÜREDI, Z., AND SIMONOVITS, M. The history of degenerate (bipartite) extremal graph problems. In *Erdős centennial*, vol. 25 of *Bolyai Soc. Math. Stud.* János Bolyai Math. Soc., Budapest, 2013, pp. 169–264.
- [21] GIRAUD, G. Sur le problème de Goodman pour les quadrangles et la majoration des nombres de Ramsey. *J. Combin. Theory (B)* 27 (1979), 237–253.
- [22] GRUSLYS, V., AND LETZTER, S. Minimising the number of triangular edges. E-print arxiv:1605.00528; accepted by *Combin. Probab. Comput.*, 2018.
- [23] GRZESIK, A., HU, P., AND VOLEC, J. Minimum number of edges that occur in odd cycles. E-print arxiv:1605.09055, 2016.
- [24] GYÁRFÁS, A., LEHEL, J., SCHELP, R. H., AND TUZA, Z. Ramsey numbers for local colorings. *Graphs Combin.* 3 (1987), 267–277.
- [25] HANCOCK, R., STADEN, K., AND TREGLOWN, A. Independent sets in hypergraphs and ramsey properties of graphs and the integers. E-print arXiv:1701.04754v2, 2017.
- [26] HATAMI, H. Graph norms and Sidorenko’s conjecture. *Israel J. Math.* 175 (2010), 125–150.
- [27] JAGGER, C. N., ŠŤOVÍČEK, P., AND THOMASON, A. Multiplicities in subgraphs. *Combinatorica* 16 (1996), 123–141.
- [28] KEEVASH, P. Hypergraph Turán problem. In *Surveys in Combinatorics*, R. Chapman, Ed., vol. 392 of *London Math. Soc. Lecture Notes Ser.* Cambridge Univ. Press, 2011, pp. 83–140.
- [29] KEEVASH, P., AND SUDAKOV, B. On the number of edges not covered by monochromatic copies of a fixed graph. *J. Combin. Theory (B)* 90 (2004), 41–53.

- [30] KIM, J. H., LEE, C., AND LEE, J. Two approaches to Sidorenko’s conjecture. *Trans. Amer. Math. Soc.* 368 (2016), 5057–5074.
- [31] KOMLÓS, J., AND SIMONOVITS, M. Szemerédi’s regularity lemma and its applications to graph theory. In *Combinatorics, Paul Erdős is Eighty*, D. Miklós, V. T. Sós, and T. S. onyi, Eds., vol. 2. Bolyai Math. Soc., 1996, pp. 295–352.
- [32] LI, J. L. X., AND SZEGEDY, B. On the logarithmic calculus and Sidorenko’s conjecture. E-print arxiv:1107.1153; accepted by *Combinatorica*, 2018.
- [33] MA, J. On edges not in monochromatic copies of a fixed bipartite graph. *J. Combin. Theory (B)* 123 (2017), 240–248.
- [34] MUBAYI, D. Counting substructures I: Color critical graphs. *Adv. Math.* 225 (2010), 2731–2740.
- [35] NIKIFOROV, V. The number of cliques in graphs of given order and size. *Trans. Amer. Math. Soc.* 363 (2011), 1599–1618.
- [36] PIKHURKO, O., AND YILMA, Z. Supersaturation problem for color-critical graphs. *J. Combin. Theory (B)* 123 (2017), 148–185.
- [37] PYBER, L. Clique covering of graphs. *Combinatorica* 6 (1986), 393–398.
- [38] RADZISZOWSKI, S. Small Ramsey numbers. *Electronic J. Combin.* DS1 (2017), 104pp.
- [39] RAMSEY, F. P. On a problem of formal logic. *Proc. London Math. Soc.* 30 (1930), 264–286.
- [40] RAZBOROV, A. On the minimal density of triangles in graphs. *Combin. Probab. Comput.* 17 (2008), 603–618.
- [41] REIHER, C. The clique density theorem. *Annals of Math.* 184 (2016), 683–707.
- [42] SIMONOVITS, M. A method for solving extremal problems in graph theory, stability problems. In *Theory of Graphs (Proc. Colloq., Tihany, 1966)*. Academic Press, 1968, pp. 279–319.
- [43] SPERFELD, K. On the minimal monochromatic K_4 -density. E-print arxiv:1106.1030, 2011.
- [44] SUDAKOV, B. Recent developments in extremal combinatorics: Ramsey and Turán type problems. In *Proceedings of the International Congress of Mathematicians. Volume IV* (2010), Hindustan Book Agency, New Delhi, pp. 2579–2606.
- [45] SZEGEDY, B. An information theoretic approach to Sidorenko’s conjecture. E-print arxiv:1406.6738v3, 2014.
- [46] THOMASON, A. A disproof of a conjecture of Erdős in Ramsey theory. *J. Lond. Math. Soc.* 39 (1989), 246–255.
- [47] THOMASON, A. Graph products and monochromatic multiplicities. *Combinatorica* 17 (1997), 125–134.
- [48] TURÁN, P. On an extremal problem in graph theory (in Hungarian). *Mat. Fiz. Lapok* 48 (1941), 436–452.