

Rapidly rotating Bose-Einstein condensates*

Alexander Fetter, Stanford University

Warwick, December, 2005

1. Physics of one vortex line in harmonic trap
2. Experimental creation and detection of vortices
3. Vortex arrays in mean-field Thomas-Fermi regime
4. Vortex arrays in mean-field quantum-Hall regime
5. Behavior for $\Omega \rightarrow \omega_{\perp}$
6. Addition of quartic potential

*for general references, see [1, 2, 3, 4]

1 Physics of one vortex line in harmonic trap

Assume general axisymmetric trap potential

$$V_{\text{tr}}(\mathbf{r}) = V_{\text{tr}}(r, z) = \frac{1}{2}M (\omega_{\perp}^2 r^2 + \omega_z^2 z^2)$$

Basic idea (Bogoliubov): for weak interparticle potentials, nearly all particles remain in condensate for $T \ll T_c$

- dilute: s -wave scattering length $a_s \ll$ interparticle spacing $n^{-1/3}$
- equivalently, require $na_s^3 \ll 1$
- assume self-consistent condensate wave function $\Psi(\mathbf{r})$
- gives nonuniform condensate density $n(\mathbf{r}) = |\Psi(\mathbf{r})|^2$
- for $T \ll T_c$, normalization requires $N = \int dV |\Psi(\mathbf{r})|^2$

- assume an energy functional

$$E[\Psi] = \int dV \left[\underbrace{\Psi^* (\mathcal{T} + V_{\text{tr}}) \Psi}_{\text{harmonic oscillator}} + \underbrace{\frac{1}{2}g|\Psi|^4}_{\text{2-body term}} \right],$$

where $\mathcal{T} = -\hbar^2\nabla^2/2M$ is kinetic energy operator and $g = 4\pi a_s\hbar^2/M$ is interaction coupling parameter

- balance of kinetic energy $\langle \mathcal{T} \rangle$ and trap energy $\langle V_{\text{tr}} \rangle$ gives mean oscillator length $d_0 = \sqrt{\hbar/M\omega_0}$ where $\omega_0 = (\omega_{\perp}^2\omega_z)^{1/3}$ is geometric mean
- balance of kinetic energy $\langle \mathcal{T} \rangle$ and interaction energy $\langle gn \rangle$ gives healing length

$$\xi = \frac{\hbar}{\sqrt{2Mgn}} = \frac{1}{\sqrt{8\pi a_s n}}$$

- with fixed normalization and μ the chemical potential, variation of $E[\Psi]$ gives Gross-Pitaevskii (GP) eqn

$$(\mathcal{T} + V_{\text{tr}} + g|\Psi|^2) \Psi = \mu\Psi$$

- can interpret nonlinear term as a Hartree potential $V_H(\mathbf{r}) = gn(\mathbf{r})$, giving interaction with nonuniform condensate density

- generalize to time-dependent GP equation

$$i\hbar\frac{\partial\Psi}{\partial t} = (\mathcal{T} + V_{\text{tr}} + V_H) \Psi$$

- this result implies that stationary solutions have time dependence $\exp(-i\mu t/\hbar)$

Introduce hydrodynamic variables

- write $\Psi(\mathbf{r}, t) = |\Psi(\mathbf{r}, t)| \exp[iS(\mathbf{r}, t)]$ with phase S
- condensate density is $n(\mathbf{r}, t) = |\Psi(\mathbf{r}, t)|^2$
- current is

$$\mathbf{j} = \frac{\hbar}{2Mi} [\Psi^* \nabla \Psi - \Psi \nabla \Psi^*] = |\Psi|^2 \frac{\hbar \nabla S}{M}$$

- identify last factor as velocity $\mathbf{v} = \hbar \nabla S / M$
- note that \mathbf{v} is irrotational so $\nabla \wedge \mathbf{v} = 0$

- general property: circulation around contour \mathcal{C} is

$$\oint_{\mathcal{C}} d\mathbf{l} \cdot \mathbf{v} = \frac{\hbar}{M} \oint_{\mathcal{C}} d\mathbf{l} \cdot \nabla S = \frac{\hbar}{M} \Delta S|_{\mathcal{C}}$$

since $\mathbf{v} = \hbar \nabla S / M$

- change of phase $\Delta S|_{\mathcal{C}}$ must be integer times 2π since Ψ is single-valued
- hence circulation in BEC is *quantized* in units of $\kappa \equiv 2\pi\hbar/M$
- rewrite time-dependent GP equation in terms of $|\Psi|$ and S

– imaginary part gives conservation of particles

$$\frac{\partial n}{\partial t} + \nabla \cdot (n\mathbf{v}) = 0$$

– real part gives generalized Bernoulli equation

Introduction of harmonic trap yields much richer system than a uniform interacting Bose gas [5]

- trap gives new *energy* scale $\hbar\omega_0$ and new *length* scale $d_0 = \sqrt{\hbar/M\omega_0}$
- assume repulsive interactions with $a_s > 0$
- trap leads to new dimensionless parameter Na_s/d_0
- typical value of ratio: $a_s/d_0 \sim 10^{-3}$
- Na_s/d_0 is large for typical $N \sim 10^6$
- repulsive interactions expand the condensate to mean radius R_0 that exceeds d_0
- neglect radial gradient of Ψ when $Na_s/d_0 \gg 1$
- GP equation simplifies and gives local density

$$\frac{4\pi a_s \hbar^2}{M} |\Psi(r, z)|^2 = \mu - V_{\text{tr}}(r, z)$$

- harmonic trap gives quadratic density variation with condensate dimensions $R_j^2 = 2\mu/(M\omega_j^2)$ [called Thomas-Fermi (TF) limit]

One vortex line in trapped BEC

First assume bulk condensate with uniform density n and a single straight vortex line along z axis

- Gross and Pitaevskii [6, 7]: take condensate wave function

$$\Psi(\mathbf{r}) = \sqrt{n} e^{i\phi} f\left(\frac{r}{\xi}\right)$$

where r and ϕ are two-dimensional polar coordinates

- speed of sound is $s = \sqrt{gn/M}$
- assume $f(0) = 0$ and $f(x) \rightarrow 1$ for $x \gg 1$
- velocity has circular streamlines with $\mathbf{v} = (\hbar/Mr) \hat{\phi}$
- this is a quantized vortex line with $\oint d\mathbf{l} \cdot \mathbf{v} = 2\pi\hbar/M$
- $v \sim s$ when $r \sim \xi$, so vortex core forms by cavitation
- equivalently, centrifugal barrier gives vortex core of radius ξ

Static behavior of straight vortex line in a trap

Axisymmetric trap with $V_{\text{tr}}(r, z) = \frac{1}{2}M (\omega_{\perp}^2 r^2 + \omega_z^2 z^2)$

- If $\omega_z/\omega_{\perp} \gg 1$, strong axial confinement gives disk-shaped condensate
- If $\omega_z/\omega_{\perp} \ll 1$, strong radial confinement gives cigar-shaped condensate
- for vortex on axis, condensate wave function is

$$\Psi(\mathbf{r}, z) = e^{i\phi} |\Psi(r, z)|$$


- velocity is $\mathbf{v} = (\hbar/Mr)\hat{\phi}$, like uniform condensate
- centrifugal energy again forces wave function to vanish for $r \lesssim \xi$
- density is now toroidal; hole along symmetry axis
- TF limit: separated length scales with

$$\begin{aligned} \xi \text{ (vortex core)} &\ll d_0 \text{ (mean oscillator length)} \\ d_0 \text{ (mean oscillator length)} &\ll R_0 \text{ (mean condensate radius)} \end{aligned}$$

- hence TF density is essentially unchanged by vortex

Energy of rotating TF condensate with one vortex

- use density of vortex-free TF condensate; cut off the logarithmic divergence at core radius ξ
- if condensate is in rotational equilibrium at angular velocity $\boldsymbol{\Omega}$, the appropriate energy functional is [8] $E'[\Psi] = E[\Psi] - \boldsymbol{\Omega} \cdot \mathbf{L}[\Psi]$ where \mathbf{L} is the angular momentum
- let E'_0 be energy of rotating vortex-free condensate
- let $E'_1(r_0, \Omega)$ be energy of a rotating condensate with straight vortex that is displaced laterally by distance r_0 from symmetry axis
- approximation of straight vortex works best for disk-shaped condensate ($\omega_z \gtrsim \omega_\perp$)
- Difference of these two energies is energy associated with formation of vortex $\Delta E'(r_0, \Omega) = E'_1(r_0, \Omega) - E'_0$
- $\Delta E'(r_0, \Omega)$ depends on position r_0 of vortex and on Ω



Plot $\Delta E'(r_0, \Omega)$ as function of ζ_0 for various fixed Ω [9], where $\zeta_0 = r_0/R_0$ is scaled displacement from center

curve (a) is $\Delta E'(r_0, \Omega)$ for $\Omega = 0$

- $\Delta E'(r_0, 0)$ decreases monotonically with increasing ζ_0
- curvature is negative at $\zeta_0 = 0$
- for no dissipation, fixed energy means constant ζ_0
- only allowed motion is uniform precession at fixed r_0
- angular velocity is given by variational Lagrangian method [10, 11, 3] $\dot{\phi}_0 \propto -\partial E(r_0)/\partial r_0$
- precession arises from nonuniform trap potential (*not image vortex*) and nonuniform condensate density
- in presence of weak dissipation, vortex moves to lower energy and slowly spirals outward

As Ω increases, curvature near $\zeta_0 = 0$ decreases

- curve (b) is when curvature near $\zeta_0 = 0$ vanishes
- it corresponds to angular velocity

$$\Omega_m = \frac{3}{2} \frac{\hbar}{M R_\perp^2} \ln \left(\frac{R_\perp}{\xi} \right)$$

- for $\Omega \gtrsim \Omega_m$, energy $\Delta E'(\zeta_0, \Omega)$ has local *minimum* near $\zeta_0 = 0$
- dissipation would now drive vortex back *toward* the symmetry axis
- Ω_m is angular velocity for onset of *metastability*
- vortex at center is *locally* stable for $\Omega > \Omega_m$, but not *globally* stable, since $\Delta E'(0, \Omega_m)$ is positive

Mean vortex density in a rotating superfluid (Feynman)


- solid-body rotation has $\mathbf{v}_{\text{sb}} = \boldsymbol{\Omega} \wedge \mathbf{r}$
- \mathbf{v}_{sb} has constant vorticity $\nabla \wedge \mathbf{v}_{\text{sb}} = 2\boldsymbol{\Omega}$
- each quantized vortex at \mathbf{r}_j has localized vorticity

$$\nabla \wedge \mathbf{v} = \frac{2\pi\hbar}{M} \delta^{(2)}(\mathbf{r} - \mathbf{r}_j) \hat{\mathbf{z}}$$

- assume \mathcal{N}_v vortices uniformly distributed in area \mathcal{A} bounded by contour \mathcal{C}
- circulation around \mathcal{C} is $\mathcal{N}_v \times 2\pi\hbar/M$
- but circulation in \mathcal{A} is also $2\boldsymbol{\Omega}\mathcal{A}$
- hence vortex density is $n_v = \mathcal{N}_v/\mathcal{A} = M\boldsymbol{\Omega}/\pi\hbar$
- area per vortex $1/n_v$ is $\pi\hbar/M\boldsymbol{\Omega} \equiv \pi l^2$ which defines radius $l = \sqrt{\hbar/M\boldsymbol{\Omega}}$ of circular cell
- intervortex spacing $\sim 2l$ decreases like $1/\sqrt{\boldsymbol{\Omega}}$
- analogous to quantized flux lines (charged vortices) in type-II superconductors

2 Experimental creation/detection of vortices in dilute trapped BEC

- first vortex made at JILA (1999) [12]
- used nearly spherical ^{87}Rb condensate containing two different hyperfine components
- use coherent (Rabi) process to control interconversion between two components
- spin up condensate by coupling the two components with a stirring perturbation
- turn off coupling, leaving one component with trapped quantized vortex surrounding nonrotating core of other component
- use selective tuning to make nondestructive image of either component



- study precession of this vortex with filled core around trap center
- can also create vortex with empty core [13]
 - theory predicts $\dot{\phi}/2\pi \approx 1.58 \pm 0.16$ Hz, and
 - experiment finds $\dot{\phi}/2\pi \approx 1.8 \pm 0.1$ Hz
- see no outward radial motion for ~ 1 s, so dissipation is small on this time scale


École Normale Supérieure (ENS) in Paris studied vortex creation in elongated rotating cigar-shaped condensate with one component [14, 15]

- used off-center toggled rotating laser beam to deform the transverse trap potential and stir the condensate at an applied frequency $\Omega/2\pi \lesssim 200$ Hz




- find vortex appears at a critical frequency $\Omega_c \approx 0.7\omega_{\perp}$ (detected by expanding the condensate, which now has a disk shape, with vortex core as expanded hole)
- vortex nucleation is dynamical process associated with surface instability (quadrupole oscillation)


- ENS group observed small vortex arrays of up to 11 vortices (arranged in two concentric rings)



- like patterns predicted and seen in superfluid ^4He [16]




- MIT group has prepared considerably larger rotating condensates in less elongated trap
- they have observed triangular vortex lattices with up to 130 vortices [17]




- like Abrikosov lattice of quantized flux lines (which are charged vortices) in type-II superconductors
- JILA group has now made large rotating condensates with several hundred vortices and angular velocity $\Omega/\omega_{\perp} \approx 0.995$ [18]
- these rapidly rotating systems open many exciting new possibilities (discussed below)

Very recently, Zwierlein *et al.* (MIT) have studied ${}^6\text{Li}$ atoms (*fermions*) in optical dipole trap [19]

- by tuning an external magnetic field, can change the scattering length (Feshbach resonance)
- regime of bound molecules of fermions ($a_s > 0$): these “bosonic” molecules can undergo BEC
- then rotate using ENS laser-beam technique
- find vortex lattice that slowly decays



- move from BEC region ($1/k_F a_s > 0$) across resonant region to BCS region ($1/k_F a_s < 0$) of unbound but attracting fermion pairs



- vortex lattice persists and survives across resonance into fermionic regime (not yet into BCS regime of weakly bound overlapping Cooper pairs)
- provides clear evidence for macroscopic coherence and superfluidity of condensed fermionic cold gas

3 Vortex arrays in mean-field Thomas-Fermi regime

As Ω increases, the mean vortex density $n_v = M\Omega/\pi\hbar$ increases linearly following the Feynman relation

- in addition, centrifugal forces expand the condensate radially, so that the area πR_\perp^2 also increases
- hence the number of vortices $\mathcal{N}_v = n_v \pi R_\perp^2 = M\Omega R_\perp^2 / \hbar$ increases faster than linearly with Ω
- conservation of particles implies that the condensate also shrinks axially
- TF approximation assumes that interaction energy $\langle g|\Psi|^4 \rangle$ and trap energy $\langle V_{\text{tr}}|\Psi|^2 \rangle$ are large relative to kinetic energy for density variations $(\hbar^2/M)\langle (\nabla|\Psi|)^2 \rangle$
- expansion of condensate means that central density eventually becomes small and TF picture fails

Quantitative description of rotating TF condensate

Kinetic energy of condensate involves

$$\frac{\hbar^2}{2M} \int dV |\nabla \Psi|^2 = \underbrace{\int dV \frac{1}{2} M v^2 |\Psi|^2}_{\text{superflow energy}} + \underbrace{\frac{\hbar^2}{2M} \int dV (\nabla |\Psi|)^2}_{\text{density variation}}$$

where $\Psi = \exp(iS)|\Psi|$ and $\mathbf{v} = \hbar \nabla S / M$ is flow velocity

- generalized TF approximation: retain the energy of superflow but ignore the energy from density variation
- this approximation will fail eventually when vortex lattice becomes dense and cores start to overlap
- in rotating frame, generalized TF energy functional is

$$E'[\Psi] = \int dV \left[\left(\frac{1}{2} M v^2 + V_{\text{tr}} - M \boldsymbol{\Omega} \cdot \mathbf{r} \wedge \mathbf{v} \right) |\Psi|^2 + \frac{1}{2} g |\Psi|^4 \right]$$

- here, \mathbf{v} is flow velocity generated by all the vortices

For $\boldsymbol{\Omega}$ along z , can rewrite $E'[\Psi]$ as

$$E'[\Psi] = \int dV \left[\frac{1}{2}M (\mathbf{v} - \boldsymbol{\Omega} \wedge \mathbf{r})^2 |\Psi|^2 + \frac{1}{2}M\omega_z^2 z^2 |\Psi|^2 + \frac{1}{2} (\omega_\perp^2 - \Omega^2) r^2 |\Psi|^2 + \frac{1}{2}g|\Psi|^4 \right]$$

- in the rotating frame, the dominant effect of the dense vortex array is that spatially averaged flow velocity $\langle \mathbf{v} \rangle$ is close to $\boldsymbol{\Omega} \wedge \mathbf{r} = \mathbf{v}_{\text{sb}}$
- hence can ignore first term in $E'[\Psi]$, giving

$$E'[\Psi] \approx \int dV \left[\frac{1}{2}M\omega_z^2 z^2 |\Psi|^2 + \frac{1}{2} (\omega_\perp^2 - \Omega^2) r^2 |\Psi|^2 + \frac{1}{2}g|\Psi|^4 \right]$$

- E' now looks exactly like TF energy for nonrotating condensate but with a *reduced* radial trap frequency $\omega_\perp^2 \rightarrow \omega_\perp^2 - \Omega^2$

Now have TF wave function that depends explicitly on Ω through the altered radial trap frequency $\omega_{\perp}^2 \rightarrow \omega_{\perp}^2 - \Omega^2$

$$|\Psi(r, z)|^2 = n(0) \left(1 - \frac{r^2}{R_{\perp}^2} - \frac{z^2}{R_z^2} \right)$$

where $R_{\perp}^2 = 2\mu/[M(\omega_{\perp}^2 - \Omega^2)]$ and $R_z^2 = 2\mu/M\omega_z^2$

- for pure harmonic trap, must have $\Omega < \omega_{\perp}$ to retain radial confinement
- normalization $\int dV |\Psi|^2 = N$ shows that

$$\frac{\mu(\Omega)}{\mu(0)} = \left(1 - \frac{\Omega^2}{\omega_{\perp}^2} \right)^{2/5}$$

in three dimensions

- central density given by $n(0) = \mu(\Omega)/g$
- $n(0)$ decreases with increasing Ω because of reduced radial confinement

- TF formulas for condensate radii show that


$$\frac{R_z(\Omega)}{R_z(0)} = \left(1 - \frac{\Omega^2}{\omega_\perp^2}\right)^{1/5}, \quad \frac{R_\perp(\Omega)}{R_\perp(0)} = \left(1 - \frac{\Omega^2}{\omega_\perp^2}\right)^{-3/10}$$

confirming axial shrinkage and radial expansion

- aspect ratio changes

$$\frac{R_z(\Omega)}{R_\perp(\Omega)} = \frac{R_z(0)}{R_\perp(0)} \left(1 - \frac{\Omega^2}{\omega_\perp^2}\right)^{1/2}$$

- this last effect provides an important diagnostic tool to determine actual angular velocity Ω [20, 18]



- measured aspect ratio [18] indicates that Ω/ω_\perp can become as large as ≈ 0.993

How uniform is the vortex array?

The analysis of the TF density profile $|\Psi_{TF}|^2 = n_{TF}$ in the rotating condensate assumed that the flow velocity \mathbf{v} was precisely the solid-body value $\mathbf{v}_{sb} = \mathbf{\Omega} \wedge \mathbf{r}$

- this led to the cancellation of the contribution

$$\int dV (\mathbf{v} - \mathbf{\Omega} \wedge \mathbf{r})^2 n_{TF}$$

in the TF energy functional

- a more careful study [21] shows that there is a small nonuniformity in the vortex lattice
- specifically, each regular vortex lattice position vector \mathbf{r}_j experiences a small displacement field $\mathbf{u}(\mathbf{r})$, so that $\mathbf{r}_j \rightarrow \mathbf{r}_j + \mathbf{u}(\mathbf{r}_j)$
- as a result, the two-dimensional vortex density changes to

$$n_v(\mathbf{r}) \approx \bar{n}_v (1 - \nabla \cdot \mathbf{u})$$

where $\bar{n}_v = M\Omega/\pi\hbar$ is the uniform Feynman value

- variation with respect to \mathbf{u} yields an Euler-Lagrange equation that can be solved to give

$$\mathbf{u}(\mathbf{r}) \approx \frac{\bar{l}^2}{4R_{\perp}^2} \ln\left(\frac{\bar{l}^2}{\xi^2}\right) \frac{\mathbf{r}}{1 - r^2/R_{\perp}^2}$$


where $\bar{l}^2 = 1/\pi\bar{n}_v$ can be taken as the mean circular cell radius inside the slowly varying logarithm

- the deformation of the regular vortex lattice is purely radial (as expected from symmetry)
- R_{\perp}^2/\bar{l}^2 is the number of vortices \mathcal{N}_v in the rotating condensate, so that the nonuniform distortion is small, of order $1/\mathcal{N}_v$ (at most a few %), even though the TF number density n_{TF} changes dramatically near edge
- correspondingly, the vortex density becomes

$$n_v(r) \approx \bar{n}_v - \frac{1}{2\pi R_{\perp}^2} \ln\left(\frac{\bar{l}^2}{\xi^2}\right) \frac{1}{(1 - r^2/R_{\perp}^2)^2}$$

(the correction is again of order $1/\mathcal{N}_v$)

- recent JILA experiments [22] confirm these predicted small distortions for relatively dense vortex lattices



Tkachenko oscillations of the vortex lattice

Tkachenko (1966) [23] studied equilibrium arrangement of a rotating vortex array as model for superfluid ^4He

- assumed two-dim incompressible fluid with straight vortices
- showed that a triangular lattice has lowest energy in rotating frame
- small perturbations about equilibrium positions had unusual collective motion in which vortices undergo nearly transverse wave of lattice distortions (like two-dimensional transverse “phonons” in vortex lattice, but with no change in fluid density)
- for long wavelengths (small k), Tkachenko found a linear dispersion relation $\omega_k \approx c_T k$
- speed of Tkachenko wave $c_T = \sqrt{\frac{1}{4}\hbar\Omega/M} = \frac{1}{2}\hbar/M\bar{l}$, where $\bar{l} = \sqrt{\hbar/M\Omega}$ is radius of circular vortex cell



In a rotating gas, the compressibility becomes important, as shown by Sonin [24, 25] and Baym [26]

- let the speed of sound in the compressible gas be c_s
- coupling between the vortices and the compressible fluid leads to generalized dispersion relation

$$\omega^2 = c_T^2 \frac{c_s^2 k^4}{4\Omega^2 + c_s^2 k^2}$$

- if $k \gg \Omega/c_s$, recover Tkachenko's result $\omega = c_T k$ (short-wavelength incompressible limit)
- but if $k \ll \Omega/c_s$ (long wavelength), mode becomes *soft* with $\omega \propto k^2$
- Sonin [25] obtains dynamical equations for waves in a nonuniform condensate, along with appropriate boundary conditions at the outer surface
- Baym [26] uses theory for uniform condensate plus approximate boundary conditions from Anglin and Crescimanno [27]

- rough agreement with JILA experiments [28] on low-lying Tkachenko modes in rapidly rotating BEC (up to $\Omega/\omega_{\perp} \approx 0.975$)



4 Vortex arrays in mean-field quantum-Hall regime

Lowest-Landau-Level (quantum-Hall) behavior

When the vortex cores overlap, kinetic energy associated with density variation around each vortex core becomes important

- hence the TF approximation breaks down (it ignores this kinetic energy from density variations)
- return to full GP energy $E'[\Psi]$ in the rotating frame.
- in this limit of rapid rotations ($\Omega \lesssim \omega_{\perp}$), Ho [29] incorporated kinetic energy *exactly*
- condensate expands and is effectively two dimensional
- for simplicity, treat a two-dimensional condensate that is *uniform* in the z direction over a length Z
- condensate wave function $\Psi(\mathbf{r}, z)$ can be written as $\sqrt{N/Z} \psi(\mathbf{r})$, where $\psi(\mathbf{r})$ is a two-dimensional wave function with unit normalization $\int d^2r |\psi|^2 = 1$

General two-dimensional energy functional in rotating frame becomes

$$E'[\psi] = \int d^2r \psi^* \left(\underbrace{\frac{p^2}{2M} + \frac{1}{2}M\omega_{\perp}^2 r^2 - \Omega L_z}_{\text{one-body oscillator } \mathcal{H}_0} + \underbrace{\frac{1}{2}g_{2D}|\psi|^2}_{\text{interaction}} \right) \psi,$$

where $\mathbf{p} = -i\hbar\nabla$, $L_z = \hat{\mathbf{z}} \cdot \mathbf{r} \times \mathbf{p}$, and $g_{2D} = Ng/Z$

One-body oscillator hamiltonian in rotating frame \mathcal{H}_0 is exactly soluble and has eigenvalues [30]

$$\epsilon_{nm} = \hbar [\omega_{\perp} + n(\omega_{\perp} + \Omega) + m(\omega_{\perp} - \Omega)]$$

where n and m are non-negative integers

- in limit $\Omega \rightarrow \omega_{\perp}$, these eigenvalues are essentially independent of m (massive degeneracy)
- n becomes the Landau level index
- lowest Landau level with $n = 0$ is separated from higher states by gap $\sim 2\hbar\omega_{\perp}$

Large radial expansion means small central density $n(0)$, so that interaction energy $gn(0)$ eventually becomes small compared to gap $2\hbar\omega_{\perp}$

Hence focus on “lowest Landau level” (LLL), with $n = 0$ and general non-negative $m \geq 0$

- LLL eigenfunctions have a very simple form

$$\psi_{0m}(\mathbf{r}) \propto r^m e^{im\phi} e^{-r^2/2d_{\perp}^2}$$

- here, $d_{\perp} = \sqrt{\hbar/M\omega_{\perp}}$ is analogous to the “magnetic length” in the Landau problem
- in terms of a complex variable $\zeta \equiv x + iy$, these LLL eigenfunctions have an extremely simple form

$$\psi_{0m} \propto \zeta^m e^{-r^2/2d_{\perp}^2}$$

with $m \geq 0$ (note that $\zeta = r e^{i\phi}$ when expressed in two-dimensional polar coordinates)

- assume that the GP wave function is a finite linear combination of these LLL eigenfunctions

$$\psi_{LLL}(\mathbf{r}) = \sum_{m \geq 0} c_m \psi_{0m}(\mathbf{r}) = f(\zeta) e^{-r^2/2d_{\perp}^2}$$

where $f(\zeta) = \sum_{m \geq 0} c_m \zeta^m$ is an *analytic function* of the complex variable ζ

- specifically, $f(\zeta)$ is a complex polynomial and thus can be factorized as $f(\zeta) = \prod_j (\zeta - \zeta_j)$ apart from overall constant
- $f(\zeta)$ vanishes at each of the points $\{\zeta_j\}$, which are the positions of the nodes of ψ_{LLL}
- in addition, phase of wave function increases by 2π whenever ζ moves around any of these zeros $\{\zeta_j\}$
- we conclude that the LLL trial solution has singly quantized vortices located at positions of zeros $\{\zeta_j\}$

- spatial variation of number density $n(\mathbf{r}) = |\psi_{LLL}(\mathbf{r})|^2$ is determined by spacing of the vortices, so that core size is comparable with the intervortex spacing $\bar{l} = \sqrt{\hbar/M\Omega}$ which is simply d_{\perp} in the limit $\Omega \approx \omega_{\perp}$
- unlike TF approximation at lower Ω , wave function ψ_{LLL} automatically includes *all* the kinetic energy
- since LLL wave functions play a crucial role in the quantum Hall effect (two-dimensional electrons in a strong magnetic field), this LLL regime has been called “mean-field quantum-Hall” limit [31]
- note that we are still in the regime governed by GP equation, so there is still a BEC
- corresponding many-body ground state is simply a Hartree product with each particle in *same* one-body solution $\psi_{LLL}(\mathbf{r})$, namely

$$\Psi_{GP}(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N) \propto \prod_{n=1}^N \psi_{LLL}(\mathbf{r}_n)$$

- this is *coherent (superfluid)* state, since a single GP state ψ_{LLL} has *macroscopic* occupation

Take this LLL trial function seriously

- for *any* LLL state ψ_{LLL} , can show that (use oscillator units with ω_{\perp} and d_{\perp} for energy and length) [29, 31, 32]

$$\int d^2r r^2 |\psi_{LLL}|^2 = 1 + \int d^2r \psi_{LLL}^* L_z \psi_{LLL}$$

- allows exact rewriting of energy functional

$$E'[\psi_{LLL}] = \Omega + \int d^2r \left[(1 - \Omega) r^2 |\psi_{LLL}|^2 + \frac{1}{2} g_{2D} |\psi_{LLL}|^4 \right]$$

- unrestricted variation would lead to inverted parabola

$$|\psi|^2 = n(r) = \frac{2}{\pi R_0^2} \left(1 - \frac{r^2}{R_0^2} \right)$$

where $\pi R_0^4 = 2g_{2D}/(1 - \Omega)$ fixes condensate radius

- looks like earlier TF profile, but here include all kinetic energy explicitly
- these results ignore vortices and violate form of ψ_{LLL}

- to include effect of vortices, study logarithm of the particle density for any LLL state
- use ψ_{LLL} to find

$$\ln n_{LLL}(\mathbf{r}) = -\frac{r^2}{d_{\perp}^2} + 2 \sum_j \ln |\mathbf{r} - \mathbf{r}_j|$$

- apply two-dimensional Laplacian: use standard result $\nabla^2 \ln |\mathbf{r} - \mathbf{r}_j| = 2\pi\delta^{(2)}(\mathbf{r} - \mathbf{r}_j)$ to obtain

$$\nabla^2 \ln n_{LLL}(\mathbf{r}) = -\frac{4}{d_{\perp}^2} + 4\pi \sum_j \delta^{(2)}(\mathbf{r} - \mathbf{r}_j)$$

- here, sum over delta functions is precisely the *vortex* density $n_v(\mathbf{r})$
- this result relates *particle* density $n_{LLL}(\mathbf{r})$ in LLL approximation to *vortex* density $n_v(\mathbf{r})$ [29, 31, 32]

$$\frac{1}{4}\nabla^2 \ln n_{LLL}(\mathbf{r}) = -\frac{1}{d_{\perp}^2} + \pi n_v(\mathbf{r})$$

- if vortex lattice is exactly uniform (so n_v is constant), then density profile is strictly Gaussian, with $n_{LLL}(\mathbf{r}) \propto \exp(-r^2/\sigma^2)$ and $\sigma^{-2} = d_{\perp}^{-2} - \pi n_v \propto \omega_{\perp} - \Omega$
- note that $\sigma^2 \gg d_{\perp}^2$
- to better minimize the energy, mean density profile \bar{n}_{LLL} should approximate inverted parabolic shape $\bar{n}_{LLL}(\mathbf{r}) \propto 1 - r^2/R_{\perp}^2$
- then find *nonuniform* vortex density with

$$n_v(r) \approx \frac{1}{\pi d_{\perp}^2} - \frac{1}{\pi R_{\perp}^2} \frac{1}{(1 - r^2/R_{\perp}^2)^2}$$

similar to result at lower Ω [21] (in both cases, small correction term is of order $\sim \mathcal{N}_v^{-1}$)

- independently, numerical work by Cooper *et al.* [33] shows that allowing the vortices in the LLL to deviate from the triangular array near the outer edge lowers the energy

5 Behavior for $\Omega \rightarrow \omega_{\perp}$

What happens beyond the “mean-field quantum Hall” regime is still subject to vigorous debate

Predict *quantum phase transition* from *coherent BEC states* to *correlated many-body states*

- define the ratio $\nu \equiv N/\mathcal{N}_v$ of the number of atoms per vortex
- because of similarities to a two-dimensional electron gas in a strong magnetic field, ν is called the “filling fraction” [34, 35]
- current experiments [18] have $N \sim 10^5$ and $\mathcal{N}_v \sim$ several hundred, so $\nu \sim$ a few hundred
- numerical studies [35] for small number of vortices ($\mathcal{N}_v \lesssim 8$) and variable N indicate that the coherent GP state is favored for $\nu \gtrsim 6$

- for smaller ν there is a sequence of *highly correlated* states similar to some known from the quantum Hall effect, in particular a bosonic version of the *Laughlin* state [35] (here $z_n = x_n + iy_n$ refers to n th particle)

$$\Psi_{\text{Lau}}(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N) \propto \prod_{n < n'}^N (z_n - z_{n'})^2 \exp\left(-\sum_{n=1}^N \frac{|z_n|^2}{2d_{\perp}^2}\right)$$

- these correlated many-body states are *qualitatively different* from coherent GP form
 - $\Psi_{GP}(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N) \propto \prod_n \psi(\mathbf{r}_n)$ is the Hartree product of N factors of *same* one-body function $\psi(\mathbf{r})$
 - the product $\prod_{n < n'} (z_n - z_{n'})^2$ in $\Psi_{\text{Lau}}(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N)$ involves $N(N-1)/2$ factors for all possible *pairs* of particles and vanishes whenever two particles are close together
 - this is the source of the correlations
 - for large N , correlated form Ψ_{Lau} is much more difficult to use

How to reach correlated regime?

- need to reduce the ratio $\nu = N/\mathcal{N}_v$ (number of atoms per vortex)
- one possibility is to use array of small condensates trapped in optical lattice
- need to rotate each condensate to a relatively high angular velocity
- several experimental groups working on this option

6 Addition of quartic potential

One way to avoid singularity when $\Omega \rightarrow \omega_{\perp}$ is to add a quartic confining potential [36, 37, 38]

- now have a total potential with quadratic and quartic terms

$$V_{\text{tr}} = \frac{1}{2}M\omega_{\perp}^2 \left(r^2 + \lambda \frac{r^4}{d_{\perp}^2} \right)$$

where the dimensionless constant λ fixes the quartic admixture

- allows access to regime $\Omega/\omega_{\perp} \geq 1$
- assume nearly uniform vortex array with $\mathbf{v} \approx \boldsymbol{\Omega} \wedge \mathbf{r}$

- studied experimentally at ENS, Paris [39], where a blue-detuned axial laser provided the weak quartic confinement ($\lambda \sim 10^{-3}$ and $\omega_{\perp}/2\pi \approx 64.8$ Hz)




FIG. 1. Pictures of the rotating gas taken along the rotation axis after 18 ms time of flight. We indicate in each picture the stirring frequency $\Omega_{\text{stir}}^{(2)}$ during the second stirring phase ($\omega_{\perp}/2\pi = 64.8$ Hz). The vertical size of each image is $306 \mu\text{m}$.

- find regular vortex lattice for $\Omega \lesssim \omega_{\perp}$
- find disordered vortex lattice for $\Omega \gtrsim \omega_{\perp}$
- near $\Omega \approx 1.05 \omega_{\perp}$, the system seems to break up
- TF theory predicts a reduced density at center, which is observed




FIG. 2. Optical thickness of the atom cloud after time of flight for $\Omega_{\text{stir}}^{(2)}/2\pi = 66$ Hz. (a) Radial distribution in the xy plane of Fig. 1(e). Continuous line: fit using the Thomas-Fermi distribution (3). (b) Distribution along the z axis averaged over $|x| < 20 \mu\text{m}$ (imaging beam propagating along y).

What is happening?

- ENS condensate is nearly spherical for $\Omega \sim \omega_{\perp}$, so three-dimensional effects are important
- they suggest repeating the experiment with strong axial confinement to see if three-dimensional effects dominate and cause instability
- GP analysis in two dimensions finds nothing like the observed break up [37, 38, 40]
- is there some sort of transition from a GP state to a highly correlated state in the regime $\Omega \gtrsim \omega_{\perp}$?
- this issue remains very uncertain

References

- [1] F. Dalfovo, S. Giorgini, L. P. Pitaevskii and S. Stringari, *Rev. Mod. Phys.* **71**, 463 (1999).
- [2] A. L. Fetter, in *Bose-Einstein Condensation in Atomic Gases*, edited by M. Inguscio, S. Stringari and C. E. Wieman (IOP Press, Amsterdam, 1999), p. 201.
- [3] A. L. Fetter and A. A. Svidzinsky, *J. Phys.: Condens. Matter* **13**, R135 (2001).
- [4] A. L. Fetter, *J. Low Temp. Phys.* **129**, 263 (2002).
- [5] G. Baym and C. J. Pethick, *Phys. Rev. Lett.* **76**, 6 (1996).
- [6] E. P. Gross, *Nuovo Cimento* **20**, 454 (1961).
- [7] L. P. Pitaevskii, *Zh. Eksp. Teor. Fiz.* **40**, 646 (1961) [*Sov. Phys. JETP* **13**, 451 (1961)].
- [8] L. D. Landau and E. M. Lifshitz, *Mechanics*, Pergamon Press, Oxford (1960); E. M. Lifshitz and L. P. Pitaevskii, *Statistical Physics*, Pergamon Press, Oxford (1980).
- [9] A. A. Svidzinsky and A. L. Fetter, *Phys. Rev. Lett.* **84**, 5919 (2000).
- [10] E. Lundh and P. Ao, *Phys. Rev. A* **61**, 063612 (2000).
- [11] S. A. McGee and M. J. Holland, *Phys. Rev. A* **63**, 043608 (2001).
- [12] M. R. Matthews, B. P. Anderson, P. C. Haljan, D. S. Hall, C. E. Wieman and E. A. Cornell, *Phys. Rev. Lett.* **83**, 2498 (1999).
- [13] B. P. Anderson, P. C. Haljan, C. E. Wieman, and E. A. Cornell, *Phys. Rev. Lett.* **85**, 2857 (2000).

- [14] K. W. Madison, F. Chevy, W. Wohlleben and J. Dalibard, Phys. Rev. Lett. **84**, 806 (2000).
- [15] K. W. Madison, F. Chevy, W. Wohlleben and J. Dalibard, J. Mod. Opt. **47**, 2725 (2000).
- [16] E. J. Yarmchuk, M. J. V. Gordon, and R. E. Packard, Phys. Rev. Lett. **79**, 214 (1979).
- [17] J. R. Abo-Shaeer, C. Raman, J. M. Vogels, and W. Ketterle, Science **292**, 476 (2001).
- [18] V. Schweikhard, I. Coddington, P. Engels, V.P. Mogendorff, and E. A. Cornell, Phys. Rev. Lett. **92**, 040404 (2004).
- [19] M. Zwierlein, C. Schunck, A. Schirotzek, C. Stan, P. Zarth, and W. Ketterle, OCTS-Workshop on Strongly Interacting Quantum Gases, Columbus, Ohio (April, 2005), unpublished; M. Zwierlein, J. R. Abo-Shaeer, A. Schirotzek, C. H. Schunck, and W. Ketterle, Nature **435**, 1047 (2005).
- [20] P. C. Haljan, I. Coddington, P. Engels, and E. A. Cornell, Phys. Rev. Lett. **87**, 210403 (2001).
- [21] D. E. Sheehy and L. Radzihovsky, Phys. Rev. A **70**, 051602(R) (2004); Phys. Rev. A **70**, 063620 (2004) .
- [22] I. Coddington, P. C. Haljan, P. Engels, V. Schweikhard, S. Tung, and E. A. Cornell, Phys. Rev. A **70**, 063607 (2004).
- [23] V. K. Tkachenko, Zh. Eksp. Teor. Fiz. **49**, 1875 (1965) [Sov. Phys. JETP **22**, 1282 (1966)]; Zh. Eksp. Teor. Fiz. **50**, 1573 (1966) [Sov. Phys. JETP **23**, 1049 (1966)]. Phys. Rev. **162**, 143 (1967). Phys. Rev. B **11**, 2049 (1975).

- [24] E. B. Sonin, Rev. Mod. Phys. **59**, 87 (1987).
- [25] E. B. Sonin, Phys. Rev. A **71**, 011603(R) (2005).
- [26] G. Baym, Phys. Rev. Lett. **91**, 110402 (2003).
- [27] J. R. Anglin and M. Crescimanno, cond-mat/0210063.
- [28] I. Coddington, P. Engels, V. Schweikhard and E. A. Cornell, Phys. Rev. Lett. **91**, 100402 (2003).
- [29] T.-L. Ho, Phys. Rev. Lett. **87**, 060403 (2001).
- [30] C. Cohen-Tannoudji, B. Diu and F. Laloë, *Quantum Mechanics* (J. Wiley & Sons, New York, 1977), Volume I, pp. 742-764.
- [31] G. Watanabe, G. Baym, and C. J. Pethick, Phys. Rev. Lett. **93**, 190401 (2004).
- [32] A. Aftalion, X. Blanc, and J. Dalibard, Phys. Rev. A **71**, 023611 (2005).
- [33] N. R. Cooper, S. Komineas, and N. Read, Phys. Rev. A **70**, 033604 (2004).
- [34] N. K. Wilkin and M. J. F. Gunn, Phys. Rev. Lett. **84**, 6 (2000).
- [35] N. R. Cooper, N. K. Wilkin, and M. J. F. Gunn, Phys. Rev. Lett. **87**, 120405 (2001).
- [36] A. L. Fetter, Phys. Rev. A **64**, 063608 (2001).
- [37] K. Kasamatsu, M. Tsubota, and M. Ueda, Phys. Rev. A **66**, 053606 (2002).
- [38] G. M. Kavoulakis and G. Baym, New J. Phys. **5**, 51.1 (2003).

- [39] V. Bretin, S. Stock, Y. Seurin, and J. Dalibard, Phys. Rev. Lett. **92**, 050403 (2004).
- [40] A. L. Fetter, B. Jackson, and S. Stringari, Phys. Rev. A **71**, 013605 (2005).