

OSCILLATORY SUMS

VOLKER BETZ, VASSILI GELFREICH AND FLORIAN THEIL

Abstract. We explore a strange class of sums with alternating, but huge terms. Like with oscillatory integrals, the final sum is much smaller than the individual terms. Unlike oscillatory integrals, we know of no easy general explanation for this effect. In one particular case however, we are able to give a somewhat surprising explanation for it.

Keywords: ...

2000 Math. Subj. Class.: ...

We all know about oscillatory integrals, where the positive and negative part of the integrand give rise to cancellations that result in a value of the integral much smaller than the values of the integrand, or in the integral being finite even though the integrand is not Lebesgue-integrable. The basic mechanism of the cancellations can be analysed and understood quite easily, and rigorously.

But what about the integrals discrete cousin, the series? As a first example, look at the sequence of finite sums $(s_n)_{n \in \mathbb{N}}$ with $s_n = \sum_{k=0}^{3n} \frac{(-n)^k}{k!}$. The sum is alternating alright, but otherwise behaves rather badly: the coefficients quickly grow in absolute value, reach an exponentially large maximum at $k = n$ and then decay to zero, being of order one around $k = en$ (all of this can be seen easily using Stirling's formula). As for the cancellation that we might hope for, the ratio between two consecutive coefficients with $k \approx n$ is e , so their difference is of the order e^n . In conclusion, when we just look at the coefficients of s_n , we have a hard time to believe that this sum could give any meaningful value.

Of course, we do know that s_n converges to zero as $n \rightarrow \infty$, as it is just the truncated Taylor series of e^{-n} , which is the value that s_n has when the upper summation index is taken to be infinity. By the Leibnitz criterion, $e^{-n} - s_n$ is bounded by the first omitted term, which for $k = 3n + 1$ is already exponentially small. Thus, after a second look there is nothing mysterious or even exceedingly interesting about s_n , despite maybe the fact that the sums are small for reasons that are not very obvious from the coefficients.

Let us make things slightly more interesting and add a perturbation. We define

$$S_n = \sum_{k=0}^n \frac{(k-n)^k}{k!}. \quad (1)$$

The coefficients show the same qualitative behaviour as those of s_n , with the important differences that this time, we do not have a Taylor series to relate S_n with a well-known function, and that taking the upper summation index to be infinity renders the expression for S_n meaningless. So, should we still expect S_n to have any sort of reasonable behaviour, or even converge as $n \rightarrow \infty$? If so, why?

Let us investigate the first question by actually calculating S_n for the first few hundred n . Surprisingly, S_n converges to zero, and very quickly so: a logarithmic plot (cf. Figure ?? a)) shows that this convergence is exponential! But that is not all. By Figure ?? a), it appears that the exponential rate of convergence of S_n is just under $1/3$; so surely




FIGURE 1. a) $\ln|S_n|$ for $n = 1, \dots, 600$. b) The first 1000 values of $(-1)^n S_n e^{n/\pi}$, coloured modulo seven.

it must be $1/\pi$. Let us cancel this exponential decay, and see what we get. Plotting $\tilde{S}_n = (-1)^n S_n e^{n/\pi}$ (cf. Figure ?? b)) makes things appear even more mysterious: now we not only have a sequence that has no reason to converge, but nevertheless does, but when the exponential decay rate is taken out, it displays a complicated and rather beautiful structure: There are seven sine curves hidden in the values of \tilde{S}_n , each of them with a period of just under 700, and a very slight overall increase in amplitude. (The factor $(-1)^n$ is there for cosmetic reasons: without it, we would see not seven but fourteen sine curves, in pairs such that for every curve the negative curve is also present. The image would not look nearly as nice.)

These images make the second question about the reason for the behaviour of S_n all the more interesting. In particular, further questions arise: why is the re-normalized sequence displaying a beautiful pattern of interlaced sine curves? What have the numbers $1/\pi$ and 7 to do with it?

In the case of (??), we will be able to answer all these questions analytically, and by doing it find a very good (but hidden) reason for S_n to converge. Before doing this, let us see whether (??) is an isolated freak phenomenon, or whether there are more of those sequences that converge without having any obvious reason to do so.

It turns out that there are many, and we have given some of those that we found in Table ???. Let us call them oscillatory sums, for their formal similarity with oscillatory integrals. Not all oscillatory sums converge, but all seem to be *much* smaller than their maximal element, or even the difference between the maximal element and the next smaller one. Also, some oscillatory sums show the interlaced sine curves when re-normalized, while some do not, see Figure ??. There are many oscillatory sums where no analytic trick like the one that we will use to treat (??) seems to work, but which nonetheless behave in a very regular way. At present, we have no idea how general this behaviour is. Have we maybe unconsciously picked sequences that are similar enough to (??) in order to behave similarly? Also, we do not know whether there is deeper reason for all of them to behave as they do, or whether one has to study them case by case. It is essentially only in the case of (??) that we have a good idea of what is going on, which we will now present.

Let us view S_n not as a sequence, but as the integer values $f(n)$ of some function f . We define

$$f(t) = \sum_{k=0}^{\lfloor t \rfloor} \frac{(k-t)^k}{k!}, \quad (t > 0)$$

where $\lfloor t \rfloor$ denotes the integer part of $t \in \mathbb{R}$. For $t < 0$ we set $f(t) = 0$. This function has some interesting properties: first of all, $f(n) = S_n$ for all $n \in \mathbb{N}$. Next, f is piecewise

S_n	max elem ($n = 600$)	Δ max elem ($n = 600$)	$ S_{600} $	exponential rate γ	graph
$\sum_{k=0}^n \frac{(k^2 - n^2)^k}{(2k)!}$	1.8×10^{235}	9.6×10^{232}	1.0×10^{33}	0.1266	Fig. ?? a)
$\sum_{k=0}^n \frac{(k^2 - n^2)^k}{(3k)!}$	3.5×10^{29}	5.0×10^{26}	6.5×10^{14}	(0.0569)	Fig. ?? b)
$\sum_{k=0}^n \frac{(\sqrt{k} - \sqrt{n})^k}{k!}$	5.1×10^7	9.1×10^5	1.9×10^{-3}	0	Fig. ?? c)
$\sum_{k=0}^{2n} e^{-\frac{1}{n}(k-n)^2} (-1)^k$	1	1.6×10^{-3}	6.3×10^{-262}	-1	Fig. ?? d)

TABLE 1. Four oscillatory sums: max elem denotes the element of maximal absolute value, Δ max elem the difference between the absolute values of the latter and the next smaller one, taken at $n = 600$. The exponential rate of growth or decay is estimated from the numerics; the value in brackets indicates that in this case, the growth is not actually exponential.




FIGURE 2. Plots of the renormalized oscillatory sums from Table ???. We plotted $s_n e^{-\gamma n}$ for $n = 1, \dots, 600$, where γ is the exponential rate given in the next to last column of Table ??.

polynomial, being a polynomial of degree n in the interval $[n, n + 1]$. f is differentiable on $(1, \infty)$, more precisely $f \in C^n(n, \infty)$ for any $n \in \mathbb{N}$. To see this, note that when t increases and crosses an integer value $n \in \mathbb{N}$, the sum in the definition of f gains the additional term $(n - t)^n/n!$. This term vanishes at $t = n$ together with its first $(n - 1)$ derivatives.

But most importantly, f is the unique solution of the equation ¹⁾

$$f'(t) = -f(t-1) \quad (t \notin \{0, 1\}) \quad (2)$$

subject to the initial condition

$$f(t) = 1 \quad \text{for } t \in [0, 1].$$

Indeed, if $t \notin \mathbb{N}$ we can differentiate the definition of f to obtain

$$f'(t) = -\sum_{k=1}^{[t]} \frac{(k-t)^{k-1}}{(k-1)!} = -\sum_{k=0}^{[t-1]} \frac{(k+1-t)^k}{k!} = -f(t-1).$$

For integer $t \geq 2$, the equation follows from the continuity of f and f' . Uniqueness follows from considering the integral form of the equation:

$$f(t) = 1 - \int_0^{t-1} f(s) ds$$

for $t \geq 1$.

So f satisfies a linear delay-differential equation with constant coefficients, which can be solved using the Laplace transform. Let

$$u(p) = \int_0^\infty e^{-pt} f(t) dt$$

be the Laplace transform of f . In order to derive the equation for u we multiply equation (??) by e^{-pt} and integrate from 1 to ∞ . Integrating the left hand side by parts we obtain

$$\begin{aligned} \int_1^\infty e^{-pt} f'(t) dt &= -e^{-p} f(1) + p \int_1^\infty e^{-pt} f(t) dt \\ &= -e^{-p} - p \int_0^1 e^{-pt} dt + pu(p) = -1 + pu(p). \end{aligned}$$

Integrating the right hand side we get

$$\int_1^\infty e^{-pt} f(t-1) dt = \int_0^\infty e^{-p(t+1)} f(t) dt = e^{-p} u(p).$$

Therefore

$$-1 + pu(p) = -e^{-p} u(p)$$

and consequently

$$u(p) = \frac{1}{p + e^{-p}}.$$

To get back to the function f , we need to invert the Laplace transform. To do so, let us first study the analytic continuation $u(z)$ of u . It has simple poles where the denominator is zero, i.e. where

$$z + e^{-z} = 0, \quad (3)$$

and it is analytic otherwise. Since (??) is equivalent to $z e^z = -1$, the poles closest to the real axis are $W(-1)$ and $\overline{W}(-1)$, where W is the Lambert function, i.e. the inverse function of $z e^z$. This function is multi-valued, and in order to get error estimates, we want the location of the next pair of poles, too.

¹⁾The authors thank D.Turaev for pointing out that $f(t)$ satisfies equation (??).




FIGURE 3. Guess what sequence you are seeing in each of these pictures!

We transform (??) into the pair of real equations

$$-x = e^{-x} \cos y \tag{4}$$

$$y = e^{-x} \sin y, \tag{5}$$

with $z = x + iy$. Since singularities come in complex conjugate pairs, we restrict to $y > 0$. (??) implies that $e^x = \sin(y)/y \leq 1$, so $x < 0$, which means that all the singularities are on the left of the real axis. Squaring and adding (??) and (??), we get $y^2 = e^{-2x} - x^2$, so for any solution $|y|$ grows monotonically with $|x|$, and vice versa. Finally, rearranging (??) and taking the logarithm yields $x = \ln(\sin(y)/y)$ whenever $\sin(y) > 0$, with no real solution otherwise. Inserted into (??), this leads to

$$\ln\left(\frac{\sin y}{y}\right) \sin y = -y \cos y. \tag{6}$$

The left hand side of (??) is defined for $(k\pi, (k+1)\pi)$ with k even, and has exactly one solution in each of these intervals. The first two of them are $y_1 = 1.33724$ and $y_2 = 7.58863$, leading to $z_1 = W(-1) \approx -0.31832 + 1.33724i$ and $z_2 \approx -2.06228 + 7.58863i$. All further solutions have even larger negative real part.

The inverse Laplace transform can now be done using the Bromwich integral:

$$f(t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \frac{e^{pt}}{p + e^{-p}} dp,$$

where as a path of integration we choose the imaginary axis, which is right of all singularities as is required. Now, shifting the path of integration to the left and using residues, we get

$$f(t) = 2\text{Re} \frac{e^{z_1 t}}{1 + z_1} + O(e^{-\text{Re}(z_2)t}).$$

With $z_j = x_j + iy_j$ this gives

$$f(t) = \frac{2}{|1 + z_1|} e^{x_1 t} \cos(y_1 t + \arg(1 + z_1)) + O(e^{x_2 t}). \tag{7}$$

So, f is a decaying exponential times a cosine function, and we are in the position to answer all of the questions that we asked before. Firstly, what does $1/\pi$ have to do with S_n ? The answer is, nothing at all, except that it happens to agree with the $-\text{Re}(W(-1))$ in the first four valid digits. The discrepancy is small enough so we could not pick it up numerically. Secondly, what has seven to do with it, and why do we see the interlaced sine functions? Given (??), the answer becomes obvious, although it may have been obvious from the start for people who are working in signal processing. Have a look at Figure ?? a) and b) and guess what you are seeing.

The surprising answer is that both of the plots actually depict the same sequence, namely $(\cos(n))_{n \in \mathbb{N}}$. The only difference is that the first plot contains the first 1000 elements, while the second one contains the first 3000 ones. Apart from telling us that we might want to be careful when drawing conclusions from looking at plots, Figure ?? b) could have suggested from the start what we have seen in (??), namely that S_n is an undersampling of a trigonometric function at a frequency that is incommensurable to the period of that function. And now the significance of the number 7 for S_n is not hard to understand any more: If we consider the defect of diophantine approximations

$$d(n) = \min_{1 \leq p, q \leq n} \left| \frac{\operatorname{Im}(z_1)}{2\pi} - \frac{p}{q} \right|,$$

then the numbers n where d jumps correspond to sampling frequencies which will give the illusion of periodic behavior. If N points are plotted then we need that $Nd(n) = O(1)$. The first solutions for $1000d(n) \leq 3$ is obtained for $n = 14$.

VOLKER BETZ
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF WARWICK
 COVENTRY, CV4 7AL, ENGLAND
<http://www.maths.warwick.ac.uk/~betz/>
E-mail address: `v.m.betz@warwick.ac.uk`