

Decay in norm of transfer operators for semiflows

by

IAN MELBOURNE (Warwick), NICOLÒ PAVIATO (Warwick) and
DALIA TERHESIU (Leiden)

Abstract. We establish exponential decay in Hölder norm of transfer operators applied to smooth observables of uniformly and nonuniformly expanding semiflows with exponential decay of correlations.

1. Introduction. Exponential decay of correlations is well-understood for large classes of uniformly and nonuniformly expanding maps; see for example [8, 13, 15, 16, 18, 23, 24, 25, 26, 29]. The typical method of proof is to establish a spectral gap for the associated transfer operator L . Such a spectral gap yields a decay rate $\|L^n v - \int v\| \leq C_v e^{-an}$ for v lying in a suitable function space, where a, C_v are positive constants. Decay of correlations is an immediate consequence of such decay for L^n .

Results on decay of correlations lead to numerous statistical limit theorems. Although not needed for results such as the central limit theorem, strong norm control on $L^n v$ is often useful for finer statistical properties. For example, rates of convergence in the central limit theorem [14] and the associated functional central limit theorem [4] rely heavily on control of operator norms.

In this paper, we consider norm decay of transfer operators for uniformly and nonuniformly expanding semiflows. Here, the standard method is to deduce decay of the correlation function from analyticity of Laplace transforms, bypassing spectral properties of L_t ; see [11, 17, 22]. As far as we know, the only result on spectral gaps for transfer operators of semiflows is due to Tsujii [27]. However, this result is for suspension semiflows over the doubling map with a C^3 roof function, where the smoothness of the roof function

2020 *Mathematics Subject Classification*: Primary 37A25; Secondary 37C10, 37C30, 60K05.

Key words and phrases: operator renewal theory, exponential mixing, nonuniformly expanding semiflows.

Received 10 April 2021; revised 22 February 2022.

Published online *.

is crucial and very restrictive. A similar result for contact Anosov flows is proved in [28]. Both of the papers [27, 28] obtain spectral gaps for L_t acting on a suitable anisotropic Banach space. In addition, a paper of Butterley [10] obtains norm decay of transfer operators, but the results are quite different from ours and the abstract setting there seems not to cover the situation considered here (see Remark 2.6 below). Apart from these, there are apparently no previous results on norm decay of transfer operators for semiflows or flows.

Recently, in [20], we showed that spectral gaps are impossible in Hölder spaces with exponent greater than $1/2$ (and in any Banach space that embeds in such a Hölder space). Nevertheless, our aim of controlling the Hölder norm of L_tv for a large class of semiflows and observables v remains viable, and our main result is the first in this direction. We consider uniformly and nonuniformly expanding semiflows satisfying a Dolgopyat-type estimate [11]. Such an estimate plays a key role in proving exponential decay of correlations for the semiflow. Theorem 2.3 below shows how to use this estimate to prove exponential decay of L_tv in a Hölder norm for smooth mean zero observables satisfying a good support condition. Apart from the Dolgopyat estimate, the main ingredient is an operator renewal equation for semiflows [21] which enables consideration of the operator Laplace transform $\int_0^\infty e^{-st} L_t dt$.

The remainder of the paper is organised as follows. In Section 2, we recall the setup for nonuniformly expanding semiflows with exponential decay of correlations and state our main result, Theorem 2.3, on decay in norm. In Section 3, we prove Theorem 2.3.

NOTATION. We use the “big O” and \ll notation interchangeably, writing $a_n = O(b_n)$ or $a_n \ll b_n$ if there are constants $C > 0$, $n_0 \geq 1$ such that $a_n \leq Cb_n$ for all $n \geq n_0$.

2. Setup and statement of the main result. In this section, we state our result on Hölder norm decay of transfer operators for uniformly and nonuniformly expanding semiflows.

Let (Y, d) be a bounded metric space with Borel probability measure μ and an at most countable measurable partition $\{Y_j\}$. Let $F : Y \rightarrow Y$ be a measure-preserving transformation such that F restricts to a measure-theoretic bijection from Y_j onto Y for each j . Let $g = d\mu/(d\mu \circ F)$ be the inverse Jacobian of F .

Fix $\eta \in (0, 1)$. Assume that there are constants $\lambda > 1$ and $C > 0$ such that $d(Fy, Fy') \geq \lambda d(y, y')$ and $|\log g(y) - \log g(y')| \leq Cd(Fy, Fy')^\eta$ for all $y, y' \in Y_j$, $j \geq 1$. In particular, F is a Gibbs–Markov map as in [2] (see also [1, 3]) with ergodic (and mixing) invariant measure μ .

Let $\varphi : Y \rightarrow [2, \infty)$ be a piecewise continuous roof function. We assume that there is a constant $C > 0$ such that

$$(2.1) \quad |\varphi(y) - \varphi(y')| \leq Cd(Fy, Fy')^\eta$$

for all $y, y' \in Y_j$, $j \geq 1$. Also, we assume exponential tails, that is, there exists $\delta_0 > 0$ such that

$$(2.2) \quad \sum_j \mu(Y_j) e^{\delta_0 |1_{Y_j} \varphi|_\infty} < \infty.$$

Define the suspension $Y^\varphi = \{(y, u) \in Y \times [0, \infty) : u \in [0, \varphi(y)]\} / \sim$ where $(y, \varphi(y)) \sim (Fy, 0)$. The suspension semiflow $F_t : Y^\varphi \rightarrow Y^\varphi$ is given by $F_t(y, u) = (y, u+t)$ modulo identifications. We define the ergodic F_t -invariant probability measure $\mu^\varphi = (\mu \times \text{Lebesgue}) / \bar{\varphi}$ where $\bar{\varphi} = \int_Y \varphi d\mu$ ⁽¹⁾.

Let $L_t : L^1(Y^\varphi) \rightarrow L^1(Y^\varphi)$ denote the transfer operator corresponding to F_t (so $\int_{Y^\varphi} L_t v w d\mu^\varphi = \int_{Y^\varphi} v w \circ F_t d\mu^\varphi$ for all $v \in L^1(Y^\varphi)$, $w \in L^\infty(Y^\varphi)$, $t > 0$) and let $R_0 : L^1(Y) \rightarrow L^1(Y)$ denote the transfer operator for F . Recall (see for example [2]) that $(R_0 v)(y) = \sum_j g(y_j) v(y_j)$ where y_j is the unique preimage of y under $F|_{Y_j}$, and there is a constant $C > 0$ such that

$$(2.3) \quad |g(y)| \leq C\mu(Y_j), \quad |g(y) - g(y')| \leq C\mu(Y_j)d(Fy, Fy')^\eta,$$

for all $y, y' \in Y_j$, $j \geq 1$.

Function space on Y^φ . Let $Y_j^\varphi = \{(y, u) \in Y^\varphi : y \in Y_j\}$. Fix $\eta \in (0, 1]$, $\delta > 0$. For $v : Y^\varphi \rightarrow \mathbb{R}$, define $|v|_{\delta, \infty} = \sup_{(y, u) \in Y^\varphi} e^{-\delta u} |v(y, u)|$ and

$$\begin{aligned} \|v\|_{\delta, \eta} &= |v|_{\delta, \infty} + |v|_{\delta, \eta}, \\ |v|_{\delta, \eta} &= \sup_{j \geq 1} \sup_{(y, u), (y', u) \in Y_j^\varphi, y \neq y'} e^{-\delta u} \frac{|v(y, u) - v(y', u)|}{d(y, y')^\eta}. \end{aligned}$$

Then $\mathcal{F}_{\delta, \eta}(Y^\varphi)$ consists of observables $v : Y^\varphi \rightarrow \mathbb{R}$ with $\|v\|_{\delta, \eta} < \infty$.

Next, define $\partial_u v$ to be the partial derivative of v with respect to u at points $(y, u) \in Y^\varphi$ with $u \in (0, \varphi(y))$ and to be the appropriate one-sided partial derivative when $u \in \{0, \varphi(y)\}$. For $m \geq 0$, define $\mathcal{F}_{\delta, \eta, m}(Y^\varphi)$ to consist of observables $v : Y^\varphi \rightarrow \mathbb{R}$ such that $\partial_u^j v \in \mathcal{F}_{\delta, \eta}(Y^\varphi)$ for $j = 0, 1, \dots, m$, with norm $\|v\|_{\delta, \eta, m} = \max_{j=0, \dots, m} \|\partial_u^j v\|_{\delta, \eta}$.

DEFINITION 2.1. Given $r > 0$, we consider the subset $\{(y, u) \in Y \times \mathbb{R} : u \in [r, \varphi(y) - r]\}$ viewed as a subset of Y^φ . We say that a function $v : Y^\varphi \rightarrow \mathbb{R}$ has *good support* if there exists $r > 0$ such that $\text{supp } v \subset \{(y, u) \in Y \times \mathbb{R} : u \in [r, \varphi(y) - r]\}$.

For functions with good support, $\partial_u v$ coincides with the derivative $\partial_t v = \lim_{h \rightarrow 0} (v \circ F_h - v)/h$ in the flow direction.

⁽¹⁾ We call such semiflows “nonuniformly expanding” since they are the continuous-time analogue of maps that are nonuniformly expanding in the sense of Young [29]. “Uniformly expanding” semiflows are those with φ bounded; they have bounded distortion as well as uniform expansion.

REMARK 2.2. It is standard to restrict to observables with good support when considering decay of correlations for semiflows, see for instance [12, 27].

Let

$$\mathcal{F}_{\delta,\eta,m}^0(Y^\varphi) = \left\{ v \in \mathcal{F}_{\delta,\eta,m}(Y^\varphi) : \int_{Y^\varphi} v d\mu^\varphi = 0 \right\}.$$

We write $\mathcal{F}_{\delta,\eta}(Y^\varphi)$ and $\mathcal{F}_{\delta,\eta}^0(Y^\varphi)$ when $m = 0$.

Function space on Y . For $v : Y \rightarrow \mathbb{R}$, define

$$\|v\|_\eta = |v|_\infty + |v|_\eta, \quad |v|_\eta = \sup_{j \geq 1} \sup_{y, y' \in Y_j, y \neq y'} |v(y) - v(y')|/d(y, y')^\eta.$$

Let $\mathcal{F}_\eta(Y)$ consist of observables $v : Y \rightarrow \mathbb{R}$ with $\|v\|_\eta < \infty$.

Dolgopyat estimate. Define the twisted transfer operators

$$\widehat{R}_0(s) : L^1(Y) \rightarrow L^1(Y), \quad \widehat{R}_0(s)v = R_0(e^{-s\varphi}v).$$

We assume that there exist $\gamma \in (0, 1)$, $\epsilon > 0$, $m_0 \geq 0$, $A, D > 0$ such that

$$(2.4) \quad \|\widehat{R}_0(s)^n\|_{\mathcal{F}_\eta(Y) \rightarrow \mathcal{F}_\eta(Y)} \leq |b|^{m_0} \gamma^n$$

for all $s = a + ib \in \mathbb{C}$ with $|a| < \epsilon$, $|b| \geq D$ and all $n \geq A \log |b|$. Such an assumption holds in the settings of [5, 6, 7, 11].

Now we can state our main result on norm decay for L_t .

THEOREM 2.3. *Under these assumptions, there exist $\epsilon > 0$, $m \geq 1$, $C > 0$ such that*

$$\|L_t v\|_{\delta,\eta,1} \leq C e^{-\epsilon t} \|v\|_{\delta,\eta,m} \quad \text{for all } t > 0$$

for all $v \in \mathcal{F}_{\delta,\eta,m}^0(Y^\varphi)$ with good support.

REMARK 2.4. Since the norm applied to v is stronger than the norm applied to $L_t v$, Theorem 2.3 does not imply a spectral gap for L_t . We note that the norm on $\mathcal{F}_{\delta,\eta,1}(Y^\varphi)$ gives no Hölder control in the flow direction when passing through points of the form $(y, \varphi(y))$. This lack of control is a barrier to mollification arguments of the type usually used to pass from smooth observables to Hölder observables. In fact, such arguments are doomed to fail at the operator level by [20, Theorem 1.1] when $\eta > 1/2$ and hence seem unlikely for any η .

REMARK 2.5. Usually, we can take $m_0 \in (0, 1)$ in (2.4), in which case $m = 3$ suffices in Theorem 2.3.

There are numerous simplifications when $\{Y_j\}$ is a finite partition. In particular, conditions (2.1) and (2.2) are redundant and we can take $\delta = 0$.

REMARK 2.6. At first glance, Theorem 2.3 has some similarities with [10, Theorem 1]. In particular, we mention formula (2.4) therein which takes the form $\|P_t \mu\|_{\mathcal{A}} \leq C_\ell e^{-\ell t} \|Z \mu\|_{\mathcal{B}}$ where $Z = \partial_t$. However, $\|\cdot\|_{\mathcal{A}}$ corresponds to a “weak” norm which would just be the L^∞ norm in our setting. Moreover,

the hypothesis in [10] that the operators $T_t : \mathcal{B} \rightarrow \mathcal{B}$ ($L_t : \mathcal{F}_{\delta,\eta,1}(Y^\varphi) \rightarrow \mathcal{F}_{\delta,\eta,1}(Y^\varphi)$ in our notation) are bounded looks to be unverifiable in our setting even for fixed t . On the other hand, the expansion in equation (2.3) of [10] is beyond our methods.

REMARK 2.7. Numerous (non)uniformly hyperbolic flows are modelled (after inducing and quotienting along stable leaves) by “Gibbs–Markov semiflows” $F_t : Y^\varphi \rightarrow Y^\varphi$ of the type considered here with the exponential tail condition (2.2). These include basic sets for Axiom A flows, Lorentz gases with finite horizon, and Lorenz attractors (see for instance [19, Section 1.1]). Whenever the Dolgopyat estimate (2.4) is satisfied in such examples, as in [5, 6, 7, 11], Theorem 2.3 guarantees exponential decay for the norm of the transfer operator for the corresponding Gibbs–Markov semiflow.

3. Proof of Theorem 2.3. Our proof of norm decay is broken into three parts. In Subsection 3.1, we recall a continuous-time operator renewal equation [21] which enables estimates of Laplace transforms of transfer operators at the level of Y . In Subsection 3.2, we show how to pass to estimates of Laplace transforms of L_t . In Subsection 3.3, we invert the Laplace transform to obtain norm decay of L_t .

3.1. Operator renewal equation. Let $\tilde{Y} = Y \times [0, 1]$ and define

$$\tilde{F} : \tilde{Y} \rightarrow \tilde{Y}, \quad \tilde{F}(y, u) = (Fy, u),$$

with transfer operator $\tilde{R} : L^1(\tilde{Y}) \rightarrow L^1(\tilde{Y})$. Also, define

$$\tilde{\varphi} : \tilde{Y} \rightarrow [2, \infty), \quad \tilde{\varphi}(y, u) = \varphi(y).$$

Define the twisted transfer operators

$$\hat{R}(s) : L^1(\tilde{Y}) \rightarrow L^1(\tilde{Y}), \quad \hat{R}(s)v = \tilde{R}(e^{-s\tilde{\varphi}}v).$$

Let $\tilde{Y}_j = Y_j \times [0, 1]$. For $v : \tilde{Y} \rightarrow \mathbb{R}$, define

$$\|v\|_\eta = |v|_\infty + |v|_\eta, \quad |v|_\eta = \sup_{j \geq 1} \sup_{(y,u), (y',u) \in \tilde{Y}_j, y \neq y'} |v(y, u) - v(y', u)| / d(y, y')^\eta.$$

Let $\mathcal{F}_\eta(\tilde{Y})$ consist of observables $v : \tilde{Y} \rightarrow \mathbb{R}$ with $\|v\|_\eta < \infty$. Let

$$\mathcal{F}_\eta^0(\tilde{Y}) = \left\{ v \in \mathcal{F}_\eta(\tilde{Y}) : \int_{\tilde{Y}} v d\tilde{\mu} = 0 \right\},$$

where $\tilde{\mu} = \mu \times \text{Leb}_{[0,1]}$.

LEMMA 3.1. *Write $s = a + ib \in \mathbb{C}$. There exist $\epsilon > 0$, $m_1 \geq 0$, $C > 0$ such that*

(a) $s \mapsto (I - \hat{R}(s))^{-1} : \mathcal{F}_\eta^0(\tilde{Y}) \rightarrow \mathcal{F}_\eta(\tilde{Y})$ is analytic on $\{|a| < \epsilon\}$;

- (b) $s \mapsto (I - \widehat{R}(s))^{-1} : \mathcal{F}_\eta(\widetilde{Y}) \rightarrow \mathcal{F}_\eta(\widetilde{Y})$ is analytic on $\{|a| < \epsilon\}$ except for a simple pole at $s = 0$;
- (c) $\|(I - \widehat{R}(s))^{-1}\|_{\mathcal{F}_\eta(\widetilde{Y}) \rightarrow \mathcal{F}_\eta(\widetilde{Y})} \leq C|b|^{m_1}$ for $|a| \leq \epsilon$, $|b| \geq 1$.

Proof. It suffices to verify these properties for $Z(s) = (I - \widehat{R}_0(s))^{-1}$ on Y . They immediately transfer to $(I - \widehat{R}(s))^{-1}$ on \widetilde{Y} since $(\widehat{R}v)(y, u) = (\widehat{R}_0v^u)(y)$ where $v^u(y) = v(y, u)$.

The arguments for passing from (2.4) to the desired properties for $Z(s)$ are standard. For completeness, we sketch these details now, recalling arguments from [5]. Define $\mathcal{F}_\eta(Y)$ with norm $\|\cdot\|_\eta$ by restricting to $u = 0$ (this coincides with the usual Hölder space on Y). Let A, D, ϵ and m_0 be as in (2.4). Increase A and D so that $D > 1$ and $|b|^{m_0} \gamma^{[A \log |b|]} \leq 1/2$ for $|b| \geq D$. Suppose that $|a| \leq \epsilon$, $|b| \geq D$. Then $\|\widehat{R}_0(s)^{[A \log |b|]}\|_\eta \leq |b|^{m_0} \gamma^{[A \log |b|]} \leq 1/2$ and $\|(I - \widehat{R}_0(s)^{[A \log |b|]})^{-1}\|_\eta \leq 2$.

As in [5, Proposition 2.5], we can shrink ϵ so that $s \rightarrow \widehat{R}_0(s)$ is continuous on $\mathcal{F}_\eta(Y)$ for $|a| \leq \epsilon$. The simple eigenvalue 1 for $\widehat{R}_0(0) = R_0$ extends to a continuous family of simple eigenvalues $\lambda(s)$ for $|s| \leq \epsilon$. Hence we can choose ϵ so that $1/2 < \lambda(a) < 2$ for $|a| \leq \epsilon$. By [5, Corollary 2.8], $\|\widehat{R}_0(s)^n\|_\eta \ll |b|\lambda(a)^n \leq |b|2^n$ for all $n \geq 1$, $|a| \leq \epsilon$, $|b| \geq D$. Hence

$$\begin{aligned} \|Z(s)\|_\eta &\leq (1 + \|\widehat{R}_0(s)\|_\eta + \dots + \|\widehat{R}_0(s)^{[A \log |b|]-1}\|_\eta) \|(I - \widehat{R}_0(s)^{[A \log |b|]})^{-1}\|_\eta \\ &\ll (\log |b|) |b| 2^{A \log |b|} \leq |b|^{m_1} \end{aligned}$$

with $m_1 = 1 + A \log 2$. This proves analyticity on the region $\{|a| < \epsilon, |b| > D\}$ with the desired estimates for property (c) on this region.

For $|a| \leq \epsilon$, $|b| \leq D$, we recall arguments from [5, proof of Lemma 2.22] (where $\widehat{R}_0(s)$ is denoted Q_s). For ϵ sufficiently small, the part of the spectrum of $\widehat{R}_0(s)$ that is close to 1 consists only of isolated eigenvalues. Also, the spectral radius of $\widehat{R}_0(s)$ is at most $\lambda(a)$ and $\lambda(a) < 1$ for $a \in [0, \epsilon]$, so $s \mapsto Z(s)$ is analytic on $\{0 < a < \epsilon\}$.

Suppose that $\widehat{R}_0(ib)v = v$ for some $v \in \mathcal{F}_\eta(Y)$, $b \neq 0$. Choose $q \geq 1$ such that $q|b| > D$. Since $\widehat{R}_0(s)$ is the L^2 adjoint of $v \mapsto e^{s\varphi}v \circ F$, we have $e^{ib\varphi}v \circ F = v$. Hence $e^{iqb\varphi}v^q \circ F = v^q$ and so $\widehat{R}_0(iqb)v^q = v^q$. But $\|Z(iqb)v^q\|_\eta < \infty$, so $v = 0$. Hence $1 \notin \text{spec } \widehat{R}_0(ib)$ for all $b \neq 0$. It follows that for all $b \neq 0$ there exists an open set $U_b \subset \mathbb{C}$ containing ib such that $1 \notin \text{spec } \widehat{R}_0(s)$ for all $s \in U_b$, and so $s \mapsto Z(s)$ is analytic on U_b .

Next, we recall that for s near to zero, $\lambda(s) = 1 + cs + O(s^2)$ where $c < 0$. Hence $s \mapsto Z(s)$ has a simple pole at zero. It follows that there exists $\epsilon > 0$ such that $s \mapsto Z(s)$ is analytic on $\{|a| < \epsilon, |b| < 2D\}$ except for a simple pole at $s = 0$. Combining this with the estimates on $\{|a| < \epsilon, |b| \geq D\}$ we have proved properties (b) and (c) for $Z(s)$.

Finally, the spectral projection π corresponding to the eigenvalue $\lambda(0)=1$ for $\widehat{R}_0(0) = R$ is given by $\pi v = \int_Y v d\mu$. Hence the pole disappears on restriction to observables of mean zero, proving property (a) for $Z(s)$. ■

Next define

$$T_t v = 1_{\widetilde{Y}} L_t(1_{\widetilde{Y}} v), \quad U_t v = 1_{\widetilde{Y}} L_t(1_{\{\widetilde{\varphi} > t\}} v)$$

and

$$\widehat{T}(s) = \int_0^\infty e^{-st} T_t dt, \quad \widehat{U}(s) = \int_0^\infty e^{-st} U_t dt.$$

By [21, Theorem 3.3], we have the operator renewal equation

$$\widehat{T} = \widehat{U}(I - \widehat{R})^{-1}.$$

PROPOSITION 3.2. *There exist $\epsilon > 0$, $C > 0$ such that $s \mapsto \widehat{U}(s) : \mathcal{F}_\eta(\widetilde{Y}) \rightarrow \mathcal{F}_\eta(\widetilde{Y})$ is analytic on $\{|a| < \epsilon\}$ and $\|\widehat{U}(s)\|_{\mathcal{F}_\eta(\widetilde{Y}) \rightarrow \mathcal{F}_\eta(\widetilde{Y})} \leq C|s|$ for $|a| \leq \epsilon$.*

Proof. By [21, Proposition 3.4],

$$(U_t v)(y, u) = \begin{cases} v(y, u - t) 1_{[t, 1]}(u), & 0 \leq t \leq 1, \\ (\widetilde{R}v_t)(y, u), & t > 1, \end{cases}$$

where $v_t(y, u) = 1_{\{t < \varphi(y) < t+1-u\}} v(y, u - t + \varphi(y))$. Consequently, $\widehat{U}(s) = \widehat{U}_1(s) + \widehat{U}_2(s)$, where

$$(\widehat{U}_1(s)v)(y, u) = \int_0^u e^{-st} v(y, u - t) dt, \quad \widehat{U}_2(s)v = \int_1^\infty e^{-st} \widetilde{R}v_t dt.$$

It is clear that $\|\widehat{U}_1(s)v\|_\eta \leq e^\epsilon \|v\|_\eta$. We focus attention on the second term

$$(\widehat{U}_2(s)v)(y, u) = \sum_j g(y_j) \int_1^\infty e^{-st} v_t(y_j, u) dt = \sum_j g(y_j) \widehat{V}(s)(y_j, u),$$

where $\widehat{V}(s)(y, u) = \int_u^1 e^{s(t-u-\varphi)} v(y, t) dt$. Clearly, $|1_{Y_j} \widehat{V}(s)|_\infty \leq e^{\epsilon |1_{Y_j} \varphi|_\infty} |v|_\infty$. Also,

$$\widehat{V}(s)(y, u) - \widehat{V}(s)(y', u) = I + J,$$

where

$$I = \int_u^1 (e^{s(t-u-\varphi(y))} - e^{s(t-u-\varphi(y'))}) v(y, t) dt,$$

$$J = \int_u^1 e^{s(t-u-\varphi(y'))} (v(y, t) - v(y', t)) dt.$$

For $y, y' \in Y_j$,

$$|I| \leq |v|_\infty \int_u^1 e^{\epsilon(|1_{Y_j}\varphi|_\infty + u - t)} |s| |\varphi(y) - \varphi(y')| dt \ll |s| |v|_\infty e^{\epsilon|1_{Y_j}\varphi|_\infty} d(Fy, Fy')^\eta$$

by (2.1), and

$$|J| \leq \int_u^1 e^{\epsilon(|1_{Y_j}\varphi|_\infty + u - t)} |v(y, t) - v(y', t)| dt \leq e^{\epsilon|1_{Y_j}\varphi|_\infty} |v|_\eta d(y, y')^\eta.$$

Hence $|\widehat{V}(s)(y, u) - \widehat{V}(s)(y', u)|_\eta \ll |s| e^{\epsilon|1_{Y_j}\varphi|_\infty} \|v\|_\eta d(Fy, Fy')^\eta$.

It follows from the estimates for $1_{Y_j}\widehat{V}(s)$ together with (2.3) that

$$\|\widehat{U}_2(s)v\|_\eta \ll \sum_j |s| \mu(Y_j) e^{\epsilon|1_{Y_j}\varphi|_\infty} \|v\|_\eta.$$

By (2.2), $\|\widehat{U}_2(s)v\|_\eta \ll |s| \|v\|_\eta$ for ϵ sufficiently small. We conclude that $\|\widehat{U}(s)v\|_\eta \ll |s| \|v\|_\eta$. ■

3.2. From \widehat{T} on \widetilde{Y} to \widehat{L} on Y^φ . Lemma 3.1 and Proposition 3.2 yield analyticity and estimates for $\widehat{T} = \widehat{U}(I - \widehat{R})^{-1}$ on \widetilde{Y} . In this subsection, we show how these properties are inherited by $\widehat{L}(s) = \int_0^\infty e^{-st} L_t dt$ on Y^φ . Recall that $\widetilde{Y} = Y \times [0, 1]$, which we view as a subset of Y^φ .

REMARK 3.3. The approach in this subsection is similar to that in [9, Section 5] but there are some important differences. The rationale behind the two-step decomposition in Propositions 3.4 and 3.5 below is that the discreteness of the decomposition in Proposition 3.4 simplifies many formulas significantly. In particular, the previously problematic term E_t in [9] becomes elementary (and vanishes for large t when φ is bounded). The decomposition in Proposition 3.5 remains continuous to simplify the estimates in Proposition 3.8.

Since the setting in [9] is different (infinite ergodic theory, reinducing), we keep the exposition here self-contained even where the estimates coincide with those in [9].

Define

$$A_n : L^1(\widetilde{Y}) \rightarrow L^1(Y^\varphi), \quad (A_n v)(y, u) = 1_{\{n \leq u < n+1\}} (L_n v)(y, u), \quad n \geq 0,$$

$$E_t : L^1(Y^\varphi) \rightarrow L^1(Y^\varphi), \quad (E_t v)(y, u) = 1_{\{[t]+1 \leq u \leq \varphi(y)\}} (L_t v)(y, u), \quad t > 0.$$

PROPOSITION 3.4. $L_t = \sum_{j=0}^{[t]} A_j 1_{\widetilde{Y}} L_{t-j} + E_t$ for $t > 0$.

Proof. For $y \in Y$, $u \in (0, \varphi(y))$,

$$\begin{aligned} (L_t v)(y, u) &= \sum_{j=0}^{[t]} 1_{\{j \leq u < j+1\}} (L_t v)(y, u) + 1_{\{[t]+1 \leq u \leq \varphi(y)\}} (L_t v)(y, u) \\ &= \sum_{j=0}^{[t]} (A_j L_{t-j} v)(y, u) + (E_t v)(y, u). \end{aligned}$$

Now use $A_n = A_n 1_{\tilde{Y}}$. ■

Next, define

$$\begin{aligned} B_t &: L^1(Y^\varphi) \rightarrow L^1(\tilde{Y}), & B_t v &= 1_{\tilde{Y}} L_t (1_{\Delta_t} v), \\ G_t &: L^1(Y^\varphi) \rightarrow L^1(\tilde{Y}), & G_t v &= B_t (\omega(t) v), \\ H_t &: L^1(Y^\varphi) \rightarrow L^1(\tilde{Y}), & H_t v &= 1_{\tilde{Y}} L_t (1_{\Delta'_t} v), \end{aligned}$$

for $t > 0$, where

$$\begin{aligned} \Delta_t &= \{(y, u) \in Y^\varphi : \varphi(y) - t \leq u < \varphi(y) - t + 1\}, \\ \Delta'_t &= \{(y, u) \in Y^\varphi : u < \varphi(y) - t\}, \\ \omega(t)(y, u) &= \varphi(y) - u - t + 1. \end{aligned}$$

PROPOSITION 3.5. $1_{\tilde{Y}} L_t = \int_0^t T_{t-\tau} B_\tau d\tau + G_t + H_t$ for $t > 0$.

Proof. Let $y \in Y$, $u \in [0, \varphi(y)]$. Then

$$\begin{aligned} \int_0^t 1_{\Delta_\tau}(y, u) d\tau &= \int_0^t 1_{\{\varphi(y) - u \leq \tau \leq \varphi(y) - u + 1\}} d\tau \\ &= 1_{\{t \geq \varphi(y) - u + 1\}} + 1_{\{\varphi(y) - u \leq t < \varphi(y) - u + 1\}} (t - \varphi(y) + u) \\ &= 1 - 1_{\{t < \varphi(y) - u + 1\}} + 1_{\{\varphi(y) - u \leq t < \varphi(y) - u + 1\}} (t - \varphi(y) + u) \\ &= 1 - 1_{\Delta'_t}(y, u) + 1_{\Delta_t}(y, u) (t - \varphi(y) + u - 1). \end{aligned}$$

Hence $\int_0^t 1_{\Delta_\tau} d\tau = 1 - 1_{\Delta_t} \omega(t) - 1_{\Delta'_t}$. It follows that

$$\begin{aligned} \int_0^t T_{t-\tau} B_\tau v d\tau &= 1_{\tilde{Y}} \int_0^t L_{t-\tau} 1_{\tilde{Y}} B_\tau v d\tau = 1_{\tilde{Y}} \int_0^t L_{t-\tau} B_\tau v d\tau \\ &= 1_{\tilde{Y}} \int_0^t L_{t-\tau} L_\tau (1_{\Delta_\tau} v) d\tau = 1_{\tilde{Y}} L_t \left(\int_0^t 1_{\Delta_\tau} v d\tau \right) \\ &= 1_{\tilde{Y}} L_t v - G_t v - H_t v, \end{aligned}$$

as required. ■

We have already defined the Laplace transforms $\widehat{L}(s)$ and $\widehat{T}(s)$ for $s = a + ib$ with $a > 0$. Similarly, define

$$\begin{aligned}\widehat{B}(s) &= \int_0^{\infty} e^{-st} B_t dt, & \widehat{E}(s) &= \int_0^{\infty} e^{-st} E_t dt, \\ \widehat{G}(s) &= \int_0^{\infty} e^{-st} G_t dt, & \widehat{H}(s) &= \int_0^{\infty} e^{-st} H_t dt.\end{aligned}$$

Also, we define the discrete transform $\widehat{A}(s) = \sum_{n=0}^{\infty} e^{-sn} A_n$.

COROLLARY 3.6. $\widehat{L}(s) = \widehat{A}(s)\widehat{T}(s)\widehat{B}(s) + \widehat{A}(s)\widehat{G}(s) + \widehat{A}(s)\widehat{H}(s) + \widehat{E}(s)$ for $a > 0$.

Proof. By Proposition 3.4,

$$\begin{aligned}\widehat{L}(s) - \widehat{E}(s) &= \int_0^{\infty} e^{-st} \sum_{j=0}^{[t]} A_j 1_{\widetilde{Y}} L_{t-j} dt = \sum_{j=0}^{\infty} e^{-sj} A_j 1_{\widetilde{Y}} \int_j^{\infty} e^{-s(t-j)} L_{t-j} dt \\ &= \widehat{A}(s) 1_{\widetilde{Y}} \int_0^{\infty} e^{-st} L_t dt = \widehat{A}(s) 1_{\widetilde{Y}} \widehat{L}(s).\end{aligned}$$

Hence $\widehat{L} = \widehat{A} 1_{\widetilde{Y}} \widehat{L} + \widehat{E}$. In addition, by Proposition 3.5, $1_{\widetilde{Y}} \widehat{L} = \widehat{T} \widehat{B} + \widehat{G} + \widehat{H}$. ■

PROPOSITION 3.7. *Let $\delta > \epsilon > 0$. Then there is a constant $C > 0$ such that*

- (a) $\|\widehat{A}(s)\|_{\mathcal{F}_{\eta}(\widetilde{Y}) \rightarrow \mathcal{F}_{\delta, \eta}(Y^{\varphi})} \leq 1$,
- (b) $\|\widehat{E}(s)\|_{\mathcal{F}_{\delta, \eta}(Y^{\varphi}) \rightarrow \mathcal{F}_{\delta, \eta}(Y^{\varphi})} \leq C$,
- (c) $\|\widehat{H}(s)\|_{\mathcal{F}_{\delta, \eta}(Y^{\varphi}) \rightarrow \mathcal{F}_{\eta}(\widetilde{Y})} \leq e^{\delta}$,

for $|a| \leq \epsilon$.

Proof. (a) Let $v \in \mathcal{F}_{\eta}(\widetilde{Y})$. Let $(y, u), (y', u) \in Y_j^{\varphi}$, $j \geq 1$. Since $(A_n v)(y, u) = 1_{\{n \leq u < n+1\}} v(y, u - n)$, we have

$$(\widehat{A}(s)v)(y, u) = \sum_{n=0}^{\infty} e^{-sn} 1_{\{n \leq u < n+1\}} v(y, u - n) = e^{-s[u]} v(y, u - [u]).$$

Hence

$$\begin{aligned}|(\widehat{A}(s)v)(y, u)| &\leq e^{\epsilon u} |v|_{\infty}, \\ |(\widehat{A}(s)v)(y, u) - (\widehat{A}(s)v)(y', u)| &\leq e^{\epsilon u} |v|_{\eta} d(y, y')^{\eta}.\end{aligned}$$

That is, $|\widehat{A}(s)v|_{\epsilon, \infty} \leq |v|_{\infty}$, $|\widehat{A}(s)v|_{\epsilon, \eta} \leq |v|_{\eta}$. So $\|\widehat{A}(s)v\|_{\delta, \eta} \leq \|\widehat{A}(s)v\|_{\epsilon, \eta} \leq \|v\|_{\eta}$.

(b) We take $C = 1/(\delta - \epsilon)$. Let $v \in \mathcal{F}_{\delta,\eta}(Y^\varphi)$. Let $(y, u), (y', u) \in Y_j^\varphi$, $j \geq 1$. Note that $(E_t v)(y, u) = 1_{\{[t]+1 \leq u\}} v(y, u - t)$, so

$$(\widehat{E}(s)v)(y, u) = \int_0^\infty e^{-st} 1_{\{[t]+1 \leq u\}} v(y, u - t) dt.$$

Hence

$$|(\widehat{E}(s)v)(y, u)| \leq \int_0^\infty e^{\epsilon t} |v|_{\delta,\infty} e^{\delta(u-t)} dt = C |v|_{\delta,\infty} e^{\delta u}$$

and

$$\begin{aligned} |(\widehat{E}(s)v)(y, u) - (\widehat{E}(s)v)(y', u)| &\leq \int_0^\infty e^{\epsilon t} |v|_{\delta,\eta} d(y, y')^\eta e^{\delta(u-t)} dt \\ &= C e^{\delta u} |v|_{\delta,\eta} d(y, y')^\eta. \end{aligned}$$

That is, $|\widehat{E}(s)v|_{\delta,\infty} \leq |v|_{\delta,\infty}$ and $|\widehat{E}(s)v|_{\delta,\eta} \leq |v|_{\delta,\eta}$.

(c) Let $v \in \mathcal{F}_{\epsilon,\eta}(Y^\varphi)$. Let $(y, u), (y', u) \in \widetilde{Y}_j$, $j \geq 1$. Then $(H_t v)(y, u) = 1_{\{t < u\}} v(y, u - t)$ and $(\widehat{H}(s)v)(y, u) = \int_0^u e^{-st} v(y, u - t) dt$. Hence

$$|\widehat{H}(s)v|_\infty \leq e^\delta |v|_{\delta,\infty}, \quad |(\widehat{H}(s)v)(y, u) - (\widehat{H}(s)v)(y', u)| \leq e^\delta |v|_{\delta,\eta} d(y, y')^\eta.$$

The result follows. ■

PROPOSITION 3.8. *There exist $\delta > \epsilon > 0$ and $C > 0$ such that*

$$\|\widehat{B}(s)\|_{\mathcal{F}_{\delta,\eta}(Y^\varphi) \rightarrow \mathcal{F}_\eta(\widetilde{Y})} \leq C|s| \quad \text{and} \quad \|\widehat{G}(s)\|_{\mathcal{F}_{\delta,\eta}(Y^\varphi) \rightarrow \mathcal{F}_\eta(\widetilde{Y})} \leq C|s|$$

for $|a| \leq \epsilon$.

Proof. Let $v \in L^1(Y^\varphi)$, $w \in L^\infty(\widetilde{Y})$. Using $F_t(y, u) = (Fy, u + t - \varphi(y))$ for $(y, u) \in \Delta_t$, we have

$$\begin{aligned} \int_{\widetilde{Y}} B_t v w d\tilde{\mu} &= \bar{\varphi} \int_{Y^\varphi} L_t(1_{\Delta_t} v) w d\mu^\varphi = \bar{\varphi} \int_{Y^\varphi} 1_{\Delta_t} v w \circ F_t d\mu^\varphi \\ &= \int_Y \int_0^{\varphi(y)} 1_{\{0 \leq u+t-\varphi(y) < 1\}} v(y, u) w(Fy, u + t - \varphi) du d\mu \\ &= \int_Y \int_{t-\varphi(y)}^t 1_{\{0 \leq u < 1\}} v(y, u + \varphi(y) - t) w(Fy, u) du d\mu \\ &= \int_{\widetilde{Y}} v_t w \circ \widetilde{F} d\tilde{\mu} = \int_{\widetilde{Y}} \widetilde{R} v_t w d\tilde{\mu}, \end{aligned}$$

where $v_t(y, u) = 1_{\{0 < u + \varphi(y) - t < \varphi(y)\}} v(y, u + \varphi(y) - t)$.

Hence $B_t v = \widetilde{R} v_t$ and it follows immediately that $G_t v = \widetilde{R}(\omega(t)v)_t$. But $(\omega(t)v)_t(y, u) = 1_{\{0 < u + \varphi(y) - t < \varphi(y)\}} (\omega(t)v)(y, u + \varphi(y) - t) = (1 - u)v_t(y, u)$, so $(G_t v)(y, u) = (1 - u)(B_t v)(y, u)$.

Next, $\widehat{B}(s)v = \widetilde{R}\widehat{V}(s)$ where

$$\begin{aligned}\widehat{V}(s)(y, u) &= \int_0^\infty e^{-st} v_t(y, u) dt = \int_u^{u+\varphi(y)} e^{-st} v(y, u + \varphi(y) - t) dt \\ &= \int_0^{\varphi(y)} e^{-s(\varphi(y)+u-t)} v(y, t) dt.\end{aligned}$$

It is immediate that

$$(3.1) \quad (\widehat{G}(s)v)(y, u) = (1 - u)(\widehat{B}(s)v)(y, u).$$

Suppose that $\delta > \epsilon > 0$ are fixed. Let $v \in \mathcal{F}_{\delta, \eta}(Y^\varphi)$. Let $(y, u), (y', u) \in \widetilde{Y}_j, j \geq 1$. Then

$$|\widehat{V}(s)(y, u)| \leq \int_0^{\varphi(y)} e^{-a(\varphi(y)+u-t)} |v|_{\delta, \infty} e^{\delta t} dt \ll e^{\delta \varphi(y)} |v|_{\delta, \infty}$$

and so $|1_{Y_j} \widehat{V}(s)|_\infty \ll e^{\delta |1_{Y_j} \varphi|_\infty} |v|_{\delta, \infty}$.

Next, suppose without loss of generality that $\varphi(y') \leq \varphi(y)$. Then

$$\widehat{V}(s)(y, u) - \widehat{V}(s)(y', u) = J_1 + J_2 + J_3,$$

where

$$\begin{aligned}J_1 &= \int_0^{\varphi(y)} (e^{-s(\varphi(y)+u-t)} - e^{-s(\varphi(y')+u-t)}) v(y, t) dt, \\ J_2 &= \int_0^{\varphi(y)} e^{-s(\varphi(y')+u-t)} (v(y, t) - v(y', t)) dt, \\ J_3 &= \int_{\varphi(y')}^{\varphi(y)} e^{-s(\varphi(y')+u-t)} v(y', t) dt.\end{aligned}$$

For notational convenience we suppose that $a \in (-\epsilon, 0)$ since the range $a \geq 0$ is simpler. Using (2.1),

$$\begin{aligned}|J_1| &\leq \int_0^{\varphi(y)} e^{\epsilon(|1_{Y_j} \varphi|_\infty + 1 - t)} |s| |\varphi(y) - \varphi(y')| |v|_{\delta, \infty} e^{\delta t} dt \\ &\ll |s| \varphi(y) e^{\delta |1_{Y_j} \varphi|_\infty} d(Fy, Fy')^\eta |v|_{\delta, \infty} \ll |s| e^{2\delta |1_{Y_j} \varphi|_\infty} d(Fy, Fy')^\eta |v|_{\delta, \infty}, \\ |J_2| &\leq \int_0^{\varphi(y)} e^{\epsilon(|1_{Y_j} \varphi|_\infty + 1 - t)} |v|_{\delta, \eta} e^{\delta t} d(y, y')^\eta dt \ll e^{\delta |1_{Y_j} \varphi|_\infty} d(y, y')^\eta |v|_{\delta, \eta}, \\ |J_3| &\leq \int_{\varphi(y')}^{\varphi(y)} e^{\epsilon(|1_{Y_j} \varphi|_\infty + 1 - t)} |v|_{\delta, \infty} e^{\delta t} dt \ll e^{2\delta |1_{Y_j} \varphi|_\infty} |v|_{\delta, \infty} d(Fy, Fy')^\eta.\end{aligned}$$

Hence

$$|\widehat{V}(s)(y, u) - \widehat{V}(s)(y, u)| \ll |s| e^{2\delta|1_{Y_j, \varphi}|_\infty} \|v\|_{\delta, \eta} d(Fy, Fy')^\eta.$$

Now, for $(y, u) \in \widetilde{Y}$,

$$(\widehat{B}(s)v)(y, u) = (\widetilde{R}\widehat{V}(s))(y, u) = \sum_j g(y_j) \widehat{V}(s)(y_j, u),$$

where y_j is the unique preimage of y under $F|_{Y_j}$. It follows from the estimates for $\widehat{V}(s)$ together with (2.3) that

$$\|\widehat{B}(s)v\|_\eta \ll |s| \sum_j \mu(Y_j) e^{2\delta|1_{Y_j, \varphi}|_\infty} \|v\|_{\delta, \eta}.$$

Shrinking δ , we deduce the desired estimate for \widehat{B} from (2.2). Finally, the estimate for \widehat{G} follows from (3.1). ■

PROPOSITION 3.9. $\int_{\widetilde{Y}} \widehat{B}(0)v d\tilde{\mu} = \bar{\varphi} \int_{Y^\varphi} v d\mu^\varphi$ for $v \in L^1(Y^\varphi)$.

Proof. By the definition of \widehat{B} ,

$$\begin{aligned} \int_{\widetilde{Y}} \widehat{B}(0)v d\tilde{\mu} &= \int_{\widetilde{Y}} \int_0^\infty L_t(1_{\Delta_t}v) dt d\tilde{\mu} = \bar{\varphi} \int_0^\infty \int_{Y^\varphi} L_t(1_{\Delta_t}v) d\mu^\varphi dt \\ &= \bar{\varphi} \int_0^\infty \int_{Y^\varphi} 1_{\Delta_t}v d\mu^\varphi dt = \bar{\varphi} \int_{Y^\varphi} \int_0^\infty 1_{\{\varphi-u < t < \varphi-u+1\}} v dt d\mu^\varphi \\ &= \bar{\varphi} \int_{Y^\varphi} v d\mu^\varphi, \end{aligned}$$

as required. ■

LEMMA 3.10. Write $s = a + ib \in \mathbb{C}$. There exist $\epsilon > 0$, $\delta > 0$, $m_2 \geq 0$, $C > 0$ such that

- (a) $s \mapsto \widehat{L}(s) : \mathcal{F}_{\delta, \eta}^0(Y^\varphi) \rightarrow \mathcal{F}_{\delta, \eta}(Y^\varphi)$ is analytic on $\{|a| < \epsilon\}$;
- (b) $s \mapsto \widehat{L}(s) : \mathcal{F}_{\delta, \eta}(Y^\varphi) \rightarrow \mathcal{F}_{\delta, \eta}(Y^\varphi)$ is analytic on $\{|a| < \epsilon\}$ except for a simple pole at $s = 0$;
- (c) $\|\widehat{L}(s)v\|_{\delta, \eta} \leq C|b|^{m_2} \|v\|_{\delta, \eta}$ for $|a| \leq \epsilon$, $|b| \geq 1$, $v \in \mathcal{F}_{\delta, \eta}(Y^\varphi)$.

Proof. Recall that

$$\widehat{L} = \widehat{A}\widehat{T}\widehat{B} + \widehat{A}\widehat{G} + \widehat{A}\widehat{H} + \widehat{E}, \quad \widehat{T} = \widehat{U}(I - \widehat{R})^{-1},$$

where \widehat{U} , \widehat{A} , \widehat{B} , \widehat{G} , \widehat{H} and \widehat{E} are analytic by Propositions 3.2, 3.7 and 3.8. Hence part (b) follows immediately from Lemma 3.1(b). Also, part (c) is a consequence of Lemma 3.1(c).

By Proposition 3.9, $\widehat{B}(0)(\mathcal{F}_{\delta, \eta}^0(Y^\varphi)) \subset \mathcal{F}_\eta^0(\widetilde{Y})$. Hence the simple pole at $s = 0$ for $(I - \widehat{R})^{-1}\widehat{B}$ disappears on restriction to $\mathcal{F}_{\delta, \eta}^0(Y^\varphi)$ by Lemma 3.1(a). This proves part (a). ■

3.3. Moving the contour of integration

PROPOSITION 3.11. *Let $m \geq 1$. Let $v \in \mathcal{F}_{\delta,\eta,m}(Y^\varphi)$ have good support. Then $\widehat{L}(s)v = \sum_{j=0}^{m-1} (-1)^j s^{-(j+1)} \partial_t^j v + (-1)^m s^{-m} \widehat{L}(s) \partial_t^m v$ for $a > 0$.*

Proof. Recall that $\text{supp } v \subset \{(y, u) \in Y^\varphi : u \in [r, \varphi(y) - r]\}$ for some $r > 0$. For $h \in [0, r]$, we can define $(\Psi_h v)(y, u) = v(y, u - h)$ and then $(\Psi_h v) \circ F_h = v$.

Let $w \in L^\infty(Y^\varphi)$ and write $\rho_{v,w}(t) = \int_{Y^\varphi} v w_t d\mu^\varphi$, where $w_t = w \circ F_t$. Then for $h \in [0, r]$,

$$\rho_{v,w}(t+h) = \int_{Y^\varphi} v w_t \circ F_h d\mu^\varphi = \int_{Y^\varphi} (\Psi_h v) \circ F_h w_t \circ F_h d\mu^\varphi = \int_{Y^\varphi} \Psi_h v w_t d\mu^\varphi.$$

Hence $h^{-1}(\rho_{v,w}(t+h) - \rho_{v,w}(t)) = \int_{Y^\varphi} h^{-1}(\Psi_h v - v) w_t d\mu^\varphi$, so

$$\rho'_{v,w}(t) = - \int_{Y^\varphi} \partial_t v w_t d\mu^\varphi = - \int_{Y^\varphi} \partial_t v w \circ F_t d\mu^\varphi = -\rho_{\partial_t v, w}(t).$$

Inductively, $\rho_{v,w}^{(j)}(t) = (-1)^j \rho_{\partial_t^j v, w}(t)$.

Now $\int_{Y^\varphi} \widehat{L}(s) v w d\mu^\varphi = \int_0^\infty e^{-st} \int_{Y^\varphi} L_t v w d\mu^\varphi dt = \int_0^\infty e^{-st} \rho_{v,w}(t) dt$, so repeatedly integrating by parts, we get

$$\begin{aligned} \int_{Y^\varphi} \widehat{L}(s) v w d\mu^\varphi &= \sum_{j=0}^{m-1} s^{-(j+1)} \rho_{v,w}^{(j)}(0) + s^{-m} \int_0^\infty e^{-st} \rho_{v,w}^{(m)}(t) dt \\ &= \sum_{j=0}^{m-1} (-1)^j s^{-(j+1)} \rho_{\partial_t^j v, w}(0) + (-1)^m s^{-m} \int_0^\infty e^{-st} \rho_{\partial_t^m v, w}(t) dt \\ &= \int_{Y^\varphi} \sum_{j=0}^{m-1} (-1)^j s^{-(j+1)} \partial_t^j v w d\mu^\varphi + (-1)^m s^{-m} \int_0^\infty e^{-st} \rho_{\partial_t^m v, w}(t) dt. \end{aligned}$$

Finally, $\int_0^\infty e^{-st} \rho_{\partial_t^m v, w}(t) dt = \int_{Y^\varphi} \widehat{L}(s) \partial_t^m v w d\mu^\varphi$ and the result follows since $w \in L^\infty(Y^\varphi)$ is arbitrary. ■

We can now estimate $\|L_t v\|_{\delta,\eta}$.

COROLLARY 3.12. *Under the assumptions of Theorem 2.3, there exist $\epsilon > 0$, $m_3 \geq 1$, $C > 0$ such that*

$$\|L_t v\|_{\delta,\eta} \leq C e^{-\epsilon t} \|v\|_{\delta,\eta,m_3} \quad \text{for all } t > 0,$$

for all $v \in \mathcal{F}_{\delta,\eta,m_3}^0(Y^\varphi)$ with good support.

Proof. Let $m_3 = m_2 + 2$. By Lemma 3.10(a), $\widehat{L}(s) : \mathcal{F}_{\delta,\eta,m_3}^0(Y^\varphi) \rightarrow \mathcal{F}_{\delta,\eta}(Y^\varphi)$ is analytic for $|a| \leq \epsilon$. The alternative expression in Proposition 3.11 is also analytic on this region (the apparent singularity at $s = 0$ is removable by the equality with the analytic function \widehat{L}). Hence we can

move the contour of integration to $s = -\epsilon + ib$ when computing the inverse Laplace transform, to obtain

$$\begin{aligned} L_t v &= \int_{-\infty}^{\infty} e^{st} \left(\sum_{j=0}^{m_3-1} (-1)^j s^{-(j+1)} \partial_t^j v + (-1)^{m_3} s^{-m_3} \widehat{L}(s) \partial_t^{m_3} v \right) db \\ &= e^{-\epsilon t} \sum_{j=0}^{m_3-1} (-1)^j \partial_t^j v \int_{-\infty}^{\infty} e^{ibt} s^{-(j+1)} db \\ &\quad + (-1)^{m_3} e^{-\epsilon t} \int_{-\infty}^{\infty} e^{ibt} s^{-m_3} \widehat{L}(s) \partial_t^{m_3} v db. \end{aligned}$$

The final term is estimated using Lemma 3.10(b, c):

$$\begin{aligned} &\left\| \int_{-\infty}^{\infty} e^{ibt} s^{-m_3} \widehat{L}(s) \partial_t^{m_3} v db \right\|_{\delta, \eta} \\ &\ll \int_{-\infty}^{\infty} (1 + |b|)^{-(m_2+2)} (1 + |b|)^{m_2} \|\partial_t^{m_3} v\|_{\delta, \eta} db \ll \|v\|_{\delta, \eta, m_3}. \end{aligned}$$

Clearly, the integrals $\int_{-\infty}^{\infty} e^{ibt} s^{-(j+1)} db$ converge absolutely for $j \geq 1$, while the integral for $j = 0$ converges as an improper Riemann integral. Hence altogether we obtain $\|L_t v\|_{\delta, \eta} \ll e^{-\epsilon t} \|v\|_{\delta, \eta, m_3}$. ■

For the proof of Theorem 2.3, it remains to estimate $\|\partial_u L_t v\|_{\delta, \eta}$. Recall that the transfer operator R_0 for F has weight function g . We have the pointwise formula

$$(R_0^k v)(y) = \sum_{F^k y' = y} g_k(y') v(y') \quad \text{where} \quad g_k = g \dots g \circ F^{k-1}.$$

Let

$$\varphi_k = \sum_{j=0}^{k-1} \varphi \circ F^j.$$

PROPOSITION 3.13. *Let $v \in L^1(Y^\varphi)$. Then for all $t > 0$, $(y, u) \in Y^\varphi$,*

$$(L_t v)(y, u) = \sum_{k=0}^{\lfloor t/2 \rfloor} \sum_{F^k y' = y} g_k(y') 1_{\{0 \leq u - t + \varphi_k(y') < \varphi(y')\}} v(y', u - t + \varphi_k(y')).$$

Proof. Recall that the roof function φ is bounded below by 2. The lap number $N_t(y, u) \in [0, t/2] \cap \mathbb{N}$ is the unique integer $k \geq 0$ such that $u + t - \varphi_k(y) \in [0, \varphi(F^k y))$. In particular, $F_t(y, u) = (F^{N_t(y, u)} y, u + t - \varphi_{N_t(y, u)}(y))$.

For $w \in L^\infty(Y^\varphi)$,

$$\begin{aligned} \int_{Y^\varphi} L_t(1_{\{N_t=k\}}v) w d\mu^\varphi &= \int_{Y^\varphi} 1_{\{N_t=k\}}v w \circ F_t d\mu^\varphi \\ &= \bar{\varphi}^{-1} \int_Y \int_0^{\varphi(y)} 1_{\{0 \leq u+t-\varphi_k(y) < \varphi(F^k y)\}} v(y, u) w(F^k y, u+t-\varphi_k(y)) du d\mu \\ &= \bar{\varphi}^{-1} \int_Y \int_0^{\varphi(F^k y)} 1_{\{0 \leq u-t+\varphi_k(y) < \varphi(y)\}} v(y, u-t+\varphi_k(y)) w(F^k y, u) du d\mu. \end{aligned}$$

Writing $v_{t,k}^u(y) = 1_{\{0 \leq u-t+\varphi_k(y) < \varphi(y)\}} v(y, u-t+\varphi_k(y))$ and $w^u(y) = w(y, u)$, we get

$$\begin{aligned} \int_{Y^\varphi} L_t(1_{\{N_t=k\}}v) w d\mu^\varphi &= \bar{\varphi}^{-1} \int_0^\infty \int_Y 1_{\{u < \varphi \circ F^k\}} v_{t,k}^u w^u \circ F^k d\mu du \\ &= \bar{\varphi}^{-1} \int_0^\infty \int_Y 1_{\{u < \varphi\}} R_0^k v_{t,k}^u w^u d\mu du = \int_{Y^\varphi} (R_0^k v_{t,k}^u)(y) w(y, u) d\mu^\varphi. \end{aligned}$$

Hence

$$(L_t v)(y, u) = \sum_{k=0}^{\lfloor t/2 \rfloor} (L_t(1_{\{N_t=k\}}v))(y, u) = \sum_{k=0}^{\lfloor t/2 \rfloor} (R_0^k v_{t,k}^u)(y).$$

The result follows from the pointwise formula for R_0^k . ■

Proof of Theorem 2.3. Let $m = m_3 + 1$. By Corollary 3.12, we have $\|L_t v\|_{\delta, \eta} \ll e^{-ct} \|v\|_{\delta, \eta, m}$.

Recall that ∂_u denotes the ordinary derivative with respect to u at $0 < u < \varphi(y)$ and denotes the appropriate one-sided derivative at $u = 0$ and $u = \varphi(y)$. Since v has good support, the indicator functions in the right-hand side of the formula in Proposition 3.13 are constant on the support of v . It follows that $\partial_u L_t v = L_t(\partial_u v)$. By Corollary 3.12,

$$\|\partial_u L_t v\|_{\delta, \eta} = \|L_t(\partial_u v)\|_{\delta, \eta} \ll e^{-ct} \|\partial_u v\|_{\delta, \eta, m_3} \leq e^{-ct} \|v\|_{\delta, \eta, m}.$$

Hence, $\|L_t v\|_{\delta, \eta, 1} \ll e^{-ct} \|v\|_{\delta, \eta, m}$ as required. ■

Acknowledgements. We are grateful to the referee for a number of helpful suggestions.

References

- [1] J. Aaronson, *An Introduction to Infinite Ergodic Theory*, Math. Surveys Monogr. 50, Amer. Math. Soc., Providence, RI, 1997.

- [2] J. Aaronson and M. Denker, *Local limit theorems for partial sums of stationary sequences generated by Gibbs–Markov maps*, Stoch. Dynam. 1 (2001), 193–237.
- [3] J. Aaronson, M. Denker and M. Urbański, *Ergodic theory for Markov fibred systems and parabolic rational maps*, Trans. Amer. Math. Soc. 337 (1993), 495–548.
- [4] M. Antoniou and I. Melbourne, *Rate of convergence in the weak invariance principle for deterministic systems*, Comm. Math. Phys. 369 (2019), 1147–1165.
- [5] V. Araújo and I. Melbourne, *Exponential decay of correlations for nonuniformly hyperbolic flows with a $C^{1+\alpha}$ stable foliation, including the classical Lorenz attractor*, Ann. Henri Poincaré 17 (2016), 2975–3004.
- [6] A. Avila, S. Gouëzel and J.-C. Yoccoz, *Exponential mixing for the Teichmüller flow*, Publ. Math. Inst. Hautes Études Sci. 104 (2006), 143–211.
- [7] V. Baladi and B. Vallée, *Exponential decay of correlations for surface semi-flows without finite Markov partitions*, Proc. Amer. Math. Soc. 133 (2005), 865–874.
- [8] R. Bowen, *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*, Lecture Notes in Math 470, Springer, Berlin, 1975.
- [9] H. Bruin, I. Melbourne and D. Terhesiu, *Rates of mixing for non-Markov infinite measure semiflows*, Trans. Amer. Math. Soc. 371 (2019), 7343–7386.
- [10] O. Butterley, *A note on operator semigroups associated to chaotic flows*, Ergodic Theory Dynam. Systems 36 (2016), 1396–1408.
- [11] D. Dolgopyat, *On the decay of correlations in Anosov flows*, Ann. of Math. 147 (1998), 357–390.
- [12] D. Dolgopyat, *Prevalence of rapid mixing in hyperbolic flows*, Ergodic Theory Dynam. Systems 18 (1998), 1097–1114.
- [13] P. Góra and A. Boyarsky, *Absolutely continuous invariant measures for piecewise expanding C^2 transformations in R^N* , Israel J. Math. 67 (1989), 272–286.
- [14] S. Gouëzel, *Berry–Esseen theorem and local limit theorem for non uniformly expanding maps*, Ann. Inst. H. Poincaré Probab. Statist. 41 (2005), 997–1024.
- [15] F. Hofbauer and G. Keller, *Ergodic properties of invariant measures for piecewise monotonic transformations*, Math. Z. 180 (1982), 119–140.
- [16] G. Keller, *Generalized bounded variation and applications to piecewise monotonic transformations*, Z. Wahrsch. Verw. Gebiete 69 (1985), 461–478.
- [17] C. Liverani, *On contact Anosov flows*, Ann. of Math. 159 (2004), 1275–1312.
- [18] C. Liverani, *Multidimensional expanding maps with singularities: a pedestrian approach*, Ergodic Theory Dynam. Systems 33 (2013), 168–182.
- [19] I. Melbourne, *Superpolynomial and polynomial mixing for semiflows and flows*, Nonlinearity 31 (2018), R268–R316.
- [20] I. Melbourne, N. Paviato and D. Terhesiu, *Nonexistence of spectral gaps in Hölder spaces for continuous time dynamical systems*, Israel J. Math. 247 (2022), 987–991.
- [21] I. Melbourne and D. Terhesiu, *Operator renewal theory for continuous time dynamical systems with finite and infinite measure*, Monatsh. Math. 182 (2017), 377–431.
- [22] M. Pollicott, *On the rate of mixing of Axiom A flows*, Invent. Math. 81 (1985), 413–426.
- [23] D. Ruelle, *Thermodynamic Formalism*, Encyclopedia Math. Appl. 5, Addison-Wesley, Reading, MA, 1978.
- [24] M. Rychlik, *Bounded variation and invariant measures*, Studia Math. 76 (1983), 69–80.
- [25] B. Saussol, *Absolutely continuous invariant measures for multidimensional expanding maps*, Israel J. Math. 116 (2000), 223–248.
- [26] Ya. G. Sinai, *Gibbs measures in ergodic theory*, Russian Math. Surveys 27 (1972), no. 4, 21–69.
- [27] M. Tsujii, *Decay of correlations in suspension semi-flows of angle-multiplying maps*, Ergodic Theory Dynam. Systems 28 (2008), 291–317.

- [28] M. Tsujii, *Quasi-compactness of transfer operators for contact Anosov flows*, *Nonlinearity* 23 (2010), 1495–1545.
- [29] L.-S. Young, *Statistical properties of dynamical systems with some hyperbolicity*, *Ann. of Math.* 147 (1998), 585–650.

Ian Melbourne, Nicolò Paviato
Mathematics Institute
University of Warwick
Coventry, CV4 7AL, UK
E-mail: i.melbourne@warwick.ac.uk
nicolo.paviato@warwick.ac.uk

Dalia Terhesiu
Mathematisch Instituut
University of Leiden
Niels Bohrweg 1
2333 CA Leiden, Netherlands
E-mail: d.e.terhesiu@math.leidenuniv.nl