

3G6 COMMUTATIVE ALGEBRA - HOMEWORK 3

DUE TUESDAY 23 FEBRUARY, 2PM

Hand in the problems in Section B *only* to the boxes outside the undergraduate office. You are encouraged to work together on the problems, but your written work should be your own.

A : WARM-UP PROBLEMS

- (1) (If you didn't do this when working on HW2) Let I be an ideal of a ring R . Show that there is a bijection between ideals in R/I and ideals in R containing I .
- (2) Let $R = \mathbb{Z}/10\mathbb{Z}$, and let $U = \{1, 5\}$. Describe $R[U^{-1}]$.
- (3) Let R be a local ring with maximal ideal \mathfrak{m} . What happens when we localize R at \mathfrak{m} ? (i.e., form $R_{\mathfrak{m}} = R[(R \setminus \mathfrak{m})^{-1}]$).
- (4) Show that $\langle x^2 + y^2, xy \rangle \subset \mathbb{C}[x, y]$ is not a free module.
- (5) Let $A = \begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix}$, where the entries are in $\mathbb{Z}/5\mathbb{Z}$. Compute the characteristic polynomial of A .

B: EXERCISES

- (1) Show that if R is a local ring with maximal ideal \mathfrak{m} , then $R_{\mathfrak{m}} \cong R$.
- (2) Let $R = K[x, y]/\langle xy \rangle$, and let $P = \langle x, y-1 \rangle$, and $P' = \langle x-1, y \rangle$ be two ideals in R . Show that P and P' are prime. Describe $\text{Spec}(R_P)$ and $\text{Spec}(R_{P'})$.
- (3) Let R be a domain, and let $f \in R$ be a nonzero non-unit. Let $U = \{1, f, f^2, f^3, \dots\}$. We write $R[1/f]$ for $R[U^{-1}]$. Prove that $R[1/f]$ is not finitely-generated as an R -module. Give an example to show that this might not be true if R is not a domain (ie give an example of R that is not a domain, and $f \in R$ with $R[1/f]$ finitely generated as an R -module. Be sure to carefully justify your answer!
- (4) Let U be a multiplicatively closed subset of a ring R . Show that if P is an ideal of R that is maximal with respect to the property $P \cap U = \emptyset$, then P is prime. Here “maximal with respect to the property ...” means that if $I \supset P$ is an ideal with $I \cap U = \emptyset$, then $I = P$.

- (5) Let $R = \mathbb{C}[x, y]$, and let $\phi: R^2 \rightarrow R^2$ be given by the matrix

$$A = \begin{pmatrix} f_{11} & f_{12} \\ f_{21} & f_{22} \end{pmatrix},$$

where $f_{ij} \in \mathbb{C}[x, y]$. Give necessary and sufficient conditions for ϕ to be an isomorphism. You must completely justify your answer.

- (6) (Eisenbud Exercise 2.6: Generalized Chinese Remainder Theorem). Let R be a ring, and let Q_1, \dots, Q_n be ideals of R such that $Q_i + Q_j = R$ for all $i \neq j$. Show that $R/(\cap_i Q_i) \cong \prod_i R/Q_i$ as follows:
- Consider the map of rings $\phi: R \rightarrow \prod_i R/Q_i$ obtained from the n projection maps $R \rightarrow R/Q_i$. Show that $\ker \phi = \cap_i Q_i$.
 - Let \mathfrak{m} be a maximal ideal of R . Show that the hypothesis that $Q_i + Q_j = R$ for all $i \neq j$ means that at most one of the Q_i is contained in \mathfrak{m} . Use this to show that ϕ is surjective.

C: EXTENSIONS

- Show that localization has the following universal property: If $\phi: R \rightarrow S$ is a ring homomorphism which takes all elements of $U \subset R$ to units of S , then there is a unique induced ring homomorphism from $R[U^{-1}]$ to S . Show that this property defines the localization up to unique isomorphism: if T is a ring for which every ring homomorphism from R to a ring S that takes elements of U to units of S factors uniquely through T , then T is uniquely isomorphic to $R[U^{-1}]$.
- Let R be a ring. Show that an ideal that is maximal with respect to not being finitely generated (ie any larger ideal is finitely generated) is prime.
- Let R be a ring. Show that an ideal that is maximal among those that are not principal is prime.
- Generalize your answer to Question B5.