

MA3D5 Galois theory

Instructions: The assignments count for 15% of the credit for the course. There are 5 worksheets, whose purpose is to ensure students do the bare minimum work required to keep up with the course; they are not intended to be difficult, and you can discuss them in the support class. Solutions must be left in the filing cabinet by the front office by the

deadlines: 10:00 am on Thu of Weeks 2, 4, 6, 8 plus Term 3, Week 1 (Thu 21st Apr 2005).

We'll arrive at the total marks by taking the first, the best 2 out of the next 3, plus the last. In other words, the first sheet covering prerequisites and the final sheet including past exam questions are intended as compulsory vacation work.

I would be grateful for comments on the worksheets or the course material.

Preliminary warm-up (not for credit)

P.1 See Section 1.4 of the notes for elementary symmetric functions. If $\sigma_1, \dots, \sigma_n$ are elementary symmetric functions of $\alpha_1, \dots, \alpha_n$, compute the elementary symmetric functions of $\alpha_1^2, \dots, \alpha_n^2$.

P.2 Suppose that $f(x) = x^2 + ax + b = 0$ has roots α, β . Find the quadratic polynomial with roots

$$(a) \alpha^2, \beta^2; \quad (b) \alpha^3, \beta^3; \quad (c) \alpha^7\beta^4, \alpha^4\beta^7.$$

P.3 Write $\varepsilon := \exp \frac{2\pi i}{5}$ for the natural primitive 5th root of 1 as in Section 1.5 of the notes; it is a root of the quartic $f(x) = x^4 + x^3 + x^2 + x + 1 = 0$. Find the quadratic equation whose two roots are $\varepsilon + \varepsilon^4$ and $\varepsilon^2 + \varepsilon^3$, and hence give radical formulas for $\cos \frac{2\pi}{5}$, $\cos \frac{4\pi}{5}$. Ditto for \sin .

P.4 Consider the “reciprocal” or “palindromic” quartic polynomial

$$f(x) = ax^4 + bx^3 + cx^2 + bx + a.$$

If α is a root, prove that so is $1/\alpha$.

Show that $\frac{1}{x^2}f(x)$ can be written as a quadratic polynomial in $y = x + \frac{1}{x}$. Use this to find a formula for the roots of f involving two square root signs.

Assignment A

Deadline: Thu 13th Jan 2005 at 10:00

A.1 (i) If $f(x) = a_0x^n + a_1x^{n-1} + \dots + a_n$ has roots $\alpha_1, \dots, \alpha_n$, use elementary symmetric functions to find the polynomial with roots $c\alpha_1, c\alpha_2, \dots, c\alpha_n$. Say why the result is not surprising.

(ii) Use elementary symmetric polynomials to find the polynomial whose roots are $1/\alpha_1, \dots, 1/\alpha_n$. Check against common sense.

A.2 Suppose that α, β, γ are the three roots of the cubic equation $f(x) = x^3 + 3px + 2q = 0$, and let ω be a primitive cube root of 1 as in Section 1.5 of the notes. Study the effect of permuting α, β, γ on the quantities

$$t_1 = \alpha + \omega\beta + \omega^2\gamma \quad \text{and} \quad t_2 = \alpha + \omega^2\beta + \omega\gamma.$$

Deduce that t_1^3, t_2^3 are invariant under the 3-cycle $(\alpha\beta\gamma)$ and are interchanged by the transposition $(\beta\gamma)$. Use elementary symmetric functions to find the quadratic polynomial with roots t_1^3, t_2^3 .

A.3 Let $\varepsilon := \exp \frac{2\pi i}{5}$ be the natural primitive 5th root of 1.

Calculate from first principles the elementary symmetric functions of

(i) the 5 roots $\varepsilon^0 = 1, \varepsilon, \varepsilon^2, \varepsilon^3, \varepsilon^4$ of $x^5 - 1$; and

(ii) the 4 roots $\varepsilon, \varepsilon^2, \varepsilon^3, \varepsilon^4$ of $x^4 + x^3 + x^2 + x + 1$.

[Hint: The 5 roots are vertexes of a regular 5-gon centred at 0.]

A.4 Carry out in detail the calculation to prove formulas (1.12) of the notes for the roots of a quartic. Namely, if

$$\left. \begin{aligned} \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 &= 0, \\ \alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_1\alpha_4 + \alpha_2\alpha_3 + \alpha_2\alpha_4 + \alpha_3\alpha_4 &= r, \\ \alpha_1\alpha_2\alpha_3 + \alpha_1\alpha_2\alpha_4 + \alpha_1\alpha_3\alpha_4 + \alpha_2\alpha_3\alpha_4 &= -s, \\ \alpha_1\alpha_2\alpha_3\alpha_4 &= t, \end{aligned} \right\}$$

and u, v, w are defined by

$$\left. \begin{aligned} 2\alpha_1 &= u + v + w, \\ 2\alpha_2 &= u - v - w, \\ 2\alpha_3 &= -u + v - w, \\ 2\alpha_4 &= -u - v + w. \end{aligned} \right\} \quad (1)$$

then prove that

$$\left. \begin{aligned} u^2 + v^2 + w^2 &= -2r, \\ uvw &= -s, \\ u^2v^2 + u^2w^2 + v^2w^2 &= r^2 - 4t. \end{aligned} \right\} \quad (2)$$

How to use Maple

This is not part of the assignment sheet or of the course, but I recommend doing it. At a unix terminal, type maple; you get a clever logo, and the prompt `>`. For example, you can calculate $\sum \alpha_i^3$ in terms of elementary symmetric functions (*and get the right answer!*) by the following few lines:

```
> s1:=a+b+c; s2:=a*b+a*c+b*c; s3:=a*b*c;
      s1 := a + b + c
      s2 := a b + a c + b c
      s3 := a b c
> expand(a^3+b^3+c^3-s1^3);
      2      2      2      2      2      2
      - 3 a b - 3 a c - 3 a b - 6 a b c - 3 a c - 3 b c - 3 b c
> expand(a^3+b^3+c^3-s1^3+3*s1*s2);
      3 a b c
> evalb(expand(s1^3-3*s1*s2+3*a*b*c) = a^3+b^3+c^3);
      true
```

Mathematica is very similar.