

## MA4L7 Algebraic curves

### Example Sheet 4. Deadline Wed 4th Mar at 12:00

**Exercise 4.1** For distinct  $a_i \in k$ , write

$$F = \prod (x - a_i) \quad \text{and} \quad G_i = \prod_{j \neq i} (x - a_j).$$

Prove that

$$\sum_i \lambda_i G_i \equiv 1$$

where

$$\lambda_i = \prod_{j \neq i} \frac{1}{a_i - a_j}$$

[Hint: Find zeroes of  $\sum \lambda_i G_i - 1$ .]

**Exercise 4.2** Let  $P_i \in \mathbb{A}^n$  be a finite set of distinct points. Prove that there exists polynomials  $f_i \in k[x_{1\dots n}]$  for which

$$f_i(P_j) = \delta_{ij} \quad (\text{Kronecker delta}).$$

[Hint: Use a projection to  $\mathbb{A}^1$  that separates the  $P_i$ , and apply Ex. 4.1.]

Deduce that there exists  $g_i \in m_{P_j}^N \subset k[x_{1\dots n}]$  (where  $m_{P_j}$  is the maximal ideal at  $P_j$ ) with  $g_i(P_i) = 1$ .

**Exercise 4.3** For  $k$  an algebraically closed field and  $I \subset k[x_{1\dots n}]$  an ideal defining a finite set  $V(I) = \{P_i\} \subset \mathbb{A}^n$ , prove that

$$k[x_{1\dots n}]/I \cong \bigoplus_{P_i} \mathcal{O}_{P_i}/I \cdot \mathcal{O}_{P_i}.$$

Use the coprime result of Ex. 4.3 together with the NSS to prove the natural map is surjective and injective. For more details, compare [Fulton, Algebraic curves, Section 9, Prop. 6].

**Exercise 4.4** Let  $C = C_a \subset \mathbb{P}^2$  be a nonsingular curve of degree  $a$  defined by the homogeneous polynomial  $F_a$ . Assume that  $C$  meets the line  $z = 0$  transversally in  $a$  points and set  $H$  for the divisor of  $z$ .

A rational function  $h \in k(C)$  can be written  $h = A/B$  with  $A, B$  homogeneous forms of the same degree  $d$ . If  $h = \mathcal{L}(C, nH)$ , prove that  $B$  is in

the ideal  $(z^n, F_a)$ , so that  $f$  can also be written as  $A'/z^n$ . This implies that the map  $k[x, y, z]_n \rightarrow \mathcal{L}(C, nH)$  discussed in Ex. 2.7 is surjective. [Hint: The assumption  $h \in k(C)$  is that its poles on  $C$  are cancelled locally by multiplication by  $z^n$ .] (Sorry, I don't have time to tidy this up. Ex. 4.3 only deals with the affine case.)

**Exercise 4.5 (Genus 1 curve as  $Q_1 \cap Q_2 \subset \mathbb{P}^3$ )** Let  $E$  be a genus 1 curve and  $P \in E$ . Assume known the treatment of  $R(E, P) = \bigoplus \mathcal{L}(E, nP)$  as  $k[x, y, z]/(f_6)$  with  $x, y, z$  of degree 1, 2, 3 and  $f_6$  the relation

$$z^2 = y^3 + ax^4y + bx^6.$$

Calculate the subring  $R(E, 4P) \subset R(E, P)$ . [Hint: Give names to the monomials of degree 4. Find two quadratic relations between these in  $\mathcal{L}(8P)$ , either as trivial coincidences between monomials, or involving multiples of  $f_6$ .]

Deduce that  $4P$  is very ample and that the image of  $\varphi_{4P}: E \rightarrow \mathbb{P}^3$  is an intersection of two quadrics.

**Exercise 4.6 (Curve of genus 3)** Let  $C$  be a curve of  $g = 3$ . Write out the dimensions  $l(nK_C)$  for  $n = 0, n = 1$  and  $n \geq 2$ . The canonical ring of  $C$  is

$$R(C, K_C) = \bigoplus_{n \geq 0} \mathcal{L}(C, nK_C).$$

Complete the calculations of  $R(C, K_C)$  given in lectures:

If  $\varphi_{K_C}: C \rightarrow \mathbb{P}^2$  is an embedding, prove that the hypersurface ring  $k[x_{0..2}]/f_4$  has the right dimension in each degree. Then the graded homomorphism  $k[x_{0..2}] \rightarrow R(C, K_C)$  is surjective by Max Noether's theorem, and the kernel consists exactly of the multiples of  $f_4$ .

If there is a quadratic relation  $q(x_{0..2}) = 0$ , prove that there is only one  $q$ . The space  $\mathcal{L}(2K_C)$  needs a further relation  $y$ . You have to figure out the number of monomials in  $x_{0..2}, y$  in each degree, and the number of monomials multiples of the relation  $q$  in each degree. You need to prove that there is a relation  $F_4$  involving  $y^2$ , and finally check that the dimension of  $k[x_{0..2}, y]/(q_2, F_4)$  matches the dimension of  $\mathcal{L}(nK_C)$  in each degree  $n \geq 4$ .

**Exercise 4.7 (Half-canonical divisor)** Let  $C$  be a curve of  $g = 3$  and  $P, Q \in C$  two points such that  $2(P + Q) \stackrel{\text{lin}}{\sim} K_C$ . Set  $A = P + Q$ , so that  $K_C = 2A$ . Assume that  $l(A) = 1$  (the cases  $l(A) = 0$  and  $l(A) = 2$  are also possible, and interesting, but left for another day).

Consider the sections ring  $R(C, A) = \bigoplus_{n \geq 0} \mathcal{L}(C, nA)$ . This clearly contains the canonical ring  $R(C, K_C) = R(C, 2A)$ . The question is to determine the possibilities for generators and relations for  $R(C, A)$ , by analogy with Ex. 4.5. Geometrically, there are two cases: if  $C = C_4 \subset \mathbb{P}^2$ , the line joining  $P, Q$  is a bitangent of  $C_4$ . If  $C$  is hyperelliptic then  $P, Q$  are ramification points of the double cover  $C \rightarrow \mathbb{P}^1$ .

Write out the dimension of  $\mathcal{L}(C, nA)$  for  $n = 0, 1, 2$  and  $n \geq 3$ .

Check that these dimensions coincide with those of the graded ring  $k[x, y_1, y_2, z]/(f_4, g_6)$ , where the generators have degrees 1, 2, 2, 3 and the relations have the indicated degrees.

In the hyperelliptic case, there is an extra quadratic relation of the form  $q(x^2, y_1, y_2) = 0$  (of weighted degree 4), and an extra generator  $t$  of degree 4, giving  $k[x, y_1, y_2, z, t]/(q_4, f_4, g_6)$ .

**Exercise 4.8 (Past exam question) Part 1.** The proof of RR used in the course was based on three main propositions. The first two of these are:

- (I) A principal divisor has degree zero:  $\deg(\operatorname{div} f) = 0$  for all  $f \in k(C)^\times$ .
- (II) There exists a sequence of divisors  $D_n$  of degree tending to  $+\infty$  such that the difference  $\deg D_n + 1 - l(D_n)$  is bounded.

Use (I) and (II) together with the standard methods of argument to prove the following results:

- (i) The maximum  $g = \max_D \{\deg D + 1 - l(D)\}$  taken over all divisors  $D$  is well defined, so that the Riemann–Roch inequality  $l(D) \geq 1 - g + \deg D$  is satisfied for every divisor  $D$ .
- (ii) With  $g$  as in (i), every divisor  $D$  of degree  $\geq g$  has  $l(D) > 0$ , so is linearly equivalent to an effective divisor.
- (iii) There exists a divisor  $D$  of degree  $g - 1$  for which  $l(D) = 0$ , so that the RR inequality is equality.
- (iv)  $l(D) = 1 - g + \deg D$  holds for every divisor  $D$  of degree  $\geq 2g - 1$ .

**Part 2.** Suppose that  $g(C) = 2$  and  $\deg D = 4$ . Prove that  $l(D - K_C) \neq 0$ , and deduce that  $\varphi_D$  is not an embedding. Show that  $\varphi_D$  is either a generically 2-to-1 map of  $C$  to a plain conic, or maps  $C$  birational to a quartic curve  $\overline{C}$  with a node or cusp as its only singularity. Explain which divisors  $D$  correspond to each case. [You may use the criteria on embeddings, and standard properties of the canonical map of a genus 2 curve.]

**Status:** Bookwork. The whole proof of RR is too long for an exam question, but it is fair to state parts of it as given, and ask for the proof of the next part. The rider Part 2 is too hard for an exam question, but was set as an earlier question.

**Exercise 4.9 (Harder question:  $g(C) = 3$  and  $\deg D = 5$  divisor)** Let  $C$  be a nonsingular curve of  $g = 3$  and  $D$  a divisor of degree 5, with  $D$  not linearly equivalent to  $K_C + P$ . Prove that the linear system  $|D|$  defines a birational map  $\varphi_D$  of  $C$  to a plane quintic  $\overline{C}_5 \subset \mathbb{P}^2$ . [Hint: Prove that  $|D|$  is a free  $g_5^2$ . Now, because 5 is a prime, the map  $\varphi_D$  cannot be a multiple cover of another curve, so must be generically one-to-one.]

A nonsingular plane quintic would have  $g = 6$ , so that  $\overline{C}_5$  must be singular. Consider the two main cases: (i)  $\overline{C}$  has distinct 3 nodes; and (ii)  $\overline{C}$  has an ordinary triple point.

If  $\overline{C}$  is a plane quintic with nodes at the three coordinate points, show that the standard quadratic transformation  $\psi: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$  given by  $(x, y, z) \mapsto (\frac{1}{x} : \frac{1}{y} : \frac{1}{z}) = (yz : xz : xy)$  takes  $\overline{C}$  to a nonsingular quartic, and the composite  $\psi \circ \varphi_D$  is the canonical embedding  $\varphi_{K_C}$  of a nonhyperelliptic  $C$  of  $g = 3$ .

If  $\overline{C}$  is a plane quintic with an ordinary triple point  $P$ , show that the linear projection from  $P$  is a double cover, so that  $\overline{C}$  is birational to a hyperelliptic curve of  $g = 3$ .

**Exercise 4.10 (Genus 6)** Let  $C$  be a curve of  $g = 6$ , and assume it has no  $g_2^1$ ,  $g_3^1$  or  $g_5^2$ . If  $D$  is a  $g_4^1$ , show that  $K - D$  has degree 6 and  $l(K - D) = 3$ . Show that  $|K - D|$  is a  $g_6^2$ , so defines a morphism  $\varphi_{K-D}: C \rightarrow \mathbb{P}^2$ .

Let  $\Gamma_6 \subset \mathbb{P}^2$  be a sextic having double points (nodes or cusps) at the 4 points  $(1, 0, 0)$ ,  $(0, 1, 0)$ ,  $(0, 0, 1)$ ,  $(1, 1, 1)$  of the standard projective frame of reference. By considering the linear system of cubics of  $\mathbb{P}^2$  passing through the 4 points, show that the resolution  $C$  has a linear system of dimension  $\geq 6$  and degree  $\leq 10$ .

Given that its resolution  $C \rightarrow \Gamma_6$  is a curve of genus 6. Show that  $C$  has 5  $g_4^1$ s and complementary  $g_6^2$ s. [Hint: Four of them are fairly obvious. The fifth comes from the pencil of conics through the 4 points.]

It is a fact that any curve of genus 6 is given either by this construction, or a different construction adapted to the case that  $C$  has a  $g_2^1$ ,  $g_3^1$  or  $g_5^2$ , or is a double cover of curve of  $g = 1$ . (The  $g_5^2$  case correspond to a plane quintic  $C_5 \subset \mathbb{P}^2$ .) Unfortunately, it would be something of a detour from the main course to discuss this rigorously or comprehensibly.